

# CANONICAL FORMS OF BOREL FUNCTIONS ON THE MILLIKEN SPACE

DISSERTATION ZUR ERLANGUNG DES DOKTORGRADES  
DER MATHEMATISCH-NATURWISSENSCHAFTLICHEN FAKULTÄT  
DER CHRISTIAN-ALBRECHTS-UNIVERSITÄT ZU KIEL

VORGELEGT VON OLAF KLEIN

KIEL, 2002

CANONICAL FORMS OF BOREL FUNCTIONS ON THE MILLIKEN SPACE  
Dissertation, 2002

**Klein, Olaf**

Christian-Albrechts-Universität zu Kiel  
Mathematisches Seminar  
Ludewig-Meyn-Straße 4  
D-24098 Kiel, Germany  
Internet: <http://www.math.uni-kiel.de>

*Fakultät:* Mathematisch-Naturwissenschaftliche Fakultät  
*Wissenschaftsfach:* Mathematik  
*Referent:* Prof. Dr. Otmar Spinas  
*Korreferent:* Prof. Dr. Peter Koepke  
*Datum der Disputation:* Montag, den 8. 7. 2002  
*Zum Druck genehmigt:* Kiel, den 17. 7. 2002

# CONTENTS

## **0 Preface**

0.1	Kurzfassung . . . . .	4
0.2	Abstract . . . . .	5
0.3	Introduction . . . . .	6

## **1 The Milliken Space . . . . . 10**

## **2 Proof of the Main Theorem . . . . . 15**

## **Appendix**

A	Corollaries . . . . .	42
B	Erklärung . . . . .	44
C	Lebenslauf . . . . .	45
D	Bibliography . . . . .	46
E	Symbols . . . . .	48
F	Index . . . . .	49

# 0. PREFACE

## 0.1 KURZFASSUNG

Das Ziel dieser Arbeit ist die Kanonisierung von Borelfunktionen  $\Delta: \Omega^\omega \rightarrow \mathbb{R}$ , wobei  $\Omega^\omega$  den Milliken-Raum bezeichnet, d.h. den Raum aller aufsteigenden unendlichen Folgen von paarweise disjunkten, nichtleeren, endlichen Teilmengen von  $\omega$ . Unser Hauptresultat bezieht sich dabei auf die metrische Topologie auf dem Milliken-Raum. Das Resultat ist eine gemeinsame Verallgemeinerung sowohl eines Satzes von Taylor (s. Satz 0.3.5) als auch eines Satzes von Prömel und Voigt (s. Satz 0.3.8).

Wir beginnen, indem wir einige Notationen einführen. Die Menge der natürlichen Zahlen bezeichnen wir mit  $\omega$ , und wir identifizieren wie üblich jede natürliche Zahl mit der Menge ihrer Vorgänger, z.B.  $k = \{0, \dots, k-1\}$ . Für die Menge aller Teilmengen von  $X$ , die die gleiche Kardinalität  $\kappa \leq \omega$  besitzen, schreiben wir  $[X]^\kappa$ . Die Menge aller endlichen Teilmengen von  $X$  bezeichnen wir mit  $[X]^{<\omega}$ . Darüber hinaus definieren wir noch  $[X]^{\leq\omega} = [X]^{<\omega} \cup [X]^\omega$ . Schließlich sei  $\Omega^{<\omega}$  noch der Raum aller aufsteigenden endlichen Folgen von paarweise disjunkten, nichtleeren, endlichen Teilmengen von  $\omega$ .

**DEFINITION.** Sei  $\gamma: \Omega^{<\omega} \rightarrow \{sm, min-sep, max-sep, min-max, sss, vss\}$ . Zu jedem  $m \in [\omega]^{<\omega}$  sei  $sm(m) = \emptyset$ ,  $min-sep(m) = \{min(m)\}$ ,  $max-sep(m) = \{max(m)\}$ ,  $min-max(m) = \{min(m), max(m)\}$  und  $sss(m) = vss(m) = m$ .

Für  $x \in \Omega^\omega$  definieren wir  $\Gamma_\gamma(x)$  wie folgt: Sei  $k(0) = 0$  und  $\langle k(i): 0 < i < N \leq \omega \rangle$  eine aufsteigende Folge derjenigen  $k$ , für die  $\gamma(x \upharpoonright (k-1)) = vss$  gilt. Darüber hinaus sei  $k(N) = \omega$ , wenn  $N < \omega$ . Damit setzen wir  $\Gamma_\gamma(x) = \langle \bigcup_{k(i) \leq j < k(i+1)} \gamma(x \upharpoonright j)(x(j)): i < N \rangle$ .

Für  $\kappa \leq \omega$  und  $a \in \Omega^\omega$  sei  $(a)^\kappa$  die Menge aller aufsteigenden Folgen von  $\kappa$  paarweise disjunkten, nichtleeren, endlichen Teilmengen von  $\omega$ , die durch Vereinigung von Stücken  $a(i)$ ,  $i \in \omega$ , entstanden sind. Jetzt können wir unser Hauptresultat angeben:

**HAUPTSATZ.** Zu jeder Borel-meßbaren Funktion  $\Delta: \Omega^\omega \rightarrow \mathbb{R}$  existiert ein  $\gamma: \Omega^{<\omega} \rightarrow \{sm, min-sep, max-sep, min-max, sss, vss\}$  und ein  $a \in \Omega^\omega$ , so daß für alle  $x, y \in (a)^\omega$  gilt

$$\Delta(x) = \Delta(y) \text{ gdw. } \Gamma_\gamma(x) = \Gamma_\gamma(y).$$

**BEMERKUNG.** Unser Hauptresultat hat sogar Gültigkeit für Abbildungen  $\Delta: \Omega^\omega \rightarrow \mathbb{R}$ , die Baire-meßbar bezüglich der H-Elmentuck-Topologie sind. Zur Definition der H-Elmentuck Topologie siehe den Beginn des ersten Kapitels. Die Bemerkung im Anschluß an Lemma 2.1 zeigt die entsprechende Änderung in

unserem Beweis auf.

Darüber hinaus gilt für das garantierte  $a \in \Omega^\omega$ , daß für keine  $x, y \in (a)^\omega$  die Menge  $\Gamma_\gamma(x)$  ein geeignetes Anfangsstück von  $\Gamma_\gamma(y)$  ist. Zur Definition von „ein geeignetes Anfangsstück“ siehe die Definition vor Lemma 2.35.

Nimmt eine gegebene Borel-meßbare Funktion nur abzählbar viele Werte an, so ist  $\Gamma_\gamma(x)$  für jede  $x \in (a)^\omega$  endlich. Die Definition von „endlich“ in diesem Zusammenhang wird direkt vor Lemma 2.36 gegeben.

Den Beweis des Hauptsatzes geben wir in Kapitel 2. In Kapitel 1 zeigen wir zunächst, daß alle analytischen Teilmengen des Milliken-Raums vollständig H-Ramsey sind – eine Eigenschaft, auf die wir in Kapitel 2 zurückgreifen werden.

## 0.2 ABSTRACT

The goal of this work is to canonize Borel measurable mappings  $\Delta: \Omega^\omega \rightarrow \mathbb{R}$ , where  $\Omega^\omega$  is the Milliken space, i.e., the space of all increasing infinite sequences of pairwise disjoint nonempty finite sets of  $\omega$ . Our main result refers to the metric topology on the Milliken space. The result is a common generalization of a theorem of Taylor (cf. Theorem 0.3.5) and a theorem of Prömel and Voigt (cf. Theorem 0.3.8).

We begin by establishing some notation. The set of nonnegative integers is denoted by  $\omega$ , and we identify each element of  $\omega$  with the set of its predecessors as usual, for instance  $k = \{0, \dots, k-1\}$ . For the set of all subsets of  $X$ , which have the same cardinality  $\kappa \leq \omega$ , we write  $[X]^\kappa$ . The collection of all finite subsets of  $X$  is denoted by  $[X]^{<\omega}$ . Moreover let  $[X]^{\leq\omega} = [X]^{<\omega} \cup [X]^\omega$ . Finally, let  $\Omega^{<\omega}$  denote the space of all increasing finite sequences of pairwise disjoint nonempty finite subsets of  $\omega$ .

DEFINITION. Let  $\gamma: \Omega^{<\omega} \rightarrow \{sm, \text{min-sep}, \text{max-sep}, \text{min-max}, \text{sss}, \text{vss}\}$ . For  $m \in [\omega]^{<\omega}$  let  $sm(m) = \emptyset$ ,  $\text{min-sep}(m) = \{\text{min}(m)\}$ ,  $\text{max-sep}(m) = \{\text{max}(m)\}$ ,  $\text{min-max}(m) = \{\text{min}(m), \text{max}(m)\}$  and  $\text{sss}(m) = \text{vss}(m) = m$ .

Let  $x \in \Omega^\omega$ . Define  $\Gamma_\gamma(x)$  as follows: Let  $k(0) = 0$  and  $\langle k(i): 0 < i < N \leq \omega \rangle$  increasingly enumerate those  $k$  such that  $\gamma(x \upharpoonright (k-1)) = \text{vss}$ . Moreover let  $k(N) = \omega$ , if  $N < \omega$ . Now let  $\Gamma_\gamma(x) = \langle \bigcup_{k(i) \leq j < k(i+1)} \gamma(x \upharpoonright j)(x(j)): i < N \rangle$ .

For  $\kappa \leq \omega$  and  $a \in \Omega^\omega$  let  $(a)^\kappa$  denote the collection of all increasing sequences of  $\kappa$  pairwise disjoint nonempty finite subsets of  $\omega$ , which are obtained by unions of some  $a(i)$ ,  $i \in \omega$ . Now we give our main result:

MAIN THEOREM. (TH) *For every Borel measurable mapping  $\Delta: \Omega^\omega \rightarrow \mathbb{R}$  there exist  $\gamma: \Omega^{<\omega} \rightarrow \{sm, \text{min-sep}, \text{max-sep}, \text{min-max}, \text{sss}, \text{vss}\}$  and  $a \in \Omega^\omega$  such that for all  $x, y \in (a)^\omega$*

$$\Delta(x) = \Delta(y) \text{ iff } \Gamma_\gamma(x) = \Gamma_\gamma(y).$$

REMARK. Our main result is even valid for mappings  $\Delta: \Omega^\omega \rightarrow \mathbb{R}$ , which are Baire measurable with respect to the H-Ellentuck topology. For the definition of H-Ellentuck topology see the beginning of chapter 1. The remark subsequent to Lemma 2.1 shows the corresponding modification of our proof.

Moreover for the guaranteed  $a \in \Omega^\omega$  it holds that for no  $x, y \in (a)^\omega$  the set  $\Gamma_\gamma(x)$  is a proper initial segment of  $\Gamma_\gamma(y)$ . For the definition of being a proper initial segment see the definition before Lemma 2.35.

Finally, if we restrict to Borel measurable mappings with a countable range, we have that  $\Gamma_\gamma(x)$  is finite for every  $x \in (a)^\omega$ . The definition of  $\Gamma_\gamma(x)$  being finite is given before Lemma 2.36.

We give the proof of the Main Theorem in chapter 2. In chapter 1 we show that every analytic subset of the Milliken space is completely H-Ramsey – a property that will be used in chapter 2.

### 0.3 INTRODUCTION

Ramsey's Theorem [Ra30] is an important extension of the *pigeon-hole principle*: If  $\omega = P_0 \cup \dots \cup P_{k-1}$  is a partition of  $\omega$  into finitely many pieces, then for some  $i < k$ ,  $P_i$  is infinite. Moreover if  $f: \omega \rightarrow \omega$ , then there exists  $A \in [\omega]^\omega$  such that  $f \upharpoonright A$  is either one to one or constant.

THEOREM 0.3.1. (Ramsey R) *Let  $k, l \in \omega$ . If  $[\omega]^k = P_0 \cup \dots \cup P_{l-1}$  is a partition of  $[\omega]^k$  into finitely many pieces, there is an infinite set  $A \in [\omega]^\omega$  such that  $[A]^k \subseteq P_i$  for some  $i < l$ .*

Ramsey's Theorem can be viewed as a canonization of finite-range functions on  $[\omega]^k$ . Later P. Erdős and R. Rado [ErRa50] canonized arbitrary such functions.

THEOREM 0.3.2. (Erdős-Rado ER) *If  $k \in \omega$  and  $f: [\omega]^k \rightarrow \omega$ , then there exists an infinite set  $X \subseteq \omega$  and a set  $\Delta(f, X) \subseteq \{0, \dots, k-1\}$  such that if  $\{x_0, \dots, x_{k-1}\}$  and  $\{y_0, \dots, y_{k-1}\}$  are in  $[X]^k$  with  $x_0 < \dots < x_{k-1}$  and  $y_0 < \dots < y_{k-1}$ , then*

$$f(\{x_0, \dots, x_{k-1}\}) = f(\{y_0, \dots, y_{k-1}\}) \text{ iff } x_i = y_i \text{ for all } i \in \Delta(f, X).$$

About twenty years later N. Hindman [Hi74] analysed the space of all finite subsets of  $\omega$ . He found the following famous result, which was a conjecture of Graham and Rothschild [GrRo71]. For simplicity of notation we identify each finite subset  $X$  of  $\omega$  with  $\langle X \rangle$ .

THEOREM 0.3.3. (Hindman) *Let  $k \in \omega$  with  $k > 0$ . If  $f: [\omega]^{<\omega} \rightarrow k$ , then there exists  $a \in \Omega^\omega$  such that  $f$  is constant on  $(a)^\omega$ .*

An elegant proof of this result has been provided by J. E. Baumgartner [Ba74]. This theorem was the basis of the work of K. R. Milliken and A. D. Taylor mentioned below. Taylor proved a canonical partition relation for finite subsets of  $\omega$  that generalizes Hindman's Theorem in much the same way that the Erdős-Rado Theorem generalizes Ramsey's Theorem. In his proof Taylor used a  $n$ -dimensional version of Theorem 0.3.3 (cf. Lemma 2.2 of [Ta76]), which was obtained independently also by Milliken (cf. Theorem 2.2 of [Mi75]).

**THEOREM 0.3.4. (Milliken-Taylor MT)** *Let  $a \in \Omega^\omega$  and  $k, l \in \omega$  with  $k, l > 0$ . If  $f: (a)^k \rightarrow l$ , then there exists  $b \in (a)^\omega$  such that  $f$  is constant on  $(b)^k$ .*

The following result of Taylor [Ta76] was stimulating for a part of this work.

**THEOREM 0.3.5. (Taylor)** *If  $f: [\omega]^{<\omega} \rightarrow \omega$ , then there exists  $a \in \Omega^\omega$  such that exactly one of (a) - (e) holds:*

- (a) *If  $m, n \in (a)^1$ , then  $f(m) = f(n)$ .*
- (b) *If  $m, n \in (a)^1$ , then  $f(m) = f(n)$  iff  $\min(m) = \min(n)$ .*
- (c) *If  $m, n \in (a)^1$ , then  $f(m) = f(n)$  iff  $\max(m) = \max(n)$ .*
- (d) *If  $m, n \in (a)^1$ , then  $f(m) = f(n)$  iff  $\min(m) = \min(n)$  and  $\max(m) = \max(n)$ .*
- (e) *If  $m, n \in (a)^1$ , then  $f(m) = f(n)$  iff  $m = n$ .*

F. Galvin and K. Prikry have shown in [GaPr73] that a similar result to Theorem 0.3.1 is valid for finite partitions of  $[\omega]^\omega$  - with the restriction that all pieces of the partitions must be Borel.

**THEOREM 0.3.6. (Galvin-Prikry GP)** *Let  $k \in \omega$  with  $k > 0$  and  $[\omega]^\omega = P_0 \cup \dots \cup P_{k-1}$  a partition of  $[\omega]^\omega$  into finitely many pieces, where each  $P_i$  is Borel. Then there is an infinite set  $A \in [\omega]^\omega$  and  $i < k$  with  $[A]^\omega \subseteq P_i$ .*

The power set of  $\omega$  can be identified with the *Cantor space*  $2^\omega$ . It can be endowed with the product topology of the discrete topology on  $\omega$ . It is a well-known fact that this topological space is completely metrizable. Thus, we can interpret the spaces  $[\omega]^k$  and  $[\omega]^\omega$  in the theorems above as topological spaces with the relative topology of  $[\omega]^{<\omega}$ . For distinction we call this topology the *metric topology* of  $[\omega]^{<\omega}$ .

A subset  $P \subseteq [\omega]^\omega$  is called *Ramsey* iff there is an infinite set  $A \in [\omega]^\omega$  such that either  $[A]^\omega \subseteq P$  or else  $[A]^\omega \cap P = \emptyset$ . By Theorem 0.3.6 every Borel set is Ramsey.

J. Silver [Si70] extended the result of Galvin-Prikry to analytic sets. Subsequent to Silver's investigation A. Mathias [Ma68] obtained a new proof of the same result.

For stronger results E. Ellentuck has introduced a finer topology on  $[\omega]^\omega$  which

is called *Ellentuck topology*. If  $A \in [\omega]^{<\omega}$  and  $B \in [\omega]^{\leq\omega}$ , then we write  $A \triangleleft B$  iff  $\max(A) < \min(B)$ , whenever both  $A$  and  $B$  are nonempty. For any  $a \in [\omega]^{<\omega}$  and  $A \in [\omega]^\omega$  with  $a \triangleleft A$  let  $[a, A]^\omega = \{S \in [\omega]^\omega : a \subseteq S \subseteq a \cup A\}$ . The Ellentuck topology then has as basic open sets all the sets of the form  $[a, A]^\omega$  for  $a \triangleleft A$ . Note that there are continuum many pairwise disjoint ones of them. Clearly, the Ellentuck topology is finer than the metric topology.

Call a set  $P \subseteq [\omega]^\omega$  *completely Ramsey* iff for every  $a \triangleleft A$  there is  $B \in [A]^\omega$  with  $[a, B]^\omega \subseteq P$  or  $[a, B]^\omega \cap P = \emptyset$ . Ellentuck [El74] has shown the following main result, which is slightly stronger than the theorem of Galvin-Prikry.

**THEOREM 0.3.7. (Ellentuck)** *Let  $P \subseteq [\omega]^\omega$ . Then  $P$  is completely Ramsey, if  $P$  has the Baire property in the Ellentuck topology.*

Moreover Galvin [El74] made the observation, that every completely Ramsey set has the Baire property. Therewith also the converse of Theorem 0.3.7 holds. An analogous result with respect to a finer topology – the  $\Sigma$ -topology – was proven by Milliken (see Theorem 4.4 of [Mi75]). Especially, we take notice of a corollary of Milliken's result: Let  $k > 0$  and  $\Omega^\omega = P_0 \cup \dots \cup P_{k-1}$  a partition of  $\Omega^\omega$  into finitely many pieces, where each  $P_i$  is Borel. Then there exists  $a \in \Omega^\omega$  and  $i < k$  with  $(a)^\omega \subseteq P_i$ . We denote it by „M“.

P. Pudlák and V. Rödl [PuRö82] canonized Borel-measurable mappings on  $[\omega]^\omega$  with a countable range. The following result of H. J. Prömel and B. Voigt [PrVo85] gives the canonization of such functions with arbitrary range.


**THEOREM 0.3.8. (Prömel-Voigt PV)** *Let  $\Delta: [\omega]^\omega \rightarrow \mathbb{R}$  be a Borel-measurable mapping. Then there exists  $A \in [\omega]^\omega$  and there exists  $\gamma: [A]^{<\omega} \rightarrow \{s, m\}$  such that the mapping  $\Gamma: [A]^\omega \rightarrow [A]^{\leq\omega}$  with  $\Gamma(X) = \{k \in X : \gamma(X \cap k) = s\}$  has the following properties:*

- (a)  $\Gamma(X) \subseteq X$  for all  $X \in [A]^\omega$ ,
- (b) for no  $X, Y \in [A]^\omega$  there exists  $k \in \Gamma(Y)$  such that  $\Gamma(X) = \Gamma(Y) \cap k$ , i.e., no  $\Gamma(X)$  is a proper initial segment of some  $\Gamma(Y)$ ,
- (c) for all  $X, Y \in [A]^\omega$  it follows that  $\Delta(X) = \Delta(Y)$  iff  $\Gamma(X) = \Gamma(Y)$ .

If we restrict to Borel-measurable mappings with a countable range, then we can find  $B \in [A]^\omega$  such that each  $\Gamma(X)$  will be finite (cf. Corollary 1 of [PrVo85]). This is slightly stronger than the corresponding result of Pudlák and Rödl.

The following figure shows the relation between some theorems mentioned above. Here  $A \rightarrow B$  means that  $A$  generalizes  $B$ .





All of these implications are pretty obvious and well-known. It was natural to search for a theorem, which stands at the place of the interrogation sign  $?_1$ . The purpose of this work is to provide such a theorem. It has  $?_2$  as a corollary – an finite-dimensional version of Theorem 0.3.5 of Taylor (cf. section A in the appendix). For the implication  $TH \rightarrow PV$  see section A in the appendix. The implication  $TH \rightarrow M$  is obvious.

# 1. THE MILLIKEN SPACE

Hindman's Theorem can be stated in equivalent form speaking about integers and their sums rather than finite sets and their unions. The sum of two integers written in binary notation looks like the characteristic function of the union of two sets, provided the integers in binary are sufficiently spread out so that no carrying occurs upon addition. But the proof of Hindman's Theorem shows that the integers can be chosen with such a property. Milliken [Mi75] stated and proved his results in the sum notation.

The following results up to 1.8 are essentially Milliken's results in the finite set notation. Also see [To98] for an axiomatic treatment of these arguments.

First of all let us expand our notation.

DEFINITION. For all  $\kappa \leq \omega$  let  $\Omega^\kappa$  denote the collection of all mappings  $f: \kappa \rightarrow [\omega]^{<\omega}$  such that  $f(i)$  is nonempty for every  $i \in \kappa$  and  $f(i) \triangleleft f(j)$  for all  $i < j < \kappa$ . Additionally, for all  $\kappa \leq \omega$  let  $\Omega^{<\kappa} = \bigcup_{i \in \kappa} \Omega^i$  and  $\Omega^{\leq \kappa} = \Omega^{<\kappa} \cup \Omega^\kappa$ .

Moreover we define  $\omega_{max}$  to be the mapping  $\omega \rightarrow [\omega]^{<\omega}$  with  $i \mapsto \{i\}$  for every  $i \in \omega$ .

If  $s$  is a mapping, we will write  $dom(s)$  to denote the domain of  $s$  and  $ran(s)$  to denote the range of  $s$ .

DEFINITION. If  $a \in \Omega^\omega$  and  $\kappa \leq \omega$ , let  $(a)^\kappa$  denote the set of all mappings  $f \in \Omega^\kappa$  such that for every  $i \in \kappa$  there exists an  $A \in [\omega]^{<\omega}$  with  $f(i) = \bigcup_{j \in A} a(j)$ . Moreover for all  $\kappa \leq \omega$  let  $(a)^{<\kappa} = \bigcup_{i \in \kappa} (a)^i$  and  $(a)^{\leq \kappa} = (a)^{<\kappa} \cup (a)^\kappa$ .

Finally, if  $s \in \Omega^{<\omega}$  and  $a \in \Omega^\omega$ , we use  $(s, a)^\omega$  to denote the set of mappings  $x \in \Omega^\omega$  such that  $x(i) = s(i)$  for every  $i \in dom(s)$  and for some  $b \in (a)^\omega$ ,  $x(i + dom(s)) = b(i)$  for every  $i \in \omega$ .

Assume that  $s, t \in \Omega^{<\omega}$  and  $a, b \in \Omega^\omega$ . We abbreviate  $s \in (t)^{<\omega}$  resp.  $s \in (b)^{<\omega}$  resp.  $a \in (b)^\omega$  as  $s \ll t$  resp.  $s \ll b$  resp.  $a \ll b$ .

Now let  $s \in \Omega^{<\omega}$  and  $t \in \Omega^{\leq \omega}$ . We write  $s < t$  iff  $s(dom(s) - 1) \triangleleft t(0)$ , whenever  $s$  and  $t$  are nonempty. If  $s < t$ , then we use  $s \hat{\ } t$  to denote the mapping  $\langle s(i): i < dom(s), t(i): i < dom(t) \rangle$ . Moreover for every  $k \in \omega$  let  $t \upharpoonright k$  denote the mapping  $\langle t(i): i < k \rangle$  and  $t \upharpoonright k$  denote the mapping  $\langle t(i): i \geq k \rangle$ .

Note that  $(a)^\omega$  and  $[a]^\omega$ ,  $t \upharpoonright k$  and  $t \upharpoonright k$  as well as  $s \ll t$  and  $s < t$  have different meanings. Regard  $\Omega^\omega$  as a topological space endowed with the neighborhood system consisting of sets of the form  $(s, a)^\omega$ , where  $s \in \Omega^{<\omega}$  and  $a \in \Omega^\omega$ . We will call  $\Omega^\omega$  the *Milliken space* and its topology the *H-Ellentuck topology*. The following results will refer to this topology till we revoke it.

Finally, for simplicity of notation we want to establish some abbreviations. If  $p$  is a mapping with  $ran(p) \leq 1$ , we will write  $p$  instead of  $p(0)$  or  $\emptyset$ . For the

remainder of this work let the lower case letters  $m, n$  be elements of  $\Omega^1$ ,  $p, q$  be elements of  $\Omega^{\leq 1}$ ,  $r, s, t$  be elements of  $\Omega^{< \omega}$ ,  $a, b, c, x, y, z$  be elements of  $\Omega^\omega$  and  $i, j, k, l$  be elements of  $\omega$ . Furthermore, let indexed letters be elements of the same space as the corresponding non-indexed letters. Moreover we stipulate that, whenever we write a concatenation like  $s \hat{\ } m$ , we have  $s < m$ .

In the remainder of this section we want to consider sets with an important property.

DEFINITION. We call  $R \subseteq \Omega^\omega$  *H-Ramsey* iff there is  $a$  such that  $(a)^\omega \subseteq R$  or  $(a)^\omega \subseteq \Omega^\omega \setminus R$ . Moreover we call  $R \subseteq \Omega^\omega$  *completely H-Ramsey* iff for every  $s$  and  $a$  there is  $b \ll a$  such that  $(s, b)^\omega \subseteq R$  or  $(s, b)^\omega \subseteq \Omega^\omega \setminus R$ .

First, we state some simple properties of those subsets, which are straightforward implications of the definitions.

LEMMA 1.1. *Every completely H-Ramsey set is H-Ramsey. Moreover the complement of a completely H-Ramsey set is completely H-Ramsey.*

We shall eventually be able to characterize the completely H-Ramsey subsets of the Milliken space as those with the Baire property. Our proofs depend heavily on ideas of Galvin and Prikry [GaPr73] and of Ellentuck [El74]. We start by proving that open sets are completely H-Ramsey.

LEMMA 1.2. *Every open set  $R \subseteq \Omega^\omega$  is completely H-Ramsey.*

PROOF. In order to prove the assertion in the lemma, we first give a definition, which goes back to Galvin and Prikry.

DEFINITION. We say  $a$  *accepts*  $s$  iff  $(s, a)^\omega \subseteq R$  and  $a$  *rejects*  $s$  iff there is no  $b \ll a$  which accepts  $s$ . Moreover we say  $a$  *decides*  $s$  iff  $a$  accepts  $s$  or  $a$  rejects  $s$ .

By the definition above it is obvious that for every  $s$  and  $a$  there exists  $b \ll a$  which decides  $s$ . The following claim improves this statement.

CLAIM 1.2.1. *For every  $s$  and  $a$  there exists  $b \ll a$  such that  $b$  decides  $s \hat{\ } t$  for every  $t \ll b$ .*

PROOF. Inductively, we construct  $b_j \in \Omega^\omega$  for every  $j < \omega$ . By the statement above there is  $b_0 \ll a$  such that  $b_0$  decides  $s$ . Assume that  $b_0, \dots, b_j$  have been constructed with the property that for every  $i \leq j$  and all  $t \ll \langle b_k(0) : k < i \rangle$  the set  $b_i$  decides  $s \hat{\ } t$ . Some applications of the statement above yield  $b_{j+1} \ll b_j \upharpoonright 1$  such that the inductive assumption is also satisfied for  $b_0, \dots, b_{j+1}$ .

Then  $b = \langle b_j(0) : j \in \omega \rangle$  has the desired property.  $\dashv$

For every  $s$  and  $a$  Claim 1.2.1 guarantees the existence of  $b \ll a$  such that  $b$  decides  $s \hat{\ } t$  for every  $t \ll b$ . If  $b$  accepts  $s$ , then  $(s, b)^\omega \subseteq R$ , and we are done. Otherwise  $b$  rejects  $s$ .

CLAIM 1.2.2. *If  $a$  decides  $s \hat{ } t$  for every  $t \ll a$  and  $a$  rejects  $s$ , then there is  $b \ll a$  which rejects  $s \hat{ } t$  for every  $t \ll b$ .*

PROOF. Inductively, we construct  $b_j \in \Omega^\omega$  for every  $j < \omega$ . We begin by showing that there is  $b_0 \ll a$  such that  $b_0$  rejects  $s \hat{ } m$  for every  $m \ll b_0$ .

Since  $a$  decides  $s \hat{ } t$  for every  $t \ll a$ , we can define a mapping  $d: (a)^1 \mapsto \{acc, rej\}$  by  $d(m) = acc$  iff  $a$  accepts  $s \hat{ } m$ . By using  $f(\{x_0, \dots, x_k\}) = d(\langle a(x_0) \cup \dots \cup a(x_k) \rangle)$  Theorem 0.3.4 of Milliken-Taylor guarantees the existence of  $b_0 \ll a$  such that  $d$  is constant on  $(b_0)^1$ . Now assume that  $b_0$  accepts  $s \hat{ } m$  for every  $m \ll b_0$ . Thus,  $(s \hat{ } m, b_0)^\omega \subseteq R$  for every  $m \ll b_0$ . But then  $(s, b_0)^\omega \subseteq R$ , as  $(s, b_0)^\omega = \bigcup \{(s \hat{ } m, b_0)^\omega : m \ll b_0\}$ , contradicting that  $a$  rejects  $s$ .

For the inductive step, using the same arguments repeatedly we can construct  $b_j \ll b_{j-1} \upharpoonright 1$  such that  $b_j$  rejects  $s \hat{ } t$  for every  $t \ll b_j$  with  $dom(t) = j + 1$ . The assertion follows by putting  $b = \langle b_j(0) : j \in \omega \rangle$ .  $\dashv$

By Claim 1.2.2 we can find  $c \ll b$  such that  $c$  rejects  $s \hat{ } t$  for all  $t \ll c$ . Suppose that  $x \in (s, c)^\omega \cap R$ . By definition of topology we have that  $(s, c)^\omega \cap R$  is open, as both  $(s, c)^\omega$  and  $R$  are open. Since every open set is a union of basic open sets and  $x \in (s, c)^\omega \cap R$ , there must be a basic open set within  $(s, c)^\omega \cap R$ , which contains  $x$ . Hence there exist  $k, l \in \omega$  with  $k \leq l$  such that  $s = x \upharpoonright k$  and  $(x \upharpoonright l, x \upharpoonright l)^\omega \subseteq R$ . Therefore  $x \upharpoonright l$  accepts  $x \upharpoonright l$ , which contradicts that  $c$  rejects  $x \upharpoonright l$ . Thus,  $(s, c)^\omega \subseteq R \setminus \Omega^\omega$ . This completes the proof of the lemma.  $\dashv$

DEFINITION. We call  $R \subseteq \Omega^\omega$  *H-Ramsey null* iff for every  $s$  and  $a$  there is  $b \ll a$  such that  $(s, b)^\omega \subseteq \Omega^\omega \setminus R$ .

Clearly, every H-Ramsey null set is completely H-Ramsey. Recall that a set is *nowhere dense* iff its closure contains no nonempty open set. Moreover a subset of the Milliken space is *meager* iff it is a countable union of nowhere dense sets. We shall be able to show that a set is H-Ramsey null iff it is meager iff it is nowhere dense.

LEMMA 1.3. *If  $N \subseteq \Omega^\omega$  is nowhere dense, then it is H-Ramsey null.*

PROOF. Recall that the *closure*  $Cls(N)$  of a set  $N$  is the smallest closed set containing  $N$ . By Lemma 1.1 and Lemma 1.2 we have that  $Cls(N)$  is completely H-Ramsey. Thus, for every  $s$  and  $a$  there is  $b \ll a$  such that  $(s, b)^\omega \subseteq Cls(N)$  or  $(s, b)^\omega \subseteq \Omega^\omega \setminus Cls(N) \subseteq \Omega^\omega \setminus N$ . By the definition of nowhere dense sets the former case cannot occur.  $\dashv$

LEMMA 1.4. *If  $M \subseteq \Omega^\omega$  is meager, then it is H-Ramsey null.*

PROOF. Let  $(N_k)_{k \in \omega}$  be a sequence of nowhere dense sets whose union is  $M$ . We may assume that  $N_k \subseteq N_{k+1}$  holds for all  $k$ .

Inductively, we construct  $b_j \in \Omega^\omega$  for every  $j < \omega$ . For any  $s$  and  $a$ , by Lemma 1.3 we can get  $b_0 \ll a$  such that  $(s, b_0)^\omega \subseteq \Omega^\omega \setminus N_0$ . Assume that  $b_0, \dots, b_j$  have

been constructed such that for all  $i \leq j$  we have  $(s \wedge t, b_i)^\omega \subseteq \Omega^\omega \setminus N_i$  for every  $t \ll \langle b_k(0) : k < i \rangle$ . Then some applications of Lemma 1.3 yield  $b_{j+1} \ll b_j \uparrow 1$  such that the inductive assumption is also satisfied for  $b_0, \dots, b_{j+1}$ .

Hence  $b = \langle b_j(0) : j \in \omega \rangle$  satisfies the assertion of the lemma.  $\dashv$

LEMMA 1.5. *Every subset of  $\Omega^\omega$  is nowhere dense iff it is meager iff it is H-Ramsey null.*

PROOF. Every nowhere dense set is meager. By Lemma 1.4 every meager set is H-Ramsey null. Therefore it remains to show that every H-Ramsey null set is nowhere dense.

For that purpose let  $R \subseteq \Omega^\omega$  be H-Ramsey null and  $(s, a)^\omega$  be an basic open set. By definition of H-Ramsey null there exists  $b \ll a$  such that  $(s, b)^\omega$  is outside  $R$ . Since  $\Omega^\omega \setminus Cls(R)$  is the union of all open sets outside  $R$ , the set  $(s, b)^\omega$  is also outside  $Cls(R)$ .

Hence there is no open set within  $Cls(R)$ , so  $R$  is nowhere dense.  $\dashv$

Now we prove our main result concerning completely H-Ramsey sets with respect to the H-Ellentuck topology. Recall that a set has the *Baire property* iff it can be expressed as the symmetric difference of an open and a meager set.

LEMMA 1.6. *Every set  $R \subseteq \Omega^\omega$  with the Baire property is completely H-Ramsey.*

PROOF. By definition of the Baire property  $R$  can be expressed as  $R = U \Delta M$  where  $U$  is open,  $M$  is meager and  $\Delta$  is the symmetric difference. Since  $M$  is meager, by Lemma 1.4 for any  $s$  and  $a$  we can choose  $b \ll a$  so that  $(s, b)^\omega \subseteq \Omega^\omega \setminus M$ . Moreover by Lemma 1.2 there is  $c \ll b$  such that  $(s, c)^\omega \subseteq U$  or  $(s, c)^\omega \subseteq \Omega^\omega \setminus U$ . In the former case we have  $(s, c)^\omega \subseteq R$ , and in the latter case we have  $(s, c)^\omega \subseteq \Omega^\omega \setminus R$ .  $\dashv$

MAIN LEMMA 1.7. *Let  $R \subseteq \Omega^\omega$ . Then  $R$  is completely H-Ramsey iff  $R$  has the Baire property.*

PROOF. Let  $R$  be completely H-Ramsey. Then we claim that  $N = R \setminus Int(R)$  is nowhere dense, where  $Int(R)$  denotes the *interior* of  $R$ , i.e., the union of all open subsets of  $R$ .

Indeed, if this fails, there are  $s$  and  $a$  such that  $(s, a)^\omega \subseteq Cls(N)$ . Since  $C$  is completely H-Ramsey, we can choose  $b \ll a$  such that  $(s, b)^\omega \subseteq R$  or  $(s, b)^\omega \subseteq \Omega^\omega \setminus R$ . We have that  $(s, b)^\omega \cap N \neq \emptyset$ , as  $(s, b)^\omega \subseteq Cls(N)$  and  $(s, a)^\omega \subseteq Cls(N) \setminus N$  would contradict that  $\Omega^\omega \setminus Cls(N)$  is the union of all open sets outside  $N$ . Therefore  $(s, b)^\omega \subseteq \Omega^\omega \setminus R$  is impossible, so that  $(s, b)^\omega \subseteq R$  holds. By definition of  $Int(R)$  we get that  $(s, b)^\omega \subseteq Int(R)$  and  $(s, b)^\omega \cap N = \emptyset$ , giving a contradiction. Thus,  $R$  can be expressed as  $R = Int(R) \Delta N$ , so  $R$  has the Baire property.

Hence, the assertion of the lemma follows by Lemma 1.6.  $\dashv$

Finally, we want to relate the Ramsey properties to the *metric topology* on  $\Omega^\omega$ . The following result and also the results of the remainder of this work will refer to this topology. Note that by definition the Milliken space is a subspace of  $([\omega]^{<\omega})^\omega$ . The latter one can be regarded as a topological space with the product topology of the discrete topology of  $[\omega]^{<\omega}$ . Hence the metric topology on  $\Omega^\omega$  is the relative topology on  $([\omega]^{<\omega})^\omega$ .

The metric topology has as basic open sets the sets of the form  $(s, \omega_{max})^\omega$  with  $s \in \Omega^{<\omega}$ . Thus, it is completely metrizable and coarser than the H-Ellentuck topology. Moreover as its basis is countable, the Milliken space is separable and therefore a Polish space.

**THEOREM 1.8.** *Every analytic subset of  $\Omega^\omega$  with respect to the metric topology is completely H-Ramsey.*

**PROOF.** First, recall that a *Souslin scheme*  $(S_f)_{f \in \omega^{<\omega}}$  on  $\Omega^\omega$  is a family of subsets of  $\Omega^\omega$  that are indexed by finite sequences of nonnegative integers. The *Souslin operation* applied to such a scheme produces the set  $\bigcup_{g \in \omega^\omega} \bigcap_{k \in \omega} S_{g \upharpoonright k}$ , where  $\omega^\omega$  denotes the set of all functions mapping  $\omega$  into  $\omega$ , and  $g \upharpoonright k$  is the restriction of  $g$  to the predecessors of  $k$  (cf. p. 198 of [Ke95]).

The analytic sets are formed by the Souslin operation applied to each Souslin scheme of closed sets (cf. Corollary (25.8) of [Ke95]). We have that all closed sets in the metric topology on  $\Omega^\omega$  are also closed with respect to the H-Ellentuck topology. Moreover every closed set has the Baire property with respect to the H-Ellentuck topology. As the Souslin operation preserves the Baire property (cf. Corollary (29.14) of [Ke95]), by Main Lemma 1.7 every analytic set is completely H-Ramsey. ◻

## 2. PROOF OF THE MAIN THEOREM

The proof of the Main Theorem requires some further results. Our first lemma is analogous to Lemma 1 of [PrVo85].

LEMMA 2.1. *Let  $\Delta: \Omega^\omega \rightarrow \mathbb{R}$  be Borel-measurable. Then there exists  $a$  such that the restriction  $\Delta \upharpoonright (a)^\omega$  is a continuous mapping.*

PROOF. Let  $(I_j)_{j \in \omega}$  be an enumeration of all open intervals in  $\mathbb{R}$  which have rational endpoints. The  $I_j$  form a basis for the topology of the reals. Inductively, we construct  $a_j \in \Omega^\omega$  for every  $j < \omega$ . Put  $a_0 = \omega_{max}$  and assume by induction that  $a_0, \dots, a_j$  have been constructed such that for all  $i < j$  and all  $s \ll \langle a_0(0), \dots, a_i(0) \rangle$  either  $(s, a_{i+1})^\omega \subseteq \Delta^{-1}(I_i)$  or  $(s, a_{i+1})^\omega \subseteq \Omega^\omega \setminus \Delta^{-1}(I_i)$ . Since  $I_j$  is open, it follows that  $\Delta^{-1}(I_j) \subseteq \Omega^\omega$  must be Borel and hence by Theorem 1.8 completely H-Ramsey. Hence we can get an  $a_{j+1} \ll a_j \upharpoonright 1$  such that for all  $s \ll \langle a_0(0), \dots, a_j(0) \rangle$  either  $(s, a_{j+1})^\omega \subseteq \Delta^{-1}(I_j)$  or  $(s, a_{j+1})^\omega \subseteq \Omega^\omega \setminus \Delta^{-1}(I_j)$ .

Then  $a = \langle a_j(0): j \in \omega \rangle$  has the desired property.  $\dashv$

REMARK. Suppose  $\Delta: \Omega^\omega \rightarrow \mathbb{R}$  is Baire measurable with respect to the H-Ellentuck topology. The same argument, using Lemma 1.6 instead of Theorem 1.8, shows that  $\Delta \upharpoonright (a)^\omega$  is continuous with respect to the metric topology on  $\Omega^\omega$  for some  $a$ .

For the remainder of this section let  $\Delta: \Omega^\omega \rightarrow \mathbb{R}$  be an arbitrary but fixed mapping.

DEFINITION. Let  $s, t$  and  $x$  be such that  $s < t, x$  and  $s = \langle s(0), \dots, s(k) \rangle$ . We abbreviate the mappings  $s \upharpoonright k \wedge \langle s(k) \cup t(0) \rangle \wedge t \upharpoonright 1$  resp.  $s \upharpoonright k \wedge \langle s(k) \cup x(0) \rangle \wedge x \upharpoonright 1$  as  $s \upharpoonright \wedge t$  resp.  $s \upharpoonright \wedge x$ . Additionally, we define  $s \upharpoonright \wedge \emptyset$  to be  $s \upharpoonright$ . Moreover we use  $s \square$  as a variable for  $s$  or  $s \upharpoonright$ .

REMARK. At the beginning of chapter 1 we had defined the H-Ellentuck topology by the basis consisting of the sets  $(s, a)^\omega$  with  $s \in \Omega^{<\omega}$  and  $a \in \Omega^\omega$ . Alternatively, it had been possible to define the basis by the sets  $(s \square, a)^\omega := \{x \in \Omega^\omega: \exists y \ll a \ x = s \square \wedge y\}$  with  $s \in \Omega^{<\omega}$  and  $a \in \Omega^\omega$ . Since  $(s \upharpoonright, a)^\omega = \bigcup_{m \ll a} (s \upharpoonright \wedge m, a)^\omega$ , the topology defined by the latter sets had not been finer.

Analogously to [PrVo85] we introduce now the terms *separating* and *mixing*.

DEFINITION. We say that  $s \square$  and  $t \square$  are separated by  $a$  iff  $\Delta(s \square \wedge x) \neq \Delta(t \square \wedge y)$  for all  $x, y \ll a$  with  $s < x, y$  and  $t < x, y$ . Moreover  $s \square$  and  $t \square$  are mixed by  $a$  iff for no  $b \ll a$  the sets  $s \square$  and  $t \square$  are separated by  $b$ . Finally,  $s \square$  and  $t \square$  are decided by  $a$  iff  $s \square$  and  $t \square$  are separated or mixed by  $a$ .

We stipulate that, whenever we write a concatenation like  $s \sqcup \hat{m}$  resp.  $s \sqcup \hat{m} \sqcup \hat{n}$ , we have  $s < m$  resp.  $s < m < n$ . The following lemma is a straightforward implication of the definition above.

LEMMA 2.2. *For every  $s, t$  and  $a$ , there exists  $b \ll a$  which decides  $s \sqcup$  and  $t \sqcup$ . If  $s \sqcup$  and  $t \sqcup$  are decided by  $b$ , then they are also decided by each  $c \ll b$ , and  $c$  decides in the same way as  $b$  does.*

LEMMA 2.3. (Transitivity of mixing) *Assume that  $r \sqcup$  and  $s \sqcup$  as well as  $s \sqcup$  and  $t \sqcup$  are mixed by  $a$ . Then also  $r \sqcup$  and  $t \sqcup$  are mixed by  $a$ .*

PROOF. Assume to the contrary that there exists  $b \ll a$  which separates  $r \sqcup$  and  $t \sqcup$ . We may assume without loss of generality that  $r, s, t < b$ . Consider the set  $A = \{x \ll b : \exists y \ll b \Delta(r \sqcup \hat{y}) = \Delta(s \sqcup \hat{x})\}$ . Then  $A$  is analytic, so by Theorem 1.8  $A$  is completely H-Ramsey. By definition of completely H-Ramsey there exists  $c \ll b$  with  $(c)^\omega \subseteq A$  or  $(c)^\omega \cap A = \emptyset$ . Both cases lead to a contradiction:

Assume first that  $(c)^\omega \subseteq A$ . Then for all  $x \ll c$  there exists  $y \ll b$  such that  $\Delta(r \sqcup \hat{y}) = \Delta(s \sqcup \hat{x})$ . Since  $r \sqcup$  and  $t \sqcup$  are separated by  $b$ , it follows that  $\Delta(r \sqcup \hat{y}) \neq \Delta(t \sqcup \hat{z})$  for every  $y, z \ll b$ . Hence we get  $\Delta(s \sqcup \hat{x}) \neq \Delta(t \sqcup \hat{z})$  for all  $x, z \ll c$ , contradicting that  $s \sqcup$  and  $t \sqcup$  are mixed by  $a$ .

Otherwise if  $(c)^\omega \cap A = \emptyset$ , then  $r \sqcup$  and  $s \sqcup$  are separated by  $c$ . ⊥

LEMMA 2.4. *For every  $a$  there exists  $b \ll a$  such that for every  $s, t \ll b$  the sets  $s \sqcup$  and  $t \sqcup$  are decided by  $b$ .*

PROOF. Inductively, we construct  $b_j \in \Omega^\omega$  for every  $j < \omega$ . By Lemma 2.2 there exists  $b_0 \ll a$  such that  $\emptyset$  and  $\emptyset$  are decided by  $b_0$ . Assume that  $b_0, \dots, b_j$  have been constructed such that for every  $i \leq j$  and for all  $s, t \ll \langle b_k(0) : k < i \rangle$  the sets  $s \sqcup$  and  $t \sqcup$  are decided by  $b_i$ . Some applications of Lemma 2.2 yield  $b_{j+1} \ll b_j \uparrow 1$  such that the inductive assumption is also satisfied for  $b_0, \dots, b_{j+1}$ . Then  $b = \langle b_j(0) : j \in \omega \rangle$  has the desired properties. ⊥

The following Lemma is modeled in the image of Theorem 2.1 of [Ta76].

LEMMA 2.5. *For every  $s$  and  $a$ , there exists  $b \ll a$  such that exactly one of the following properties holds:*

- (a) *If  $m, n \ll b$ , then  $s \hat{m}$  and  $s \hat{n}$  are mixed by  $b$ .*
- (b) *If  $m, n \ll b$ , then  $s \hat{m}$  and  $s \hat{n}$  are mixed by  $b$  iff  $\min(m) = \min(n)$ .*
- (c) *If  $m, n \ll b$ , then  $s \hat{m}$  and  $s \hat{n}$  are mixed by  $b$  iff  $\max(m) = \max(n)$ .*
- (d) *If  $m, n \ll b$ , then  $s \hat{m}$  and  $s \hat{n}$  are mixed by  $b$  iff  $\min(m) = \min(n)$  and  $\max(m) = \max(n)$ .*
- (e) *If  $m, n \ll b$ , then  $s \hat{m}$  and  $s \hat{n}$  are mixed by  $b$  iff  $m = n$ .*



PROOF. Lemma 2.4 guarantees the existence of  $b_0 \ll a$  such that  $s \wedge m$  and  $s \wedge n$  are decided by  $b_0$  for every  $m, n \ll b_0$ . Let  $F$  be the set of all functions  $f$  such that  $\text{dom}(f) = 3$  and  $\text{ran}(f) \subseteq 2$ . Define  $g: (b_0)^3 \rightarrow F$  as follows:

$g(h)(0) = 0$  iff  $s \wedge \langle h(0) \cup h(1) \cup h(2) \rangle$  and  $s \wedge \langle h(0) \rangle$  are mixed by  $b_0$ .

$g(h)(1) = 0$  iff  $s \wedge \langle h(0) \cup h(1) \cup h(2) \rangle$  and  $s \wedge \langle h(2) \rangle$  are mixed by  $b_0$ .

$g(h)(2) = 0$  iff  $s \wedge \langle h(0) \cup h(1) \cup h(2) \rangle$  and  $s \wedge \langle h(0) \cup h(2) \rangle$  are mixed by  $b_0$ .

By Theorem 0.3.4 of Milliken-Taylor there exists  $b_1 \ll b_0$  and a function  $f = \langle f(0), f(1), f(2) \rangle \in F$  such that  $g((b_1)^3) = \{f\}$ . We claim first that  $f$  cannot be  $\langle 0, 0, 1 \rangle$  or  $\langle 1, 0, 1 \rangle$  or  $\langle 0, 1, 1 \rangle$ . The first two are ruled out by the observation that if  $f(1) = 0$ , then we must have  $f(2) = 0$ . Indeed, if  $f(2) \neq 0$ , then  $s \wedge \langle (b_1(0) \cup b_1(1)) \cup b_1(2) \cup b_1(3) \rangle$  and  $s \wedge \langle (b_1(0) \cup b_1(1)) \cup b_1(3) \rangle$  are separated by  $b_1$ . But since  $f(1) = 0$ , both of these are mixed with  $s \wedge \langle b_1(3) \rangle$ . By transitivity of mixing we get a contradiction. Similarly, the third one is ruled out since if  $f(0) = 0$ , then we must have  $f(2) = 0$ . This leaves five possibilities for  $f$ .

We will show that these five possibilities correspond to the five clauses (a) - (e) of this lemma. By construction we are guaranteed that exactly one case holds in the assertion.

*Case (a).*  $f = \langle 0, 0, 0 \rangle$ . Let  $b = \langle b_1(i) : i > 1 \rangle$  and  $m, n \ll b$ . Since  $f(1) = 0$ ,  $s \wedge m$  and  $s \wedge \langle b_1(0) \cup b_1(1) \cup m \rangle$  as well as  $s \wedge n$  and  $s \wedge \langle b_1(0) \cup b_1(1) \cup n \rangle$  are mixed by  $b$ . Moreover because  $f(0) = 0$ , we have that  $s \wedge \langle b_1(0) \cup b_1(1) \cup m \rangle$  and  $s \wedge \langle b_1(0) \rangle$  as well as  $s \wedge \langle b_1(0) \cup b_1(1) \cup n \rangle$  and  $s \wedge \langle b_1(0) \rangle$  are mixed by  $b$ . By transitivity of mixing it follows that  $s \wedge m$  and  $s \wedge n$  are mixed by  $b$  whenever  $m, n \ll b$ , so  $b$  satisfies clause (a) of the lemma.

*Case (b).*  $f = \langle 0, 1, 0 \rangle$ . Let  $b = \langle b_1(3i) \cup b_1(3i + 1) \cup b_1(3i + 2) : i < \omega \rangle$ . Suppose first that  $m, n \ll b$  with  $\min(m) = \min(n)$ . Then  $m = \langle b(k) \cup p \rangle$  and  $n = \langle b(k) \cup q \rangle$  for some  $k$  and some  $p, q \ll b \upharpoonright k$ . Since  $f(0) = 0$ ,  $s \wedge m$  and  $s \wedge \langle b_1(3k) \rangle$  as well as  $s \wedge n$  and  $s \wedge \langle b_1(3k) \rangle$  are mixed by  $b$ . By transitivity of mixing we obtain that  $s \wedge m$  and  $s \wedge n$  are mixed by  $b$ .

Conversely, if  $m, n \ll b$  and  $\min(m) < \min(n)$ , then  $m = \langle b(k) \cup p \rangle$  for some  $k$  and  $p \ll b \upharpoonright k$ , and  $b(k) < n$ . Thus,  $s \wedge m$  and  $s \wedge \langle b(k) \cup n \rangle$  are mixed by  $b$ , since both are mixed with  $s \wedge \langle b_1(3k) \rangle$  by virtue of the fact that  $f(0) = 0$ . But since  $f(1) = 1$ , we have that  $s \wedge \langle b(k) \cup n \rangle$  and  $s \wedge n$  are separated by  $b$ , and so – by transitivity of mixing – we must have that  $s \wedge m$  and  $s \wedge n$  are separated by  $b$ .

Thus,  $s \wedge m$  and  $s \wedge n$  are mixed by  $b$  iff  $\min(m) = \min(n)$ , so  $b$  satisfies clause (b) of the lemma.

*Case (c).*  $f = \langle 1, 0, 0 \rangle$ . Let  $b = \langle b_1(3i) \cup b_1(3i + 1) \cup b_1(3i + 2) : i < \omega \rangle$  like in case (b). If  $m, n \ll b$  and  $\max(m) = \max(n)$ , then  $m = \langle p \cup b(k) \rangle$  and  $n = \langle q \cup b(k) \rangle$  for some  $k$  and  $p, q \ll b \upharpoonright k$ . Since  $f(1) = 0$  we have that  $s \wedge m$  and  $s \wedge \langle b_1(3k + 2) \rangle$  as well as  $s \wedge n$  and  $s \wedge \langle b_1(3k + 2) \rangle$  are mixed by  $b$ . Hence  $s \wedge m$  and  $s \wedge n$  are mixed by  $b$ .

Conversely, if  $m, n \ll b$  and  $\max(m) < \max(n)$ , then  $n = \langle q \cup b(k) \rangle$  for some  $k$  and  $q \ll b \upharpoonright k$ , and  $m < b(k)$ . Thus,  $s \wedge n$  and  $s \wedge \langle m \cup b(k) \rangle$  are mixed by  $b$ , because both are mixed with  $s \wedge \langle b_1(3k + 2) \rangle$  since  $f(1) = 0$ . But  $s \wedge \langle m \cup b(k) \rangle$  and  $s \wedge m$  are separated by  $b$  since  $f(0) = 1$ . So we must have that  $s \wedge m$  and  $s \wedge n$  are separated by  $b$ .

Hence  $s \wedge m$  and  $s \wedge n$  are mixed by  $b$  iff  $\max(m) = \max(n)$ , and hence  $b$  satisfies clause (c) of the lemma.

CLAIM 2.5.1. *Let  $s$  and  $a$  be such that  $s \wedge m$  and  $s \wedge n$  are decided by  $a$  for every  $m, n \ll a$ . If  $s \wedge \langle h(0) \cup h(1) \cup h(2) \rangle$  and  $s \wedge \langle h(0) \rangle$  are separated by  $a$  for all  $h \in (a)^3$ , then there exists  $b \ll a$  such that  $s \wedge m$  and  $s \wedge n$  are separated by  $b$  for every  $m, n \ll b$  with  $\max(m) < \max(n)$ .*

PROOF. Let  $a_0, a_1$  be elements of  $\Omega^\omega$ . We construct  $b$  inductively. Put  $b(0) = a(0)$  and suppose that  $b(0), \dots, b(k-1)$  have been constructed such that  $s \wedge m$  and  $s \wedge n$  are separated by  $a$  for all  $m, n \ll b \upharpoonright k$  with  $\max(m) < \max(n)$ . Let  $a_0 \ll a$  with  $b(k) < a_0(0)$ . Choose  $\langle b(k) \rangle \ll a_0$  such that  $s \wedge m$  and  $s \wedge \langle p \cup b(k) \rangle$  are separated by  $a$  for every  $m, p \ll b \upharpoonright k$ . This is possible, since otherwise for all  $\langle b(k) \rangle \ll a_0$  there would exist  $m, p \ll b \upharpoonright k$  such that  $s \wedge m$  and  $s \wedge \langle p \cup b(k) \rangle$  are mixed by  $a$ . Theorem 0.3.4 would yield  $a_1 \ll a_0$  and fixed  $m, p \ll b \upharpoonright k$  such that  $s \wedge m$  and  $s \wedge \langle p \cup b(k) \rangle$  are mixed by  $a$  for every  $\langle b(k) \rangle \ll a_1$ . By transitivity of mixing (Lemma 2.3) we get that  $s \wedge \langle p \cup m \rangle$  and  $s \wedge \langle p \cup n \rangle$  are mixed by  $a$  for all  $m, n \ll a_1$ . Choosing  $h = \langle p \cup a_1(0), a_1(1), a_1(2) \rangle$  we get a contradiction to the assumption of the lemma. This completes the construction of  $b$ .  $\dashv$

Case (d).  $f = \langle 1, 1, 0 \rangle$ . To handle case (d) we choose  $b_2 \ll b_1$  as guaranteed to exist by Claim 2.5.1. Let  $b = \langle b_1(3i) \cup b_1(3i+1) \cup b_1(3i+2) : i < \omega \rangle$ . We claim that if  $m, n \ll b$ , then  $s \wedge m$  and  $s \wedge n$  are mixed by  $b$  iff  $\min(m) = \min(n)$  and  $\max(m) = \max(n)$ .

Suppose first that  $\min(m) = \min(n)$  and  $\max(m) = \max(n)$ . Then for some  $i < j$  we have that  $s \wedge m$  and  $s \wedge \langle b_2(3i) \cup b_2(3j+2) \rangle$  as well as  $s \wedge n$  and  $s \wedge \langle b_2(3i) \cup b_2(3j+2) \rangle$  are mixed by  $b$ , since  $f(2) = 0$ . By transitivity of mixing we get that  $s \wedge m$  and  $s \wedge n$  are mixed by  $b$ .

For the converse, suppose that either  $\min(m) \neq \min(n)$  or  $\max(m) \neq \max(n)$ . If  $\max(m) \neq \max(n)$ , then clearly  $s \wedge m$  and  $s \wedge n$  are separated by  $b$ , by construction according to Claim 2.5.1. Hence we can assume that  $\max(m) = \max(n)$  and  $\min(m) < \min(n)$ . Let  $m = \langle b(k) \cup p \cup b(l) \rangle$  for some  $k < l$  and some  $p \ll \langle b(i) : k < i < l \rangle$  and  $b(k) < n$ . But then  $s \wedge m$  and  $s \wedge \langle b_2(3k) \cup b_2(3l+2) \rangle$  as well as  $s \wedge \langle b_2(3k) \cup b_2(3l+2) \rangle$  and  $s \wedge \langle b(k) \cup n \rangle$  are mixed by  $b$ , since  $f(2) = 0$ . However, since  $f(1) = 1$  we have that  $s \wedge \langle b(k) \cup n \rangle$  and  $s \wedge n$  are separated by  $b$ , and by the transitivity of mixing it follows that  $s \wedge m$  and  $s \wedge n$  are separated by  $b$ . Thus, we have shown that  $b$  satisfies clause (d) of the lemma.

DEFINITION. For some given  $s$  and  $a$  we will say that  $t$  and  $b$  are *compatible* iff  $t < b$  and  $s \wedge \langle p \cup m \rangle$  and  $s \wedge \langle q \cup m \rangle$  are separated by  $a$  for every  $m \ll b$  and for all  $p, q \ll t$  with  $\max(p) \neq \max(q)$ . Note that  $p$  and  $q$  can be empty as agreed in

the introduction. We will say that  $t$  and  $b$  are *very compatible* iff they are compatible and, moreover, there exists  $n \ll b$  and there exists  $c \ll b$  such that  $t \wedge n$  and  $c$  are compatible.

CLAIM 2.5.2. *Let  $s$  and  $a$  be such that  $s \wedge m$  and  $s \wedge n$  are decided by  $a$  for every  $m, n \ll a$ . Suppose that  $s \wedge \langle h(0) \cup h(1) \cup h(2) \rangle$  and  $s \wedge \langle h(0) \cup h(2) \rangle$  are separated by  $a$  for all  $h \in (a)^3$ . Then if  $t$  and  $b$  are compatible where  $b, t \ll a$ , then  $t$  and  $b$  are in fact very compatible.*

PROOF. Suppose that  $t$  and  $b$  are compatible but not very compatible. Then for every  $m \ll b$  and for all  $c \ll b$  with  $m < c$  there exists  $n \ll c$  and there exists  $p, q \ll t \wedge m$  such that  $\max(p) \neq \max(q)$  and  $s \wedge \langle p \cup n \rangle$  and  $s \wedge \langle q \cup n \rangle$  are mixed by  $a$ . Notice that we cannot have both  $p, q \ll t$  since  $t$  and  $b$  are compatible. Thus, we better use instead of any such  $q$  a mapping of the form  $\langle q \cup m \rangle$  with the restriction  $q \ll t$ . Now two applications of Theorem 0.3.4 yield  $c \ll b$  and fixed  $p, q \ll t$  such that  $s \wedge \langle p \cup n \rangle$  and  $s \wedge \langle q \cup m \cup n \rangle$  are mixed by  $a$  for every  $m \wedge n \ll c$ . We get mixing for all  $m \wedge n$  because of our assumption above. Choosing  $h \in (c)^3$  we obtain that  $s \wedge \langle p \cup h(2) \rangle$  and  $s \wedge \langle q \cup h(0) \cup h(2) \rangle$  as well as  $s \wedge \langle p \cup h(2) \rangle$  and  $s \wedge \langle q \cup h(0) \cup h(1) \cup h(2) \rangle$  are mixed by  $a$  since  $\langle h(0), h(2) \rangle, \langle h(0) \cup h(1), h(2) \rangle \ll c$ . Thus, by transitivity of mixing  $s \wedge \langle q \cup h(0) \cup h(1) \cup h(2) \rangle$  and  $s \wedge \langle q \cup h(0) \cup h(2) \rangle$  are mixed by  $a$ , contradicting the condition imposed in the lemma. This completes the proof of the claim.  $\dashv$

Case (e).  $f = \langle 1, 1, 1 \rangle$ . To handle case (e) we construct  $b_2 \ll b_1$  inductively. To this end we build a sequence  $\{(b_2(i), c_i) : i < \omega\}$  such that  $b_2 \upharpoonright (i+1)$  and  $c_i$  are compatible for every  $i < \omega$  with  $c_i \in \Omega^\omega$ . Let  $b_2(0) = b_1(0) \cup b_1(1)$  and  $c_0 = b_1 \upharpoonright 2$ . Notice that  $b_2 \upharpoonright 1$  and  $c_0$  are compatible since  $f(1) = 1$  and  $f(2) = 1$ . Suppose now that  $b_2 \upharpoonright (k+1)$  and  $c_k$  have been constructed and are compatible. Since  $f(2) = 1$ , Claim 2.5.2 applies and hence we have that  $b_2 \upharpoonright (k+1)$  and  $c_k$  are very compatible. Thus, there exists  $\langle b_2(k+1) \rangle \ll c_k$  and there exists  $c_{k+1} \ll c_k$  such that  $b_2 \upharpoonright (k+2)$  and  $c_{k+1}$  are compatible. This completes the construction.

Now we claim that if  $m, n \ll b_2$  with  $m \neq n$  and  $\max(m) = \max(n)$ , then we have that  $s \wedge m$  and  $s \wedge n$  are separated by  $b_2$ . To see this, let  $b_2(k)$  be the last piece of  $b_2$  occurring in  $(m \cup n) \setminus (m \cap n)$ . Then we can assume without loss of generality that  $m = \langle p \cup b_2(k) \cup m_0 \rangle$  and  $n = \langle q \cup m_0 \rangle$  for some  $p, q \ll b_2 \upharpoonright k$  and some  $m_0 \ll b_2$  with  $b_2(k) < m_0$ . Since  $b_2 \upharpoonright (k+1)$  and  $c_k$  are compatible,  $m_0 \ll c_k$  and  $\max(p \cup b_2(k)) \neq \max(q)$  we have that  $s \wedge \langle p \cup b_2(k) \cup m_0 \rangle$  and  $s \wedge \langle q \cup m_0 \rangle$  are separated by  $b_2$ . Thus,  $s \wedge m$  and  $s \wedge n$  are separated by  $b_2$ . Since  $f(0) = 1$  and  $b_2 \ll b_1$ , Claim 2.5.1 applies and we can choose  $b \ll b_2$  such that  $s \wedge m$  and  $s \wedge n$  are separated by  $b$  whenever  $m, n \ll b$  and  $\max(m) < \max(n)$ .

Finally, notice that if  $s \wedge m$  and  $s \wedge n$  are separated by  $b$ , we must have  $m \neq n$  by definition of separated. So we can conclude that  $s \wedge m$  and  $s \wedge n$  are mixed by  $b$  iff  $m \neq n$ .

This completes the proof of case (e) and with it, the proof of Lemma 2.5.  $\dashv$

The following definition is based on the five cases of Lemma 2.5.

DEFINITION. We say that  $s \sqsupseteq$  is *strongly mixed by  $a$*  iff  $s \sqsupseteq \wedge m$  and  $s \sqsupseteq \wedge n$  are mixed by  $a$  for every  $m, n \ll a$ . Moreover  $s$  is *min-separated by  $a$*  iff for every  $m, n \ll a$  the sets  $s \wedge m$  and  $s \wedge n$  are mixed by  $a$  iff  $\min(m) = \min(n)$ . Furthermore,  $s \sqsupseteq$  is *max-separated by  $a$*  iff for every  $m, n \ll a$  the sets  $s \sqsupseteq \wedge m$  and  $s \sqsupseteq \wedge n$  are mixed by  $a$  iff  $\max(m) = \max(n)$ . Moreover we say that  $s$  is *min-max-separated by  $a$*  iff for every  $m, n \ll a$  the sets  $s \wedge m$  and  $s \wedge n$  are mixed by  $a$  iff  $\min(m) = \min(n)$  and  $\max(m) = \max(n)$ . Finally,  $s \sqsupseteq$  is *strongly separated by  $a$*  iff for every  $m, n \ll a$  the sets  $s \sqsupseteq \wedge m$  and  $s \sqsupseteq \wedge n$  are mixed by  $a$  iff  $m = n$ .

Furthermore, we say  $s \sqsupseteq$  is *separated in some sense by  $a$*  iff  $s \sqsupseteq$  is min-separated, max-separated, min-max-separated or strongly separated by  $a$ . Moreover  $s$  is *completely decided by  $a$*  iff  $s$  is strongly mixed by  $a$  or  $s$  is separated in some sense by  $a$ .

LEMMA 2.6. For every  $s$  and  $a$  the following properties hold.

- (a) Let  $s$  be strongly mixed by  $a$ . Then  $s \wedge m \uparrow$  is strongly mixed by  $a$  for every  $m \ll a$ .
- (b) Let  $s$  be min-separated by  $a$ . Then  $s \wedge m \uparrow$  is strongly mixed by  $a$  for every  $m \ll a$ .
- (c) Let  $s$  be max-separated by  $a$ . Then  $s \wedge m \uparrow$  is max-separated by  $a$  for every  $m \ll a$ .
- (d) Let  $s$  be min-max-separated by  $a$ . Then  $s \wedge m \uparrow$  is max-separated by  $a$  for every  $m \ll a$ .
- (e) Let  $s$  be strongly separated by  $a$ . Then  $s \wedge m \uparrow$  is strongly separated by  $a$  for every  $m \ll a$ .

PROOF. Obvious from the definition. -1

LEMMA 2.7. For every  $s$  and  $a$  the following properties hold.

- (a) Let  $s \sqsupseteq$  be strongly mixed by  $a$ . Then  $s \sqsupseteq$  and  $s \sqsupseteq \wedge m \sqsupseteq$  as well as  $s \sqsupseteq \wedge m \sqsupseteq$  and  $s \sqsupseteq \wedge n \sqsupseteq$  are mixed by  $a$  for every  $m, n \ll a$ .
- (b) Let  $s$  be min-separated by  $a$ . Then  $s \wedge m \sqsupseteq$  and  $s \wedge n \sqsupseteq$  are mixed by  $a$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$ .
- (c) Let  $s \sqsupseteq$  be max-separated by  $a$ . Then  $s \sqsupseteq$  and  $s \sqsupseteq \wedge m \uparrow$  as well as  $s \sqsupseteq \wedge m \uparrow$  and  $s \sqsupseteq \wedge n \uparrow$  are mixed by  $a$  for every  $m, n \ll a$ .
- (d) Let  $s$  be min-max-separated by  $a$ . Then  $s \wedge m \uparrow$  and  $s \wedge n \uparrow$  are mixed by  $a$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$ .

PROOF. Case (a). Let  $s \sqsupseteq$  be strongly mixed by  $a$ . First, we prove that  $s \sqsupseteq$  and  $s \sqsupseteq$

$\wedge m \square$  are mixed by  $a$  for every  $m \ll a$ . Assume to the contrary that  $s \square$  and  $s \square \wedge m$  resp.  $s \square$  and  $s \square \wedge m \uparrow$  are not mixed by  $a$  for some  $m \ll a$ . Hence there exists  $b \ll a$  such that  $s \square$  and  $s \square \wedge m$  resp.  $s \square$  and  $s \square \wedge m \uparrow$  are separated by  $b$ . Since  $s \square$  is strongly mixed by  $a$ , by Lemma 2.2 we get that  $s \square \wedge m$  and  $s \square \wedge n$  are also mixed by  $b$  for every  $m, n \ll a$ .

Now choose  $k$  minimal such that  $m < b(k)$ . By definition of separation we must have that  $s \square \wedge \langle b(k) \rangle$  and  $s \square \wedge m$  resp.  $s \square \wedge \langle b(k) \rangle$  and  $s \square \wedge m \uparrow \wedge \langle b(k) \rangle$  are separated by  $b$ . However, since  $\langle b(k) \rangle \ll a$ , both facts contradict that  $s \square$  is strongly mixed by  $a$ .

By transitivity of mixing the second assertion, that  $s \square \wedge m \square$  and  $s \square \wedge n \square$  are mixed by  $a$  for every  $m, n \ll a$ , follows from the first one.

*Case (b).* Let  $s$  be min-separated by  $a$ . Assume to the contrary that  $s \wedge m$  and  $s \wedge n \uparrow$  resp.  $s \wedge m \uparrow$  and  $s \wedge n \uparrow$  are not mixed by  $a$  for some  $m, n \ll a$  with  $\min(m) = \min(n)$ . Hence there exists  $b \ll a$  such that  $s \wedge m$  and  $s \wedge n \uparrow$  resp.  $s \wedge m \uparrow$  and  $s \wedge n \uparrow$  are separated by  $b$ . Since  $s$  is min-separated by  $a$ , by Lemma 2.2 we get that  $s \wedge m$  and  $s \wedge n$  are also mixed by  $b$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$ .

Now choose  $k$  minimal such that  $m, n < b(k)$ . By definition of separation we must have that  $s \wedge m$  and  $s \wedge n \uparrow \wedge \langle b(k) \rangle$  resp.  $s \wedge m \uparrow \wedge \langle b(k) \rangle$  and  $s \wedge n \uparrow \wedge \langle b(k) \rangle$  are separated by  $b$ . However, since  $\langle b(k) \rangle \ll a$ , both facts contradict that  $s$  is min-separated by  $a$ .

*Case (c).* Let  $s \square$  be max-separated by  $a$ . First, we prove that  $s \square$  and  $s \square \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . Assume to the contrary that  $s \square$  and  $s \square \wedge m \uparrow$  are not mixed by  $a$  for some  $m \ll a$ . Hence there exists  $b \ll a$  such that  $s \square$  and  $s \square \wedge m \uparrow$  are separated by  $b$ . Since  $s \square$  is max-separated by  $a$ , by Lemma 2.2 we get that  $s \square \wedge m$  and  $s \square \wedge n$  are also mixed by  $b$  for every  $m, n \ll a$  with  $\max(m) = \max(n)$ .

Now choose  $k$  minimal such that  $m < b(k)$ . By definition of separation we must have that  $s \square \wedge \langle b(k) \rangle$  and  $s \square \wedge m \uparrow \wedge \langle b(k) \rangle$  are separated by  $b$ . However, since  $\langle b(k) \rangle \ll a$ , this contradicts that  $s \square$  is max-separated by  $a$ .

By transitivity of mixing the second assertion, that  $s \square \wedge m \uparrow$  and  $s \square \wedge n \uparrow$  are mixed by  $a$  for every  $m, n \ll a$ , follows from the first one.

*Case (d).* Let  $s$  be min-max-separated by  $a$ . Assume to the contrary that  $s \wedge m \uparrow$  and  $s \wedge n \uparrow$  are not mixed by  $a$  for some  $m, n \ll a$  with  $\min(m) = \min(n)$ . Hence there exists  $b \ll a$  such that  $s \wedge m \uparrow$  and  $s \wedge n \uparrow$  are separated by  $b$ . Since  $s$  is min-max-separated by  $a$ , by Lemma 2.2 we get that  $s \wedge m$  and  $s \wedge n$  are also mixed by  $b$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$  and  $\max(m) = \max(n)$ .

Now choose  $k$  minimal such that  $m, n < b(k)$ . By definition of separation we must have that  $s \wedge m \uparrow \wedge \langle b(k) \rangle$  and  $s \wedge n \uparrow \wedge \langle b(k) \rangle$  are separated by  $b$ . However, since  $\langle b(k) \rangle \ll a$ , this contradicts that  $s$  is min-max-separated by  $a$ .  $\dashv$

LEMMA 2.8. *For every  $a$  there exists  $b \ll a$  which completely decides every  $s \ll b$ .*

PROOF. Inductively, we construct  $b_j \in \Omega^\omega$  for every  $j < \omega$ . By Lemma 2.5 there exists  $b_0 \ll a$  such that  $b_0$  completely decides  $\emptyset$ . Assume by induction that  $b_0, \dots, b_j$  have been constructed such that for every  $i \leq j$  and all  $s \ll \langle b_k(0): k < i \rangle$  the set  $b_i$  completely decides  $s$ . Some applications of Lemma 2.5 yield  $b_{j+1} \ll b_j \uparrow 1$  such that the inductive assumption is also satisfied for  $b_0, \dots, b_{j+1}$ . Then  $b = \langle b_j(0): j \in \omega \rangle$  has the desired properties.  $\dashv$

DEFINITION. We say that  $a$  is *canonical for  $\Delta$*  iff  $a$  satisfies the following properties:

- (a) The mapping  $\Delta 1(a)^\omega$  is continuous.
- (b) If  $s, t \ll a$ , then  $s \sqcap$  and  $t \sqcap$  are decided by  $a$ .
- (c) Every  $s \ll a$  is completely decided by  $a$ .
- (d) Let  $s, t \ll a$ . Then  $s \sqcap$  and  $s \sqcap \wedge m \sqcap$  are either mixed by  $a$  for all  $m \ll a$  or separated by  $a$  for all  $m \ll a$ . Equally  $s \sqcap \wedge m \sqcap$  and  $t \sqcap \wedge m \sqcap$  as well as  $s \sqcap \wedge m \sqcap$  and  $t \sqcap \wedge m \sqcap \wedge n \sqcap$  are in each case either mixed by  $a$  for all  $m, n \ll a$  or separated by  $a$  for all  $m, n \ll a$ .
- (e) If  $s \ll a$ , then either for every  $x \ll a$  and all  $k \in \omega$  the set  $s \sqcap \wedge (x \uparrow k)$  is strongly mixed by  $a$  or for every  $x \ll a$  there exists  $k \in \omega$  such that  $s \sqcap \wedge (x \uparrow k)$  is separated in some sense by  $a$ .
- (f) We have that either for every  $x \ll a$  there exists  $k \in \omega$  such that  $x \uparrow j$  is strongly mixed by  $a$  for every  $j \geq k$  or for every  $x \ll a$  there exists no  $k \in \omega$  such that  $x \uparrow j$  is strongly mixed by  $a$  for every  $j \geq k$ .
- (g) There exists  $b$  with  $a = \langle b(3i) \cup b(3i+1) \cup b(3i+2): i < \omega \rangle$  such that the properties (a) to (e) are even true for  $b$  instead of  $a$ .

LEMMA 2.9. *There exists  $a$  which is canonical for  $\Delta$ .*

PROOF. First, observe by Lemma 2.2 that if  $s \sqcap$  and  $t \sqcap$  are decided by  $a$ , then they are also decided by each  $b \ll a$ , and  $b$  decides in the same way as  $a$  does. Hence by Lemma 2.1, 2.4 and 2.8 we are guaranteed that there exists  $b_0$ , which satisfies the properties (a) to (c) of canonical.

Now we turn to property (d). We show that there exists  $b_1 \ll b_0$  such that for every  $s \ll b_1$  the sets  $s$  and  $s \wedge m$  are either mixed by  $b_1$  for all  $m \ll b_1$  or separated by  $b_1$  for all  $m \ll b_1$ .

Inductively, we construct  $c_j \in \Omega^\omega$  for every  $j < \omega$ . By Theorem 0.3.4 we can find  $c_0 \ll b_0$  such that the sets  $\emptyset$  and  $\emptyset \wedge m$  are either mixed by  $c_0$  for every  $m \ll c_0$  or separated by  $c_0$  for every  $m \ll c_0$ . Assume that  $c_0, \dots, c_j$  have been constructed such that for all  $i \leq j$  and for all  $s \ll \langle c_l(0): l < i \rangle$  the sets  $s$  and  $s \wedge m$  are either mixed by  $c_i$  for every  $m \ll c_i$  or separated by  $c_i$  for every  $m \ll c_i$ . Again, invoking Theorem 0.3.4 there exists  $c_{j+1} \ll c_j \uparrow 1$  such that the inductive assumption is also satisfied for  $c_0, \dots, c_{j+1}$ .

Then  $b_1 = \langle c_j(0) : j \in \omega \rangle$  has the desired property. Applying some similar inductions, we get  $b_1$  fulfilling (a) to (d) of canonical.

Now we turn to property (e). Inductively, we construct  $c_j \in \Omega^\omega$  for every  $j < \omega$ . Consider the set  $C = \{x \ll b_1 : \forall k \in \omega \ x \upharpoonright k \text{ is strongly mixed by } b_1\}$ . Since  $C$  is closed, by Theorem 1.8 there exists  $c_0 \ll b_1$  such that  $(c_0)^\omega \subseteq C$  or  $(c_0)^\omega \subseteq \Omega^\omega \setminus C$ . Assume that  $c_0, \dots, c_j$  have been constructed such that for all  $i \leq j$  and for all  $s \ll \langle c_l(0) : l < i \rangle$  either for every  $x \ll c_i$  and all  $k \in \omega$  the set  $s \sqcap \hat{\ } (x \upharpoonright k)$  is strongly mixed by  $c_i$  or for every  $x \ll c_i$  there exists  $k \in \omega$  such that  $s \sqcap \hat{\ } (x \upharpoonright k)$  is separated in some sense by  $c_i$ . For every  $s \ll \langle c_l(0) : l < j \rangle$  consider the sets  $C_{s \sqcap \hat{\ }} = \{x \ll c_j : \forall k \in \omega \ s \sqcap \hat{\ } \langle c_l(0) \rangle \sqcap \hat{\ } (x \upharpoonright k) \text{ is strongly mixed by } c_j\}$ . Again, all  $C_{s \sqcap \hat{\ }}$  are closed. Hence some applications of Theorem 1.8 yield  $c_{j+1} \ll c_j \upharpoonright 1$  such that the inductive assumption is also satisfied for  $c_0, \dots, c_{j+1}$ . Then  $b_2 = \langle c_j(0) : j \in \omega \rangle$  satisfies the properties (a) to (e) of canonical.

Now we turn to property (f). We consider the set  $U = \{x \ll b_2 : \exists k_x \in \omega \ \forall j \geq k_x \ x \upharpoonright j \text{ is strongly mixed by } b_2\}$ . By (e) of canonical for all  $x \in U$  we have the open set  $(x \upharpoonright k_x, b_2)^\omega$  such that  $x \in (x \upharpoonright k_x, b_2)^\omega \subseteq U$ . Since  $U = \bigcup_{x \in U} (x \upharpoonright k_x, b_2)^\omega$  holds, we get that  $U$  is open. Now by Theorem 1.2 there exists  $b \ll b_2$  such that  $(b)^\omega \subseteq U$  or  $(b)^\omega \subseteq \Omega^\omega \setminus U$ . Hence  $b$  additionally satisfies property (f) of canonical.

Finally, let  $a = \langle b(3i) \cup b(3i+1) \cup b(3i+2) : i < \omega \rangle$ . Hence  $a$  has the properties (a) to (g). This completes the proof.  $\dashv$

For the remainder of this work let  $a$  be canonical for  $\Delta$ .

LEMMA 2.10. *Let  $s \ll a$ .*

- (a) *Let  $s$  be min-separated by  $a$ . If  $x, y \ll a$ , then  $\Delta(s \hat{\ } x) = \Delta(s \hat{\ } y)$  implies  $\min(x(0)) = \min(y(0))$ .*
- (b) *Let  $s \sqcap$  be max-separated by  $a$ . If  $x, y \ll a$ , then  $\Delta(s \sqcap \hat{\ } x) = \Delta(s \sqcap \hat{\ } y)$  implies  $\max(x(0)) = \max(y(0))$ .*
- (c) *Let  $s$  be min-max-separated by  $a$ . If  $x, y \ll a$ , then  $\Delta(s \hat{\ } x) = \Delta(s \hat{\ } y)$  implies  $\min(x(0)) = \min(y(0))$  and  $\max(x(0)) = \max(y(0))$ .*
- (d) *Let  $s \sqcap$  be strongly separated by  $a$ . If  $x, y \ll a$ , then  $\Delta(s \sqcap \hat{\ } x) = \Delta(s \sqcap \hat{\ } y)$  implies that there exists  $k$  such that  $x(0) = y(0) \cap k$  or  $y(0) = x(0) \cap k$ , i.e., either  $x(0)$  is an initial segment of  $y(0)$  or conversely.*

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \sqcap \hat{\ } x) = \Delta(s \sqcap \hat{\ } y)$ . Notice that we can assume without loss of generality that  $\max(x(0)) < \max(y(0))$ ; since  $\max(x(0)) = \max(y(0))$  together with the hypothesis of each of the four cases implies that  $s \sqcap \hat{\ } \langle x(0) \rangle$  and  $s \sqcap \hat{\ } \langle y(0) \rangle$  are mixed by  $a$ , and the assertion follows by Lemma 2.5.

First of all, we show that if  $s \sqcap$  is separated in some sense by  $a$  and  $\max(x(0)) < \max(y(0))$ , we must have that  $\min(x(0)) = \min(y(0))$ . For that purpose assume to the contrary that  $\min(x(0)) \neq \min(y(0))$ . We distinguish three cases.

First, let  $\max(x(0)) < \min(y(0))$ . Since  $\Delta(s \hat{x}) = \Delta(s \hat{y})$ , we have that  $s \hat{\langle x(0) \rangle}$  and  $s$  are mixed by  $a$ . Hence by (d) of canonical we must have that  $s \hat{m}$  and  $s$  are mixed by  $a$  for all  $m \ll a$ . By transitivity of mixing it follows that  $s \hat{m}$  and  $s \hat{n}$  are mixed by  $a$  for every  $m, n \ll a$ . But this contradicts that  $s$  is separated in some sense by  $a$ .

Next, suppose that  $\min(x(0)) < \min(y(0))$  and  $\max(x(0)) > \min(y(0))$ . Let  $v$  be the part of  $x(0)$  below  $\min(y(0))$ . Since  $\Delta(s \hat{x}) = \Delta(s \hat{y})$ , we have that  $s \hat{\langle v \rangle \uparrow}$  and  $s$  are mixed by  $a$ . Thus, by (d) of canonical we must have that  $s \hat{m \uparrow}$  and  $s$  are mixed by  $a$  for all  $m \ll a$ . Now let  $w$  denote the part of  $y(0)$  less than or equal to  $\max(x(0))$ . Hence we have that  $s \hat{\langle x(0) \rangle}$  and  $s \hat{\langle w \rangle \uparrow}$  are mixed by  $a$ . By transitivity of mixing  $s \hat{\langle x(0) \rangle}$  and  $s$  are mixed by  $a$ . Therefore, by (d) of canonical we must have that  $s \hat{m}$  and  $s$  are mixed by  $a$  for all  $m \ll a$ . Again, by transitivity of mixing it follows that  $s \hat{m}$  and  $s \hat{n}$  are mixed by  $a$  for every  $m, n \ll a$ . But this contradicts that  $s$  is separated in some sense by  $a$ .

Finally, assume that  $\min(x(0)) > \min(y(0))$ . Let  $v$  be the part of  $y(0)$  below  $\min(x(0))$ . Since  $\Delta(s \hat{x}) = \Delta(s \hat{y})$ , we have that  $s$  and  $s \hat{\langle v \rangle \uparrow}$  are mixed by  $a$ . Thus, by (d) of canonical we must have that  $s$  and  $s \hat{m \uparrow}$  are mixed by  $a$  for all  $m \ll a$ . Now let  $w$  denote the part of  $y(0)$  less than or equal to  $\max(x(0))$ . Hence we have that  $s \hat{\langle x(0) \rangle}$  and  $s \hat{\langle w \rangle \uparrow}$  are mixed by  $a$ . By transitivity of mixing  $s \hat{\langle x(0) \rangle}$  and  $s$  are mixed by  $a$ . Therefore, by (d) of canonical we must have that  $s \hat{m}$  and  $s$  are mixed by  $a$  for all  $m \ll a$ . Again, by transitivity of mixing it follows that  $s \hat{m}$  and  $s \hat{n}$  are mixed by  $a$  for every  $m, n \ll a$ . But this contradicts that  $s$  is separated in some sense by  $a$ .

Therewith we must have  $\min(x(0)) = \min(y(0))$ . This already proves case (a) of this lemma.

Now we prove case (b) and (c) in one step. For that purpose let  $s$  be max-separated or min-max-separated by  $a$ . Recall that we can assume without loss of generality  $\min(x(0)) = \min(y(0))$  and  $\max(x(0)) < \max(y(0))$ .

Let  $v$  be the part of  $y(0)$  less than or equal to  $\max(x(0))$ . Hence we have that  $s \hat{\langle x(0) \rangle}$  and  $s \hat{\langle v \rangle \uparrow}$  are mixed by  $a$ . Additionally, the cases (c) and (d) of Lemma 2.7 yield that  $s \hat{m \uparrow}$  and  $s \hat{n \uparrow}$  are mixed by  $a$  for all  $m, n \ll a$  with  $\min(m) = \min(n)$ . Therefore  $s \hat{\langle v \rangle \uparrow}$  and  $s \hat{\langle x(0) \rangle \uparrow}$  are mixed by  $a$ , because  $\min(v) = \min(x(0))$ . By transitivity of mixing we get that  $s \hat{\langle x(0) \rangle}$  and  $s \hat{\langle x(0) \rangle \uparrow}$  are mixed by  $a$ . Moreover by (d) of canonical we must have that  $s \hat{m}$  and  $s \hat{m \uparrow}$  are mixed by  $a$  for every  $m \ll a$ . Altogether, we have that  $s \hat{\langle x(0) \rangle}$  and  $s \hat{\langle x(0) \rangle \uparrow}$ ,  $s \hat{\langle x(0) \rangle \uparrow}$  and  $s \hat{\langle y(0) \rangle \uparrow}$  as well as  $s \hat{\langle y(0) \rangle \uparrow}$  and  $s \hat{\langle y(0) \rangle}$  are mixed by  $a$ . Again, by transitivity of mixing we obtain that  $s \hat{\langle x(0) \rangle}$  and  $s \hat{\langle y(0) \rangle}$  are mixed by  $a$ . But this contradicts the fact that  $s$  is max-separated or min-max-separated by  $a$ .

Hence we must have that  $\max(x(0)) = \max(y(0))$ , and the assertion follows by Lemma 2.5.

Finally, we prove case (d) of this lemma. For that purpose let  $s$  be strongly separated by  $a$ . Recall that we can assume without loss of generality  $\min(x(0)) =$



$\min(y(0))$  and  $\max(x(0)) < \max(y(0))$ . Suppose to the contrary that  $x(0)$  is not an initial segment of  $y(0)$ .

Let  $b$  with  $a \ll b$  be as in (g) of canonical. Moreover let  $v$  denote the longest common initial segment of  $x(0)$  and  $y(0)$ . Choose  $k$  with  $\min(x(0) \triangle y(0)) \in b(k)$ . Since  $x(0)$  is not an initial segment of  $y(0)$ , we have that  $s \sqcap \langle v \rangle \uparrow$  and  $s \sqcap \langle v \rangle \uparrow \wedge \langle b(k) \rangle \uparrow$  are mixed by  $b$ . Hence by (d) of canonical we must have that  $s \sqcap \langle m \rangle \uparrow$  and  $s \sqcap \langle m \rangle \uparrow \wedge \langle n \rangle \uparrow$  are mixed by  $b$  for all  $m, n \ll b$ . Furthermore, let  $w$  denote the part of  $y(0)$  less than or equal to  $\max(x(0))$ . Therewith we get that  $s \sqcap \langle x(0) \rangle$  and  $s \sqcap \langle w \rangle \uparrow$  are mixed by  $b$ , too. Since  $v$  is an initial segment of  $w$ , we get with the result above that  $s \sqcap \langle w \rangle \uparrow$  and  $s \sqcap \langle v \rangle \uparrow$  are mixed by  $b$ . Hence by transitivity of mixing  $s \sqcap \langle x(0) \rangle$  and  $s \sqcap \langle v \rangle \uparrow$  are mixed by  $b$ . Moreover since  $v$  is an initial segment of  $x(0)$ , property (d) of canonical yields that  $s \sqcap \langle m \rangle \uparrow$  and  $s \sqcap \langle m \rangle \uparrow \wedge \langle n \rangle$  are mixed by  $b$  for all  $m, n \ll b$ . Thus, equally  $s \sqcap \langle y(0) \rangle$  and  $s \sqcap \langle v \rangle \uparrow$  are mixed by  $b$ , and by transitivity we obtain that  $s \sqcap \langle x(0) \rangle$  and  $s \sqcap \langle y(0) \rangle$  are mixed by  $b$ . Since  $a \ll b$ , we must have that  $s \sqcap \langle x(0) \rangle$  and  $s \sqcap \langle y(0) \rangle$  are also mixed by  $a$ . But this contradicts that  $s \sqcap$  is strongly separated by  $a$ .

Hence we must have that  $x(0)$  is an initial segment of  $y(0)$ . This completes the proof.  $\dashv$

LEMMA 2.11. *Let  $s, t \ll a$ . Suppose  $s \sqcap$  and  $t \sqcap$  are mixed by  $a$  and  $s \sqcap$  is separated in some sense by  $a$ . If  $x, y \ll a$  such that  $\Delta(s \sqcap \langle x \rangle) = \Delta(t \sqcap \langle y \rangle)$ , then  $\max(x(0)) > \min(y(0))$ .*

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \sqcap \langle x \rangle) = \Delta(t \sqcap \langle y \rangle)$ . Assume to the contrary that  $\max(x(0)) < \min(y(0))$ . Note that  $\max(x(0)) = \min(y(0))$  is impossible by (g) of canonical.

Choose  $0 < k \leq \text{dom}(t)$  maximal with  $\max(t(k-1)) \leq \max(x(0))$  if possible, otherwise choose  $k = 0$ . Moreover if  $k < \text{dom}(t)$ , let  $v$  denote the part of  $t(k)$  less than or equal to  $\max(x(0))$ . Thus, if  $k = \text{dom}(t)$  or  $v = \emptyset$ , we have that  $s \sqcap \langle x(0) \rangle$  and  $t \uparrow k \sqcap$  are mixed by  $a$ . Otherwise we have that  $s \sqcap \langle x(0) \rangle$  and  $t \uparrow k \wedge \langle v \rangle \uparrow$  are mixed by  $a$ .

Moreover since  $s \sqcap$  and  $t \sqcap$  are mixed by  $a$ , there exist  $x_0, y_0 \ll a$  with  $s < x_0, y_0$  and  $t < x_0, y_0$  such that  $\Delta(s \sqcap \langle x_0 \rangle) = \Delta(t \sqcap \langle y_0 \rangle)$ .

Now assume that we are in the first case, where  $s \sqcap \langle x(0) \rangle$  and  $t \uparrow k \sqcap$  are mixed by  $a$ . If  $k < \text{dom}(t)$ , we can choose  $y_1 \ll a$  by  $y_1 = \langle t(i) : k \leq i < \text{dom}(t) \rangle \wedge y_0$  such that  $\Delta(s \sqcap \langle x_0 \rangle) = \Delta(t \uparrow k \sqcap \langle y_1 \rangle)$ . By choice of  $k$  we have  $s < x_0, y_1$  and  $t \uparrow k < x_0, y_1$ . Hence by (b) of canonical we must have that  $s \sqcap$  and  $t \uparrow k \sqcap$  are mixed by  $a$ .

Next, suppose that we are in the case, where  $s \sqcap \langle x(0) \rangle$  and  $t \uparrow k \wedge \langle v \rangle \uparrow$  are mixed by  $a$ . Let  $w$  be the part of  $t(k)$  above  $\max(v)$ . If  $k < \text{dom}(t) - 1$ , choose  $y_1 \ll a$  by  $y_1 = \langle w \rangle \wedge \langle t(i) : k < i < \text{dom}(t) \rangle \wedge y_0$ , otherwise choose  $y_1 = \langle w \rangle \wedge y_0$ . Therewith we have that  $\Delta(s \sqcap \langle x_0 \rangle) = \Delta(t \uparrow k \wedge \langle v \rangle \uparrow \wedge \langle y_1 \rangle)$  with  $s < x_0, y_1$  and  $t \uparrow k \wedge \langle v \rangle < x_0, y_1$ . Thus, by (b) of canonical we get that  $s \sqcap$  and  $t \uparrow k \wedge \langle v \rangle \uparrow$  are mixed by  $a$ .

Since  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ , by transitivity of mixing we can conclude that  $s \sqsupseteq \wedge \langle x(0) \rangle$  and  $s \sqsupseteq$  are mixed by  $a$ , contradicting all cases of Lemma 2.10.  $\dashv$

LEMMA 2.12. *Let  $s, t \ll a$ . If  $x, y \ll a$  with  $\min(x(0)) = \min(y(0))$  such that  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , then  $s \sqsupseteq \wedge m \uparrow$  and  $t \sqsupseteq \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ .*

PROOF. Let  $b$  with  $a \ll b$  be as in (g) of canonical. Choose  $k$  with  $\min(x(0)) \in b(k)$ . Since  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , we have that  $s \sqsupseteq \wedge \langle b(k) \rangle \uparrow$  and  $t \sqsupseteq \wedge \langle b(k) \rangle \uparrow$  are mixed by  $b$ . By (d) of canonical we must have that  $s \sqsupseteq \wedge m \uparrow$  and  $t \sqsupseteq \wedge m \uparrow$  are mixed by  $b$  for every  $m \ll b$ . Since  $a \ll b$ , by Lemma 2.2 we also have that  $s \sqsupseteq \wedge m \uparrow$  and  $t \sqsupseteq \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ .  $\dashv$

LEMMA 2.13. *Let  $s, t \ll a$ . Suppose  $s$  and  $t$  are mixed by  $a$  and both  $s$  and  $t$  are min-separated by  $a$ . If  $x, y \ll a$  such that  $\Delta(s \wedge x) = \Delta(t \wedge y)$ , then  $\min(x(0)) = \min(y(0))$ .*

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \wedge x) = \Delta(t \wedge y)$ . Assume to the contrary that  $\min(x(0)) \neq \min(y(0))$ . By symmetry we can suppose without loss of generality that  $\min(x(0)) < \min(y(0))$ . Moreover by Lemma 2.11 it suffices to prove that the assumption that  $\min(x(0)) < \min(y(0))$  and  $\max(x(0)) > \min(y(0))$  leads to a contradiction.

Let  $v$  be the part of  $x(0)$  below  $\min(y(0))$ . Since  $\Delta(s \wedge x) = \Delta(t \wedge y)$ , we must have that  $s \wedge \langle v \rangle \uparrow$  and  $t$  are mixed by  $a$ . Moreover since  $s$  and  $t$  are mixed by  $a$ , by transitivity of mixing we get that  $s \wedge \langle v \rangle \uparrow$  and  $s$  are mixed by  $a$ , contradicting case (a) of Lemma 2.10.  $\dashv$

LEMMA 2.14. *Let  $s, t \ll a$ . Suppose  $s$  and  $t$  are mixed by  $a$  and both  $s$  and  $t$  are min-separated by  $a$ . Then  $s \wedge m$  and  $t \wedge n$  are mixed by  $a$  for all  $m, n \ll a$  with  $\min(m) = \min(n)$ .*

PROOF. Since  $s$  and  $t$  are mixed by  $a$ , there exist  $x, y \ll a$  such that  $\Delta(s \wedge x) = \Delta(t \wedge y)$ . By Lemma 2.13 we have that  $\min(x(0)) = \min(y(0))$ .

Moreover by Lemma 2.12 we get that  $s \wedge m \uparrow$  and  $t \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . Additionally, case (b) of Lemma 2.7 yields that  $s \wedge m$  and  $s \wedge m \uparrow$  as well as  $t \wedge m \uparrow$  and  $t \wedge n$  are mixed by  $a$  for all  $m, n \ll a$  with  $\min(m) = \min(n)$ . Thus, by transitivity of mixing we get that  $s \wedge m$  and  $t \wedge n$  are mixed by  $a$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$ .  $\dashv$

LEMMA 2.15. *Let  $s, t \ll a$ . Suppose  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$  and both  $s \sqsupseteq$  and  $t \sqsupseteq$  are max-separated by  $a$ . If  $x, y \ll a$  such that  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , then  $\max(x(0)) = \max(y(0))$ .*

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ . Assume to the contrary that  $\max(x(0)) \neq \max(y(0))$ . By symmetry we can suppose without loss of

generality that  $\max(x(0)) < \max(y(0))$ . Moreover by Lemma 2.11 it suffices to prove that the assumption that  $\max(x(0)) < \max(y(0))$  and  $\max(x(0)) > \min(y(0))$  leads to a contradiction.

So let  $w$  be the part of  $y(0)$  less than or equal to  $\max(x(0))$ . Since  $\Delta(s \sqcup x) = \Delta(t \sqcup y)$ , we have that  $s \sqcup \langle x(0) \rangle$  and  $t \sqcup \langle w \rangle \uparrow$  are mixed by  $a$ . Additionally, case (c) of Lemma 2.7 yields that  $t \sqcup$  and  $t \sqcup \langle w \rangle \uparrow$  are mixed by  $a$ . Moreover since  $s \sqcup$  and  $t \sqcup$  are mixed by  $a$ , by transitivity of mixing it follows that  $s \sqcup$  and  $s \sqcup \langle x(0) \rangle$  are mixed by  $a$ , contradicting case (b) of Lemma 2.10.  $\dashv$

LEMMA 2.16. *Let  $s, t \ll a$ . Suppose  $s \sqcup$  and  $t \sqcup$  are mixed by  $a$  and both  $s \sqcup$  and  $t \sqcup$  are max-separated by  $a$ . Then  $s \sqcup \langle m \rangle$  and  $t \sqcup \langle n \rangle$  are mixed by  $a$  for all  $m, n \ll a$  with  $\max(m) = \max(n)$ .*

PROOF. Since  $s \sqcup$  and  $t \sqcup$  are mixed by  $a$ , there exist  $x, y \ll a$  such that  $\Delta(s \sqcup x) = \Delta(t \sqcup y)$ . By Lemma 2.15 we have that  $\max(x(0)) = \max(y(0))$ . Hence by definition of mixing we must have that  $s \sqcup \langle x(0) \rangle$  and  $t \sqcup \langle y(0) \rangle$  are mixed by  $a$ . Moreover we have that  $t \sqcup \langle x(0) \rangle$  and  $t \sqcup \langle y(0) \rangle$  are mixed by  $a$ , because  $t \sqcup$  is max-separated by  $a$ . By transitivity of mixing we get that  $s \sqcup \langle x(0) \rangle$  and  $t \sqcup \langle x(0) \rangle$  are mixed by  $a$ . Thus, (d) of canonical yields that  $s \sqcup \langle m \rangle$  and  $t \sqcup \langle m \rangle$  are mixed by  $a$  for all  $m \ll a$ .

Again, since  $t \sqcup$  is max-separated by  $a$ , we have that  $t \sqcup \langle m \rangle$  and  $t \sqcup \langle n \rangle$  are mixed by  $a$  for every  $m, n \ll a$  with  $\max(m) = \max(n)$ . Finally, by transitivity of mixing we obtain that  $s \sqcup \langle m \rangle$  and  $t \sqcup \langle n \rangle$  are mixed by  $a$  for all  $m, n \ll a$  with  $\max(m) = \max(n)$ .  $\dashv$

LEMMA 2.17. *Let  $s, t \ll a$ . Suppose  $s$  and  $t$  are mixed by  $a$  and both  $s$  and  $t$  are min-max-separated by  $a$ . If  $x, y \ll a$  such that  $\Delta(s \wedge x) = \Delta(t \wedge y)$ , then  $\min(x(0)) = \min(y(0))$  and  $\max(x(0)) = \max(y(0))$ .*

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \wedge x) = \Delta(t \wedge y)$ . First, assume to the contrary that  $\min(x(0)) \neq \min(y(0))$ . By symmetry we can suppose without loss of generality that  $\min(x(0)) < \min(y(0))$ . Moreover by Lemma 2.11 it suffices to prove that the assumption that  $\min(x(0)) < \min(y(0))$  and  $\max(x(0)) > \min(y(0))$  leads to a contradiction.

Let  $v$  be the part of  $x(0)$  below  $\min(y(0))$ . Since  $\Delta(s \wedge x) = \Delta(t \wedge y)$  we must have that  $s \wedge \langle v \rangle \uparrow$  and  $t$  are mixed by  $a$ . Since  $s$  and  $t$  are mixed by  $a$ , by transitivity of mixing we get that  $s \wedge \langle v \rangle \uparrow$  and  $s$  are mixed by  $a$ , contradicting case (c) of Lemma 2.10.

Hence we must have  $\min(x(0)) = \min(y(0))$ . Now assume to the contrary that  $\max(x(0)) \neq \max(y(0))$ . Equally by symmetry we can suppose without loss of generality that  $\max(x(0)) < \max(y(0))$ .

Let  $w$  be the part of  $y(0)$  less than or equal to  $\max(x(0))$ . Therewith we have that  $s \wedge \langle x(0) \rangle$  and  $t \wedge \langle w \rangle \uparrow$  are mixed by  $a$ . Since  $\min(x(0)) = \min(y(0))$ , Lemma 2.12 yields that  $s \wedge \langle w \rangle \uparrow$  and  $t \wedge \langle w \rangle \uparrow$  are mixed by  $a$ . By transitivity of mixing

we get that  $s \wedge \langle x(0) \rangle$  and  $s \wedge \langle w \rangle \uparrow$  are mixed by  $a$ , contradicting case (c) of Lemma 2.10. This completes the proof.  $\dashv$

LEMMA 2.18. *Let  $s, t \ll a$ . Suppose  $s$  and  $t$  are mixed by  $a$  and both  $s$  and  $t$  are min-max-separated by  $a$ . Then  $s \wedge m \uparrow$  and  $t \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . Moreover  $s \wedge m$  and  $t \wedge n$  are mixed by  $a$  for all  $m, n \ll a$  with  $\min(m) = \min(n)$  and  $\max(m) = \max(n)$ .*

PROOF. Since  $s$  and  $t$  are mixed by  $a$ , there exist  $x, y \ll a$  such that  $\Delta(s \wedge x) = \Delta(t \wedge y)$ . By Lemma 2.17 we get that  $\min(x(0)) = \min(y(0))$  and  $\max(x(0)) = \max(y(0))$ . Therefore, Lemma 2.12 yields that  $s \wedge m \uparrow$  and  $t \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ , which is our first assertion.

Additionally, by definition of mixing we have that  $s \wedge \langle x(0) \rangle$  and  $t \wedge \langle y(0) \rangle$  are mixed by  $a$ . Moreover we have that  $t \wedge \langle x(0) \rangle$  and  $t \wedge \langle y(0) \rangle$  are mixed by  $a$ , because  $t$  is min-max-separated by  $a$ . By transitivity of mixing we get that  $s \wedge \langle x(0) \rangle$  and  $t \wedge \langle x(0) \rangle$  are mixed by  $a$ . Thus, (d) of canonical yields that  $s \wedge m$  and  $t \wedge m$  are mixed by  $a$  for all  $m \ll a$ .

Again, since  $t$  is min-max-separated by  $a$ , we have that  $t \wedge m$  and  $t \wedge n$  are mixed by  $a$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$  and  $\max(m) = \max(n)$ . Finally, by transitivity of mixing we obtain that  $s \wedge m$  and  $t \wedge n$  are mixed by  $a$  for all  $m, n \ll a$  with  $\min(m) = \min(n)$  and  $\max(m) = \max(n)$ .  $\dashv$

LEMMA 2.19. *Let  $s, t \ll a$ . Suppose  $s \sqcap$  and  $t \sqcap$  are mixed by  $a$  and both  $s \sqcap$  and  $t \sqcap$  are strongly separated by  $a$ . If  $x, y \ll a$  such that  $\Delta(s \sqcap \wedge x) = \Delta(t \sqcap \wedge y)$ , then there exists  $k$  such that  $x(0) = y(0) \cap k$  or  $y(0) = x(0) \cap k$ , i. e., either  $x(0)$  is an initial segment of  $y(0)$  or conversely.*

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \sqcap \wedge x) = \Delta(t \sqcap \wedge y)$ . First, assume to the contrary that  $\min(x(0)) \neq \min(y(0))$ . By symmetry we can suppose without loss of generality that  $\min(x(0)) < \min(y(0))$ . Moreover by Lemma 2.11 it suffices to prove that the assumption that  $\min(x(0)) < \min(y(0))$  and  $\max(x(0)) > \min(y(0))$  leads to a contradiction.

Let  $v$  be the part of  $x(0)$  below  $\min(y(0))$ . Since  $\Delta(s \sqcap \wedge x) = \Delta(t \sqcap \wedge y)$ , we must have that  $s \sqcap \wedge \langle v \rangle \uparrow$  and  $t \sqcap$  are mixed by  $a$ . Moreover since  $s \sqcap$  and  $t \sqcap$  are mixed by  $a$ , by transitivity of mixing we get that  $s \sqcap \wedge \langle v \rangle \uparrow$  and  $s \sqcap$  are mixed by  $a$ , contradicting case (d) of Lemma 2.10.

Hence we must have that  $\min(x(0)) = \min(y(0))$ . Now assume to the contrary that neither  $x(0)$  is an initial segment of  $y(0)$  nor conversely. By symmetry we can suppose without loss of generality that  $x(0)$  is not an initial segment of  $y(0)$ .

Let  $v$  denote the longest common initial segment of  $x(0)$  and  $y(0)$ . Moreover choose  $k$  with  $\min(x(0) \triangle y(0)) \in a(k)$ . Since  $\Delta(s \sqcap \wedge x) = \Delta(t \sqcap \wedge y)$ , we have that either  $s \sqcap \wedge \langle v \rangle \uparrow \wedge \langle a(k) \rangle \sqcap$  and  $t \sqcap \wedge \langle v \rangle \uparrow$  or  $s \sqcap \wedge \langle v \rangle \uparrow$  and  $t \sqcap \wedge \langle v \rangle \uparrow \wedge \langle a(k) \rangle \uparrow$  are mixed by  $a$ . Additionally, by Lemma 2.12 we have that  $s \sqcap \wedge \langle v \rangle \uparrow$  and  $t \sqcap \wedge \langle v \rangle \uparrow$  are mixed by  $a$ , because  $\min(x(0)) = \min(y(0))$ . Thus, by transitivity of mixing

we get in the first case that  $s \sqcap \langle v \rangle \uparrow \wedge \langle a(k) \rangle \sqcap$  and  $s \sqcap \langle v \rangle \uparrow$ , in the second case that  $t \sqcap \langle v \rangle \uparrow$  and  $t \sqcap \langle v \rangle \uparrow \wedge \langle a(k) \rangle \uparrow$  are mixed by  $a$ . Both cases contradict case (d) of Lemma 2.10.  $\dashv$

Now we want to analyse the case that  $s \sqcap$  is strongly separated by  $a$ . Since  $a$  is canonical for  $\Delta$ , we are able to distinguish exactly two possibilities.

DEFINITION. Let  $s \ll a$ . Suppose that  $s \sqcap$  is strongly separated by  $a$ . We say that  $s \sqcap$  is still strongly separated by  $a$  iff  $s \sqcap \wedge m$  and  $s \sqcap \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . Moreover  $s \sqcap$  is very strongly separated by  $a$  iff  $s \sqcap \wedge m$  and  $s \sqcap \wedge m \uparrow$  are separated by  $a$  for every  $m \ll a$ .

The following lemma is a straightforward implication of the definition above.

LEMMA 2.20. Let  $s \ll a$ .

- (a) Let  $s \sqcap$  be still strongly separated by  $a$ . Then  $s \sqcap \wedge m \uparrow$  is still strongly separated by  $a$  for every  $m \ll a$ .
- (b) Let  $s \sqcap$  be very strongly separated by  $a$ . Then  $s \sqcap \wedge m \uparrow$  is very strongly separated by  $a$  for every  $m \ll a$ .

LEMMA 2.21. Let  $s, t \ll a$ . Suppose  $s \sqcap$  and  $t \sqcap$  are mixed by  $a$  and both  $s \sqcap$  and  $t \sqcap$  are very strongly separated by  $a$ . If  $x, y \ll a$  such that  $\Delta(s \sqcap \wedge x) = \Delta(t \sqcap \wedge y)$ , then  $x(0) = y(0)$ .

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \sqcap \wedge x) = \Delta(t \sqcap \wedge y)$ . By Lemma 2.19 we have that  $x(0)$  is an initial segment of  $y(0)$  or conversely. Moreover by symmetry we can suppose without loss of generality that  $x(0)$  is an initial segment of  $y(0)$ .

Assume to the contrary that  $\max(x(0)) < \max(y(0))$ . Since  $\Delta(s \sqcap \wedge x) = \Delta(t \sqcap \wedge y)$ , we have that  $s \sqcap \wedge \langle x(0) \rangle$  and  $t \sqcap \wedge \langle x(0) \rangle \uparrow$  are mixed by  $a$ . By (d) of canonical we get that  $s \sqcap \wedge m$  and  $t \sqcap \wedge m \uparrow$  are mixed by  $a$  for all  $m \ll a$ . Since  $\min(x(0)) = \min(y(0))$ , by Lemma 2.12 we also have that  $s \sqcap \wedge m \uparrow$  and  $t \sqcap \wedge m \uparrow$  are mixed by  $a$  for all  $m \ll a$ . Finally, by transitivity of mixing we can conclude that  $s \sqcap \wedge m$  and  $s \sqcap \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . But this contradicts our assumption that  $s \sqcap$  is very strongly separated by  $a$ .  $\dashv$

LEMMA 2.22. Let  $s, t \ll a$ . Suppose  $s \sqcap$  and  $t \sqcap$  are mixed by  $a$  and both  $s \sqcap$  and  $t \sqcap$  are very strongly separated by  $a$ . Then  $s \sqcap \wedge m$  and  $t \sqcap \wedge m$  are mixed by  $a$  for all  $m \ll a$ .

PROOF. Otherwise by (d) of canonical we would have that  $s \sqcap \wedge m$  and  $t \sqcap \wedge m$  are separated by  $a$  for every  $m \ll a$ . By Lemma 2.21 this would contradict that  $s \sqcap$  and  $t \sqcap$  are mixed by  $a$ , so the assertion follows.  $\dashv$

LEMMA 2.23. Let  $s, t \ll a$ . Suppose  $s \sqcap$  and  $t \sqcap$  are mixed by  $a$  and  $s \sqcap$  is strongly

*mixed by  $a$ . Moreover assume that  $t \sqsupseteq$  is either min-separated, min-max-separated or strongly separated by  $a$ . If  $x, y \ll a$  such that  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , then  $\max(x(0)) < \min(y(0))$ .*

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ . Assume to the contrary that  $\max(x(0)) > \min(y(0))$ . By Lemma 2.11 we must have that  $\min(x(0)) < \max(y(0))$ . We distinguish three cases.

For the first case suppose that  $\min(x(0)) > \min(y(0))$  and  $\min(x(0)) < \max(y(0))$ . Let  $v$  denote the part of  $y(0)$  below  $\min(x(0))$ . Since  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , we have that  $s \sqsupseteq$  and  $t \sqsupseteq \wedge \langle v \rangle \uparrow$  are mixed by  $a$ . Moreover since  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ , by transitivity of mixing we get that  $t \sqsupseteq$  and  $t \sqsupseteq \wedge \langle v \rangle \uparrow$  are mixed by  $a$ . But this contradicts case (a), (c) and (d) of Lemma 2.10.

Next, assume that  $\min(x(0)) \leq \min(y(0))$  and  $\max(x(0)) \geq \max(y(0))$ . Let  $v$  be the part of  $x(0)$  less than or equal to  $\max(y(0))$ . Since  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , we have that  $s \sqsupseteq \wedge \langle v \rangle \sqsupseteq$  and  $t \sqsupseteq \wedge \langle y(0) \rangle$  are mixed by  $a$ . Additionally, by (a) of Lemma 2.7 we have that  $s \sqsupseteq$  and  $s \sqsupseteq \wedge \langle v \rangle \sqsupseteq$  are mixed by  $a$ , because  $s \sqsupseteq$  is strongly mixed by  $a$ . Moreover since  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ , by transitivity of mixing we obtain that  $t \sqsupseteq$  and  $t \sqsupseteq \wedge \langle y(0) \rangle$  are mixed by  $a$ . Equally, this contradicts case (a), (c) and (d) of Lemma 2.10.

Finally, suppose that  $\min(x(0)) \leq \min(y(0))$  and  $\max(x(0)) < \max(y(0))$ . Let  $v$  denote the part of  $y(0)$  less than or equal to  $\max(x(0))$ . Since  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , we have that  $s \sqsupseteq \wedge \langle x(0) \rangle$  and  $t \sqsupseteq \wedge \langle v \rangle \uparrow$  are mixed by  $a$ . Additionally, by (a) of Lemma 2.7 we have that  $s \sqsupseteq$  and  $s \sqsupseteq \wedge \langle x(0) \rangle$  are mixed by  $a$ , because  $s \sqsupseteq$  is strongly mixed by  $a$ . Moreover since  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ , by transitivity of mixing we get that  $t \sqsupseteq$  and  $t \sqsupseteq \wedge \langle v \rangle \uparrow$  are mixed by  $a$ , a contradiction as above.  $\dashv$

LEMMA 2.24. *Let  $s, t \ll a$ . Suppose  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ ,  $s \sqsupseteq$  is strongly mixed by  $a$  and  $t \sqsupseteq$  is max-separated by  $a$ . If  $x, y \ll a$  such that  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , then  $\max(x(0)) < \max(y(0))$ .*

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ . Assume to the contrary that  $\max(x(0)) \geq \max(y(0))$ . By Lemma 2.11 we must have  $\min(x(0)) < \max(y(0))$ . We distinguish two cases.

First, suppose that  $\max(x(0)) > \max(y(0))$  and  $\min(x(0)) < \max(y(0))$ . Let  $v$  denote the part of  $x(0)$  less than or equal to  $\max(y(0))$ . Since  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , we have that  $s \sqsupseteq \wedge \langle v \rangle \uparrow$  and  $t \sqsupseteq \wedge \langle y(0) \rangle$  are mixed by  $a$ . Moreover by (a) of Lemma 2.7 we have that  $s \sqsupseteq \wedge \langle v \rangle \uparrow$  and  $s \sqsupseteq$  are mixed by  $a$ , because  $s \sqsupseteq$  is strongly mixed by  $a$ . Finally, since  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ , by transitivity of mixing we can conclude that  $t \sqsupseteq \wedge \langle y(0) \rangle$  and  $t \sqsupseteq$  are mixed by  $a$ . But this contradicts case (b) of Lemma 2.10.

Next, assume that  $\max(x(0)) = \max(y(0))$ . By definition of mixing we have that  $s \sqsupseteq \wedge \langle x(0) \rangle$  and  $t \sqsupseteq \wedge \langle y(0) \rangle$  are mixed by  $a$ . Since  $s \sqsupseteq$  is strongly mixed by  $a$ , by (a) of Lemma 2.7 we get that also  $s \sqsupseteq \wedge \langle x(0) \rangle$  and  $s \sqsupseteq$  are mixed by  $a$ . Moreover we have that  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ . Therefore, by transitivity of mixing we get

that  $t \sqcap \langle y(0) \rangle$  and  $t \sqcap$  are mixed by  $a$ , which equally contradicts case (b) of Lemma 2.10.  $\dashv$

LEMMA 2.25. *Let  $s, t \ll a$ . Suppose  $s$  and  $t$  are mixed by  $a$ ,  $s$  is min-separated by  $a$  and  $t$  is min-max-separated by  $a$ . If  $x, y \ll a$  such that  $\Delta(s \wedge x) = \Delta(t \wedge y)$ , then  $\min(x(0)) = \min(y(0))$  and  $\max(x(0)) < \max(y(0))$ .*

PROOF. Let  $x, y \ll a$  be such that  $\Delta(s \wedge x) = \Delta(t \wedge y)$ . First of all, we prove that we must have  $\min(x(0)) = \min(y(0))$ . For that purpose assume to the contrary that  $\min(x(0)) \neq \min(y(0))$ . Two applications of Lemma 2.11 yield that  $\max(x(0)) > \min(y(0))$  and  $\min(x(0)) < \max(y(0))$ . We distinguish two more cases.

First, suppose that  $\min(x(0)) < \min(y(0))$  and  $\max(x(0)) > \min(y(0))$ . Let  $v$  denote the part of  $x(0)$  below  $\min(y(0))$ . Since  $\Delta(s \wedge x) = \Delta(t \wedge y)$ , we have that  $s \wedge \langle v \rangle \uparrow$  and  $t$  are mixed by  $a$ . Moreover  $s$  and  $t$  are mixed by  $a$ . Hence by transitivity of mixing we get that  $s \wedge \langle v \rangle \uparrow$  and  $s$  are mixed by  $a$ . But this contradicts case (a) of Lemma 2.10.

Next, assume that  $\min(x(0)) > \min(y(0))$  and  $\min(x(0)) < \max(y(0))$ . Let  $v$  be the part of  $y(0)$  below  $\min(x(0))$ . Since  $\Delta(s \wedge x) = \Delta(t \wedge y)$ , we have that  $s$  and  $t \wedge \langle v \rangle \uparrow$  are mixed by  $a$ . Moreover  $s$  and  $t$  are mixed by  $a$ . Hence by transitivity of mixing we get that  $t$  and  $t \wedge \langle v \rangle \uparrow$  are mixed by  $a$ . This contradicts case (c) of Lemma 2.10.

Hence we have  $\min(x(0)) = \min(y(0))$ . Now we show that we also have that  $\max(x(0)) < \max(y(0))$ . For that purpose assume to the contrary that  $\max(x(0)) \geq \max(y(0))$ .

Let  $v$  denote the part of  $x(0)$  less than or equal to  $\max(y(0))$ . Since  $\Delta(s \wedge x) = \Delta(t \wedge y)$ , we have that  $s \wedge \langle v \rangle \sqcap$  and  $t \wedge \langle y(0) \rangle$  are mixed by  $a$ . Additionally, by (b) of Lemma 2.7 we get that  $s \wedge \langle y(0) \rangle \uparrow$  and  $s \wedge \langle v \rangle \sqcap$  are mixed by  $a$ , because  $\min(v) = \min(y(0))$ . Moreover since  $\min(x(0)) = \min(y(0))$ , Lemma 2.12 yields that  $s \wedge \langle y(0) \rangle \uparrow$  and  $t \wedge \langle y(0) \rangle \uparrow$  are also mixed by  $a$ . Altogether, by transitivity of mixing we get that  $t \wedge \langle y(0) \rangle$  and  $t \wedge \langle y(0) \rangle \uparrow$  are mixed by  $a$ . But this contradicts case (c) of Lemma 2.10.  $\dashv$

LEMMA 2.26. *Let  $s, t \ll a$ . Suppose  $s$  and  $t$  are mixed by  $a$ ,  $s$  is min-separated by  $a$  and  $t$  is min-max-separated by  $a$ . Then  $s \wedge m \uparrow$  and  $t \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ .*

PROOF. Since  $s$  and  $t$  are mixed by  $a$ , there exist  $x, y \ll a$  such that  $\Delta(s \wedge x) = \Delta(t \wedge y)$ . By Lemma 2.25 we must have that  $\min(x(0)) = \min(y(0))$ . Hence Lemma 2.12 yields that  $s \wedge m \uparrow$  and  $t \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ .  $\dashv$

LEMMA 2.27. *Let  $s, t \ll a$ .*

- (a) Suppose  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$  and both  $s \sqsupseteq$  and  $t \sqsupseteq$  are still strongly separated by  $a$ . Then  $s \sqsupseteq \wedge m \sqsupseteq$  and  $t \sqsupseteq \wedge m \sqsupseteq$  are mixed by  $a$  for every  $m \ll a$ .
- (b) Suppose  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ ,  $s \sqsupseteq$  is still strongly separated by  $a$  and  $t \sqsupseteq$  is very strongly separated by  $a$ . Then  $s \sqsupseteq \wedge m \sqsupseteq$  and  $t \sqsupseteq \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . Moreover  $s \sqsupseteq \wedge m \sqsupseteq$  and  $t \sqsupseteq \wedge m$  are separated by  $a$  for every  $m \ll a$ .
- (c) Suppose  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$  and both  $s \sqsupseteq$  and  $t \sqsupseteq$  are very strongly separated by  $a$ . Then  $s \sqsupseteq \wedge m \uparrow$  and  $t \sqsupseteq \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . Moreover  $s \sqsupseteq \wedge m$  and  $t \sqsupseteq \wedge m \uparrow$  are separated by  $a$  for every  $m \ll a$ .

PROOF. Since  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ , there exist  $x, y \ll a$  such that  $\Delta(s \sqsupseteq \wedge x) = \Delta(t \sqsupseteq \wedge y)$ . In each of the three cases both  $s \sqsupseteq$  and  $t \sqsupseteq$  are strongly separated by  $a$ . Hence by Lemma 2.19 we must have that  $x(0)$  is an initial segment of  $y(0)$  or conversely. Since  $\min(x(0)) = \min(y(0))$ , by Lemma 2.12 we have that  $s \sqsupseteq \wedge m \uparrow$  and  $t \sqsupseteq \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ .

The rest of the result follows directly by the definition of being still and very strongly separated, using the transitivity of mixing.  $\dashv$

LEMMA 2.28. Let  $s, t \ll a$ . Suppose  $s$  and  $t \sqsupseteq$  are mixed by  $a$  and  $s$  is min-separated by  $a$ . Then  $t \sqsupseteq$  is neither max-separated nor strongly separated by  $a$ .

PROOF. Since  $s$  and  $t \sqsupseteq$  are mixed by  $a$ , there exist  $x, y \ll a$  such that  $\Delta(s \wedge x) = \Delta(t \sqsupseteq \wedge y)$ . Assume to the contrary that  $t \sqsupseteq$  is either max-separated or strongly separated by  $a$ . Two applications of Lemma 2.11 yield that  $\min(x(0)) < \max(y(0))$  and  $\max(x(0)) > \min(y(0))$ . We distinguish five cases.

For the first case suppose that  $\min(x(0)) < \min(y(0))$  and  $\max(x(0)) > \min(y(0))$ . Let  $v$  be the part of  $x(0)$  below  $\min(y(0))$ . Since  $\Delta(s \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , we have that  $s \wedge \langle v \rangle \uparrow$  and  $t \sqsupseteq$  are mixed by  $a$ . Moreover  $s$  and  $t \sqsupseteq$  are also mixed by  $a$ . Hence by transitivity of mixing we obtain that  $s \wedge \langle v \rangle \uparrow$  and  $s$  are mixed by  $a$ . But this contradicts case (a) of Lemma 2.10.

Next, assume that  $t \sqsupseteq$  is max-separated by  $a$  and  $\min(x(0)) > \min(y(0))$  as well as  $\min(x(0)) < \max(y(0))$ . Choose  $k$  with  $\min(x(0)) \in a(k)$ . Let  $w$  denote the part of  $y(0)$  less than or equal to  $\max(a(k))$ . If  $\max(a(k)) < \max(y(0))$ , we get that  $s \wedge \langle a(k) \rangle \sqsupseteq$  and  $t \sqsupseteq \wedge \langle w \rangle \uparrow$  are mixed by  $a$ , because  $\Delta(s \wedge x) = \Delta(t \sqsupseteq \wedge y)$ . Otherwise, we must have that  $\max(a(k)) = \max(y(0))$ , since  $\min(x(0)) < \max(y(0))$ . Then we have that  $s \wedge \langle a(k) \rangle \sqsupseteq$  and  $t \sqsupseteq \wedge \langle y(0) \rangle$  are mixed by  $a$ . In the former case, by (c) of Lemma 2.7 we get that  $t \sqsupseteq \wedge \langle w \rangle \uparrow$  and  $t \sqsupseteq$  are mixed by  $a$ . Moreover since  $s$  and  $t \sqsupseteq$  are mixed by  $a$ , by transitivity of mixing we obtain that  $s \wedge \langle a(k) \rangle \sqsupseteq$  and  $s$  are mixed by  $a$ . This contradicts case (a) of Lemma 2.10. If we are in the latter case, we additionally have that  $t \sqsupseteq \wedge \langle y(0) \rangle$  and  $t \sqsupseteq \wedge \langle a(k) \rangle$  are mixed by  $a$ , because  $t \sqsupseteq$  is max-separated by  $a$  and  $\max(a(k)) = \max(y(0))$ . By transitivity of mixing we can conclude that  $s \wedge \langle a(k) \rangle \sqsupseteq$  and  $t \sqsupseteq \wedge \langle a(k) \rangle$  are mixed by  $a$ . Moreover by (d) of canonical we must have that  $s \wedge m \sqsupseteq$  and  $t \sqsupseteq \wedge m$  are mixed by  $a$  for all  $m \ll a$ .



Finally, by (b) of Lemma 2.7 we have that  $s \wedge m \uparrow$  and  $s \wedge n \uparrow$  are mixed by  $a$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$ . Again, by transitivity of mixing we obtain that  $t \uparrow \wedge m$  and  $t \uparrow \wedge n$  are mixed by  $a$  for all  $m, n \ll a$  with  $\min(m) = \min(n)$ . But this contradicts that  $t \uparrow$  is max-separated by  $a$ .

For the third case suppose that  $t \uparrow$  is strongly separated by  $a$  and  $\min(x(0)) > \min(y(0))$  as well as  $\min(x(0)) < \max(y(0))$ . Let  $v$  be the part of  $y(0)$  below  $\min(x(0))$ . Since  $\Delta(s \wedge x) = \Delta(t \uparrow \wedge y)$ , we have that  $s$  and  $t \uparrow \wedge \langle v \rangle \uparrow$  are mixed by  $a$ . Moreover  $s$  and  $t \uparrow$  are mixed by  $a$ . Hence by transitivity of mixing we obtain that  $t \uparrow$  and  $t \uparrow \wedge \langle v \rangle \uparrow$  are mixed by  $a$ . But this contradicts case (d) of Lemma 2.10.

Now assume that  $t \uparrow$  is max-separated by  $a$  and  $\min(x(0)) = \min(y(0))$ . By Lemma 2.12 we get that  $s \wedge m \uparrow$  and  $t \uparrow \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . Additionally, by (c) of Lemma 2.7 we have that  $t \uparrow \wedge m \uparrow$  and  $t \uparrow$  are mixed by  $a$  for all  $m \ll a$ , because  $t \uparrow$  is max-separated by  $a$ . Moreover since  $s$  and  $t \uparrow$  are mixed by  $a$ , by transitivity of mixing we obtain that  $s \wedge m \uparrow$  and  $s$  are mixed by  $a$  for every  $m \ll a$ . This contradicts case (a) of Lemma 2.10.

Finally, suppose that  $t \uparrow$  is strongly separated by  $a$  and  $\min(x(0)) = \min(y(0))$ . Equally, by Lemma 2.12 we get  $s \wedge m \uparrow$  and  $t \uparrow \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . Additionally, by (b) of Lemma 2.7 we have that  $s \wedge m \uparrow$  and  $s \wedge n \uparrow$  are mixed by  $a$  for all  $m, n \ll a$  with  $\min(m) = \min(n)$ . Thus, by transitivity of mixing we get that  $t \uparrow \wedge m \uparrow$  and  $t \uparrow \wedge n \uparrow$  are mixed by  $a$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$ . But this contradicts case (d) of Lemma 2.10.  $\dashv$

LEMMA 2.29. *Let  $s, t \ll a$ . Suppose  $s \uparrow$  and  $t \uparrow$  are mixed by  $a$  and  $s \uparrow$  is max-separated by  $a$ . Then  $t \uparrow$  is neither min-max-separated nor strongly separated by  $a$ .*

PROOF. Since  $s \uparrow$  and  $t \uparrow$  are mixed by  $a$ , there exist  $x, y \ll a$  such that  $\Delta(s \uparrow \wedge x) = \Delta(t \uparrow \wedge y)$ . Assume to the contrary that  $t \uparrow$  is either min-max-separated or strongly separated by  $a$ . Two applications of Lemma 2.11 yield that  $\max(x(0)) > \min(y(0))$  and  $\min(x(0)) < \max(y(0))$ . Now we distinguish three cases.

For the first case suppose that  $\min(x(0)) < \min(y(0))$  and  $\max(x(0)) > \min(y(0))$ . Let  $b$  with  $a \ll b$  be as in (g) of canonical. Moreover let  $v$  denote the part of  $x(0)$  below  $\min(y(0))$ . Choose  $k$  with  $\min(y(0)) \in b(k)$ . Additionally, let  $w$  denote the part of  $x(0)$  less than or equal to  $\max(b(k))$ . Since  $\Delta(s \uparrow \wedge x) = \Delta(t \uparrow \wedge y)$ , we have that  $s \uparrow \wedge \langle v \rangle \uparrow$  and  $t \uparrow$  as well as  $s \uparrow \wedge \langle w \rangle \uparrow$  and  $t \uparrow \wedge \langle b(k) \rangle \uparrow$  are mixed by  $b$ . By (c) and (g) of canonical we have that  $s \uparrow$  is also max-separated by  $b$ . Therefore, by (c) of Lemma 2.7 we get that  $s \uparrow \wedge \langle v \rangle \uparrow$  and  $s \uparrow \wedge \langle w \rangle \uparrow$  are mixed by  $b$ . Thus, by transitivity of mixing we obtain that  $t \uparrow$  and  $t \uparrow \wedge \langle w \rangle \uparrow$  are mixed by  $b$ . Now (d) and (g) of canonical yield that  $t \uparrow$  and  $t \uparrow \wedge m \uparrow$  are mixed by  $b$  for every  $m \ll b$ . Finally, since  $a \ll b$ , we can conclude that  $t \uparrow$  and  $t \uparrow \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . But this contradicts case (c) and (d) of Lemma 2.10.

Next, suppose that  $\min(x(0)) > \min(y(0))$  and  $\min(x(0)) < \max(y(0))$ . Let  $v$  be the part of  $y(0)$  below  $\min(x(0))$ . We have that  $s \uparrow$  and  $t \uparrow \wedge \langle v \rangle \uparrow$  are mixed by  $a$ .

Since  $s \sqsupseteq$  and  $t \sqsupseteq$  are mixed by  $a$ , we get that  $t \sqsupseteq$  and  $t \sqsupseteq \wedge \langle v \rangle \uparrow$  are mixed by  $a$ , a contradiction as above.

Finally, assume that  $\min(x(0)) = \min(y(0))$ . By Lemma 2.12 we have that  $s \sqsupseteq \wedge m \uparrow$  and  $t \sqsupseteq \wedge m \uparrow$  are mixed by  $a$  for all  $m \ll a$ . Moreover by (c) of Lemma 2.7 we have that  $s \sqsupseteq \wedge m \uparrow$  and  $s \sqsupseteq \wedge n \uparrow$  are mixed by  $a$  for every  $m, n \ll a$ , because  $s \sqsupseteq$  is max-separated by  $a$ . By transitivity of mixing we get that  $t \sqsupseteq \wedge m \uparrow$  and  $t \sqsupseteq \wedge n \uparrow$  are mixed by  $a$  for every  $m, n \ll a$ .

This contradicts case (c) and (d) of Lemma 2.10.  $\dashv$

LEMMA 2.30. *Let  $s, t \ll a$ . Suppose  $s$  and  $t \sqsupseteq$  are mixed by  $a$  and  $s$  is min-max-separated by  $a$ . Then  $t \sqsupseteq$  is not strongly separated by  $a$ .*

PROOF. Since  $s$  and  $t \sqsupseteq$  are mixed by  $a$ , there exist  $x, y \ll a$  such that  $\Delta(s \wedge x) = \Delta(t \sqsupseteq \wedge y)$ . Assume to the contrary that  $t \sqsupseteq$  is strongly separated by  $a$ . Two applications of Lemma 2.11 yield that  $\max(x(0)) > \min(y(0))$  and  $\min(x(0)) < \max(y(0))$ . Now we distinguish three cases.

For the first case suppose that  $\min(x(0)) < \min(y(0))$  and  $\max(x(0)) > \min(y(0))$ . Let  $v$  be the part of  $x(0)$  below  $\min(y(0))$ . Since  $\Delta(s \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , we have that  $s \wedge \langle v \rangle \uparrow$  and  $t \sqsupseteq$  are mixed by  $a$ . Moreover  $s$  and  $t \sqsupseteq$  are mixed by  $a$ . Hence by transitivity of mixing we get that  $s \wedge \langle v \rangle \uparrow$  and  $s$  are mixed by  $a$ . But this contradicts case (c) of Lemma 2.10.

Next, suppose that  $\min(x(0)) > \min(y(0))$  and  $\min(x(0)) < \max(y(0))$ . Let  $v$  be the part of  $y(0)$  below  $\min(x(0))$ . Since  $\Delta(s \wedge x) = \Delta(t \sqsupseteq \wedge y)$ , we have that  $s$  and  $t \sqsupseteq \wedge \langle v \rangle \uparrow$  are mixed by  $a$ . Moreover  $s$  and  $t \sqsupseteq$  are mixed by  $a$ . Hence by transitivity of mixing we get that  $t \sqsupseteq$  and  $t \sqsupseteq \wedge \langle v \rangle \uparrow$  are mixed by  $a$ , contradicting case (d) of Lemma 2.10.

Finally, assume that  $\min(x(0)) = \min(y(0))$ . By Lemma 2.12 we get that  $s \wedge m \uparrow$  and  $t \sqsupseteq \wedge m \uparrow$  are mixed by  $a$  for all  $m \ll a$ . Moreover by (d) of Lemma 2.7 we have that  $s \wedge m \uparrow$  and  $s \wedge n \uparrow$  are mixed by  $a$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$ , because  $s$  is min-max-separated by  $a$ . By transitivity of mixing we get that  $t \sqsupseteq \wedge m \uparrow$  and  $t \sqsupseteq \wedge n \uparrow$  are mixed by  $a$  for every  $m, n \ll a$  with  $\min(m) = \min(n)$ . This contradicts case (d) of Lemma 2.10.  $\dashv$

Now we define the parameter  $\gamma$  of the mapping  $\Gamma_\gamma$  which will canonize our given  $\Delta$ .

DEFINITION. For given canonical  $a$  define  $\gamma: (a)^{<\omega} \rightarrow \{sm, \text{min-sep}, \text{max-sep}, \text{min-max}, \text{sss}, \text{vss}\}$  as follows: Let  $\gamma(s) = sm$  iff  $s$  is strongly mixed by  $a$ ; moreover let  $\gamma(s) = \text{min-sep}$  iff  $s$  is min-separated by  $a$ ; let  $\gamma(s) = \text{max-sep}$  iff  $s$  is max-separated by  $a$ ; let  $\gamma(s) = \text{min-max}$  iff  $s$  is min-max-separated by  $a$ ; let  $\gamma(s) = \text{sss}$  iff  $s$  is still strongly separated by  $a$ ; finally, let  $\gamma(s) = \text{vss}$  iff  $s$  is very strongly separated by  $a$ .

Recall that  $\Gamma_\gamma$  is defined as follows:

For  $m \in [\omega]^{<\omega}$  let  $s \wedge m(m) = \emptyset$ ,  $\text{min-sep}(m) = \{\min(m)\}$ ,  $\text{max-sep}(m) =$

$\{\max(m)\}$ ,  $\min\text{-}\max(m) = \{\min(m), \max(m)\}$  and  $\text{sss}(m) = \text{vss}(m) = m$ .

Let  $x \in (a)^\omega$ . Define  $\Gamma_\gamma(x)$  as follows: Let  $k(0) = 0$  and  $\langle k(i): 0 < i < N \leq \omega \rangle$  increasingly enumerate those  $k$  such that  $\gamma(x \uparrow (k-1)) = \text{vss}$ . Moreover let  $k(N) = \omega$ , if  $N < \omega$ . Now let  $\Gamma_\gamma(x) = \langle \bigcup_{k(i) \leq j < k(i+1)} \gamma(x \uparrow j)(x(j)): i < N \rangle$ .

Finally, we need three more definitions in order to give our last few lemmas.

DEFINITION. Let  $x, y \ll a$  and  $k \in \omega$ .

If possible, choose  $i > 0$  maximal such that  $x(i-1) \triangleleft a(k)$ , otherwise choose  $i = 0$ . Additionally, let  $v$  denote the part of  $x(i)$  below  $\min(a(k))$ . Now define  $x_{k\downarrow}$  as follows: If  $v = \emptyset$ , let  $x_{k\downarrow} = x \uparrow i$ , otherwise let  $x_{k\downarrow} = x \uparrow i \wedge \langle v \rangle \uparrow$ .

Next, choose  $i$  minimal such that  $\min(a(k)) \leq \min(x(i))$ . Additionally if  $i > 0$ , let  $v$  denote the part of  $x(i-1)$  larger than or equal to  $\min(a(k))$ , otherwise let  $v = \emptyset$ . Now define  $x_{k\uparrow}$  as follows: If  $v = \emptyset$ , let  $x_{k\uparrow} = x \uparrow i$ , otherwise let  $x_{k\uparrow} = \langle v \rangle \wedge x \uparrow i$ .

Finally, if possible, choose  $0 < i < \text{dom}(\Gamma_\gamma(x))$  resp.  $0 < j < \text{dom}(\Gamma_\gamma(y))$  maximal such that  $\Gamma_\gamma(x)(i-1) \triangleleft a(k)$  resp.  $\Gamma_\gamma(y)(j-1) \triangleleft a(k)$ , otherwise choose  $i = 0$  resp.  $j = 0$ . Additionally, let  $v$  resp.  $w$  denote the part of  $\Gamma_\gamma(x)(i)$  resp.  $\Gamma_\gamma(y)(j)$  below  $\min(a(k))$ . Now we say that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$  iff  $\Gamma_\gamma(x) \uparrow i = \Gamma_\gamma(y) \uparrow j$  and  $v = w$ .

REMARK. By definition of  $x_{k\downarrow}$  and  $x_{k\uparrow}$  it follows that  $x_{k\downarrow} \wedge x_{k\uparrow} = x$  for every  $x \ll a$  and  $k \in \omega$ .

LEMMA 2.31. Let  $x, y \ll a$ . Suppose that  $x_{i\downarrow}$  and  $y_{i\downarrow}$  are mixed by  $a$  for every  $i < \omega$ . Then  $\Delta(x) = \Delta(y)$ .

PROOF. For every  $i < \omega$  let  $x_i, y_i \ll a$  be such that  $\Delta(x_{i\downarrow} \wedge x_i) = \Delta(y_{i\downarrow} \wedge y_i)$ . These sets exist, because  $x_{i\downarrow}$  and  $y_{i\downarrow}$  are mixed by  $a$ . Moreover by definition of  $x_{k\downarrow}$  we obtain that  $\lim_{i < \omega} x_{i\downarrow} \wedge x_i = x$  and  $\lim_{i < \omega} y_{i\downarrow} \wedge y_i = y$ . By (a) of canonical we have that  $\Delta \uparrow (a)^\omega$  is continuous. Hence we get that  $\Delta(x) = \lim_{i < \omega} \Delta(x_{i\downarrow} \wedge x_i)$  and  $\Delta(y) = \lim_{i < \omega} \Delta(y_{i\downarrow} \wedge y_i)$ . Thus,  $\lim_{i < \omega} \Delta(x_{i\downarrow} \wedge x_i)$  and  $\lim_{i < \omega} \Delta(y_{i\downarrow} \wedge y_i)$  exist. Finally, since  $\Delta(x_{i\downarrow} \wedge x_i) = \Delta(y_{i\downarrow} \wedge y_i)$  for every  $i < \omega$ , we get that  $\lim_{i < \omega} \Delta(x_{i\downarrow} \wedge x_i) = \lim_{i < \omega} \Delta(y_{i\downarrow} \wedge y_i)$ , so we are done.  $\dashv$

LEMMA 2.32. Let  $x, y \ll a$  and  $k \in \omega$ . Suppose that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ . Then  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ .

PROOF. We prove the assertion in the lemma by induction on  $k$ .

Suppose first that  $k = 0$ . By definition of  $x_{k\downarrow}$  we have that  $x_{0\downarrow} = \emptyset$  and  $y_{0\downarrow} = \emptyset$ . Thus, by definition of mixing we have that  $x_{0\downarrow}$  and  $y_{0\downarrow}$  are mixed by  $a$ .

Now assume that the assertion is true for some  $k$ . We show that it is also true for  $k+1$ . For that purpose suppose that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k+1$ . Hence  $\Gamma_\gamma(x)$  also corresponds with  $\Gamma_\gamma(y)$  up to  $k$ . By inductual assumption we

have that  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . Additionally, assume without loss of generality that  $x_{k+1\downarrow} \neq x_{k\downarrow}$  or  $y_{k+1\downarrow} \neq y_{k\downarrow}$ . We distinguish ten cases.

For the first case suppose that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are strongly mixed by  $a$ . We have that  $sm(m) = \emptyset$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we have that either  $x_{k+1\downarrow} = x_{k\downarrow}$  or  $x_{k+1\downarrow} = x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  and that either  $y_{k+1\downarrow} = y_{k\downarrow}$  or  $y_{k+1\downarrow} = y_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$ . By (a) of Lemma 2.7 we get that  $x_{k\downarrow}$  and  $x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  as well as  $y_{k\downarrow}$  and  $y_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  are mixed by  $a$ . Moreover since  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ , by transitivity of mixing we obtain that  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

Next, assume that  $x_{k\downarrow}$  is strongly mixed by  $a$ . Moreover suppose that  $y_{k\downarrow}$  is either min-separated, min-max-separated or strongly separated by  $a$ . We have that  $sm(m) = \emptyset$  as well as  $min-sep(m) = \{min(m)\}$ ,  $min-max(m) = \{min(m), max(m)\}$  and  $sss(m) = vss(m) = m$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we must have that  $x_{k+1\downarrow} = x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow}$ . By (a) of Lemma 2.7 we get that  $x_{k\downarrow}$  and  $x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  are mixed by  $a$ . Therefore, since  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ , by transitivity of mixing we obtain that  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

Now assume that  $x_{k\downarrow}$  is strongly mixed by  $a$  and  $y_{k\downarrow}$  is max-separated by  $a$ . We have that  $sm(m) = \emptyset$  and  $max-sep(m) = \{max(m)\}$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we must have that either  $x_{k+1\downarrow} = x_{k\downarrow}$  or  $x_{k+1\downarrow} = x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  and that either  $y_{k+1\downarrow} = y_{k\downarrow}$  or  $y_{k+1\downarrow} = y_{k\downarrow} \hat{\ } \langle a(k) \rangle \uparrow$ . By (a) of Lemma 2.7 we get that  $x_{k\downarrow}$  and  $x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  are mixed by  $a$ . Moreover by (c) of Lemma 2.7 we get that  $y_{k\downarrow}$  and  $y_{k\downarrow} \hat{\ } \langle a(k) \rangle \uparrow$  are mixed by  $a$ . Since  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ , by transitivity of mixing we obtain that  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

For the fourth case suppose that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are min-separated by  $a$ . We have that  $min-sep(m) = \{min(m)\}$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we must have that  $x_{k+1\downarrow} = x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$ . By Lemma 2.14 we get that  $x_{k\downarrow} \hat{\ } \langle a(k) \rangle$  and  $y_{k\downarrow} \hat{\ } \langle a(k) \rangle$  are mixed by  $a$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . Moreover by (b) of Lemma 2.7 we get that  $x_{k\downarrow} \hat{\ } \langle a(k) \rangle$  and  $x_{k\downarrow} \hat{\ } \langle a(k) \rangle \uparrow$  as well as  $y_{k\downarrow} \hat{\ } \langle a(k) \rangle$  and  $y_{k\downarrow} \hat{\ } \langle a(k) \rangle \uparrow$  are mixed by  $a$ . Therefore, possibly by transitivity of mixing, we obtain that  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

Next, assume that  $x_{k\downarrow}$  is min-separated by  $a$  and  $y_{k\downarrow}$  is min-max-separated by  $a$ . We have that  $min-sep(m) = \{min(m)\}$  and  $min-max(m) = \{min(m), max(m)\}$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we must have that  $x_{k+1\downarrow} = x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow} \hat{\ } \langle a(k) \rangle \uparrow$ . By Lemma 2.26 we get that  $x_{k\downarrow} \hat{\ } \langle a(k) \rangle \uparrow$  and  $y_{k\downarrow} \hat{\ } \langle a(k) \rangle \uparrow$  are mixed by  $a$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . Moreover by (b) of Lemma 2.7 we get that  $x_{k\downarrow} \hat{\ } \langle a(k) \rangle \uparrow$  and  $x_{k\downarrow} \hat{\ } \langle a(k) \rangle$  are mixed by  $a$ . Therefore, possibly by transitivity of mixing, we obtain that  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

We observe that if  $x_{k\downarrow}$  is min-separated by  $a$ , then by Lemma 2.28  $y_{k\downarrow}$  is neither max-separated nor strongly separated by  $a$ .

For the sixth case suppose that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are max-separated by  $a$ . We have that  $\text{max-sep}(m) = \{\text{max}(m)\}$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we must have that either  $((x_{k+1\downarrow} = x_{k\downarrow} \vee x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow) \wedge (y_{k+1\downarrow} = y_{k\downarrow} \vee y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow))$  or that  $(x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow) \wedge (y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow)$ . By (c) of Lemma 2.7 we get that  $x_{k\downarrow}$  and  $x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  as well as  $y_{k\downarrow}$  and  $y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  are mixed by  $a$ . Moreover by Lemma 2.16 we get that  $x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  are mixed by  $a$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . Therefore, possibly by transitivity of mixing, we obtain that  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

We observe that if  $x_{k\downarrow}$  is max-separated by  $a$ , then by Lemma 2.29  $y_{k\downarrow}$  is neither min-max-separated nor strongly separated by  $a$ .

For the seventh case assume that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are min-max-separated by  $a$ . We have that  $\text{min-max}(m) = \{\text{min}(m), \text{max}(m)\}$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we must have that either  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  or that  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$ . By Lemma 2.18 we get that  $x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  as well as  $x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  are mixed by  $a$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . Therefore,  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

We observe that if  $x_{k\downarrow}$  is min-max-separated by  $a$ , then by Lemma 2.30  $y_{k\downarrow}$  is not strongly separated by  $a$ .

For the eighth case suppose that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are still strongly separated by  $a$ . We have that  $\text{sss}(m) = m$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we must have that  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \square$ . By (a) of Lemma 2.27 we get that  $x_{k\downarrow} \wedge \langle a(k) \rangle \square$  and  $y_{k\downarrow} \wedge \langle a(k) \rangle \square$  are mixed by  $a$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . Therefore,  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

Next, assume that  $x_{k\downarrow}$  is still strongly separated by  $a$  and  $y_{k\downarrow}$  is very strongly separated by  $a$ . We have that  $\text{sss}(m) = \text{vss}(m) = m$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we must have that  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$ . By (b) of Lemma 2.27 we get that  $x_{k\downarrow} \wedge \langle a(k) \rangle \square$  and  $y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  are mixed by  $a$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . Therefore,  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

Finally, suppose that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are very strongly separated by  $a$ . We have that  $\text{vss}(m) = m$  for every  $m \ll a$ . Since  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k + 1$ , we must have that either  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  or that  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$ . By (c) of Lemma 2.27 we get that  $x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  are mixed by  $a$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . Moreover by Lemma 2.22 we get that  $x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  are mixed by  $a$ . Therefore,  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ .

Altogether, by symmetry we can conclude that in every case  $x_{k+1\downarrow}$  and  $y_{k+1\downarrow}$  are mixed by  $a$ . This completes the proof.  $\dashv$

LEMMA 2.33. *Let  $x, y \ll a$ . Suppose that  $\Gamma_\gamma(x) = \Gamma_\gamma(y)$ . Then  $\Delta(x) = \Delta(y)$ .*

PROOF. First, we observe that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $i$  for every  $i < \omega$ , because  $\Gamma_\gamma(x) = \Gamma_\gamma(y)$ . Hence by Lemma 2.32 we get that  $x_{i\downarrow}$  and  $y_{i\downarrow}$  are mixed by  $a$  for all  $i < \omega$ . Thus, Lemma 2.31 yields that  $\Delta(x) = \Delta(y)$ .  $\dashv$

LEMMA 2.34. *Let  $x, y \ll a$ . Suppose that  $\Gamma_\gamma(x) \neq \Gamma_\gamma(y)$ . Then  $\Delta(x) \neq \Delta(y)$ .*

PROOF. Since  $\Gamma_\gamma(x) \neq \Gamma_\gamma(y)$ , we can choose  $k$  maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ . By Lemma 2.32 we get that  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . We show that  $\Delta(x) \neq \Delta(y)$ . For that purpose we distinguish nine cases.

For the first case assume that  $x_{k\downarrow}$  is strongly mixed by  $a$ . Moreover suppose that  $y_{k\downarrow}$  is either min-separated, min-max-separated or strongly separated by  $a$ . We have that  $sm(m) = \emptyset$  as well as  $min-sep(m) = \{min(m)\}$ ,  $min-max(m) = \{min(m), max(m)\}$  and  $sss(m) = vss(m) = m$  for every  $m \ll a$ . Since  $k$  is chosen maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ , we must have that either  $x_{k+1\downarrow} = x_{k\downarrow}$  or  $x_{k+1\downarrow} = x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  and that  $y_{k+1\downarrow} = y_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$ . This implies that  $max(x_{k\uparrow}(0)) > min(y_{k\uparrow}(0))$ . Thus, by Lemma 2.23 we obtain that  $\Delta(x_{k\downarrow} \hat{\ } x_{k\uparrow}) \neq \Delta(y_{k\downarrow} \hat{\ } y_{k\uparrow})$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ .

Next, assume that  $x_{k\downarrow}$  is strongly mixed by  $a$  and  $y_{k\downarrow}$  is max-separated by  $a$ . We have that  $sm(m) = \emptyset$  and  $max-sep(m) = \{max(m)\}$  for every  $m \ll a$ . Since  $k$  is chosen maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ , we must have that either  $x_{k+1\downarrow} = x_{k\downarrow}$  or  $x_{k+1\downarrow} = x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  and that  $y_{k+1\downarrow} = y_{k\downarrow} \hat{\ } \langle a(k) \rangle$ . This implies that  $max(x_{k\uparrow}(0)) \geq max(y_{k\uparrow}(0))$ . Thus, by Lemma 2.24 we obtain that  $\Delta(x_{k\downarrow} \hat{\ } x_{k\uparrow}) \neq \Delta(y_{k\downarrow} \hat{\ } y_{k\uparrow})$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ .

We observe that we cannot have that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are strongly mixed by  $a$ . This would contradict the choice of  $k$ , because  $sm(m) = \emptyset$  for all  $m \ll a$ .

For the third case suppose that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are min-separated by  $a$ . We have that  $min-sep(m) = \{min(m)\}$  for every  $m \ll a$ . Moreover we have chosen  $k$  maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ . Therefore, by symmetry we must have without loss of generality that  $x_{k+1\downarrow} = x_{k\downarrow}$  and  $y_{k+1\downarrow} = y_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$ . This implies that  $min(x_{k\uparrow}(0)) > min(y_{k\uparrow}(0))$ . Thus, by Lemma 2.13 we obtain that  $\Delta(x_{k\downarrow} \hat{\ } x_{k\uparrow}) \neq \Delta(y_{k\downarrow} \hat{\ } y_{k\uparrow})$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ .

Next, assume that  $x_{k\downarrow}$  is min-separated by  $a$  and  $y_{k\downarrow}$  is min-max-separated by  $a$ . We have that  $min-sep(m) = \{min(m)\}$  and  $min-max(m) = \{min(m), max(m)\}$  for every  $m \ll a$ . Since  $k$  is chosen maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ , we must have that either  $x_{k+1\downarrow} = x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow}$ ,  $x_{k+1\downarrow} = x_{k\downarrow}$  and  $y_{k+1\downarrow} = y_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  or that  $x_{k+1\downarrow} = x_{k\downarrow} \hat{\ } \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow} \hat{\ } \langle a(k) \rangle$ . This implies that  $min(x_{k\uparrow}(0)) \neq min(y_{k\uparrow}(0))$  or  $max(x_{k\uparrow}(0)) \geq max(y_{k\uparrow}(0))$ . Thus, by Lemma 2.25 we obtain that  $\Delta(x_{k\downarrow} \hat{\ } x_{k\uparrow}) \neq \Delta(y_{k\downarrow} \hat{\ } y_{k\uparrow})$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ .

We observe that if  $x_{k\downarrow}$  is min-separated by  $a$ , then by Lemma 2.28  $y_{k\downarrow}$  is neither max-separated nor strongly separated by  $a$ .

For the fifth case suppose that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are max-separated by  $a$ . We have that  $max-sep(m) = \{max(m)\}$  for every  $m \ll a$ . Moreover we have chosen  $k$

maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ . Therefore, by symmetry we must have without loss of generality that either  $x_{k+1\downarrow} = x_{k\downarrow}$  or  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and that  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle$ . This implies that  $\max(x_{k\uparrow}(0)) > \max(y_{k\uparrow}(0))$ . Thus, by Lemma 2.15 we obtain that  $\Delta(x_{k\downarrow} \wedge x_{k\uparrow}) \neq \Delta(y_{k\downarrow} \wedge y_{k\uparrow})$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ .

We observe that if  $x_{k\downarrow}$  is max-separated by  $a$ , then by Lemma 2.29  $y_{k\downarrow}$  is neither min-max-separated nor strongly separated by  $a$ .

For the sixth case assume that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are min-max-separated by  $a$ . We have that  $\min\text{-max}(m) = \{\min(m), \max(m)\}$  for every  $m \ll a$ . Moreover we have chosen  $k$  maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ . Therefore, by symmetry we must have without loss of generality that either  $x_{k+1\downarrow} = x_{k\downarrow}$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \square$  or that  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle$ . This implies that  $\min(x_{k\uparrow}(0)) > \min(y_{k\uparrow}(0))$  or  $\max(x_{k\uparrow}(0)) > \max(y_{k\uparrow}(0))$ . Thus, by Lemma 2.17 we obtain that  $\Delta(x_{k\downarrow} \wedge x_{k\uparrow}) \neq \Delta(y_{k\downarrow} \wedge y_{k\uparrow})$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ .

We observe that if  $x_{k\downarrow}$  is min-max-separated by  $a$ , then by Lemma 2.30  $y_{k\downarrow}$  is not strongly separated by  $a$ .

For the seventh case suppose that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are still strongly separated by  $a$ . We have that  $\text{sss}(m) = m$  for every  $m \ll a$ . Moreover we have chosen  $k$  maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ . Therefore, by symmetry we must have without loss of generality that  $x_{k+1\downarrow} = x_{k\downarrow}$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \square$ . This implies that neither  $x_{k\uparrow}(0)$  is an initial segment of  $y_{k\uparrow}(0)$  nor conversely. Thus, by Lemma 2.19 we obtain that  $\Delta(x_{k\downarrow} \wedge x_{k\uparrow}) \neq \Delta(y_{k\downarrow} \wedge y_{k\uparrow})$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ .

Next, assume that  $x_{k\downarrow}$  is still strongly separated by  $a$  and  $y_{k\downarrow}$  is very strongly separated by  $a$ . We have that  $\text{sss}(m) = \text{vss}(m) = m$  for every  $m \ll a$ . Since  $k$  is chosen maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ , we must have that either  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow}$ ,  $x_{k+1\downarrow} = x_{k\downarrow}$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle \square$  or that  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle$ . The former two cases imply that neither  $x_{k\uparrow}(0)$  is an initial segment of  $y_{k\uparrow}(0)$  nor conversely. Thus, by Lemma 2.19 we obtain that  $\Delta(x_{k\downarrow} \wedge x_{k\uparrow}) \neq \Delta(y_{k\downarrow} \wedge y_{k\uparrow})$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . In the latter case by (b) of Lemma 2.27 we get that  $x_{k\downarrow} \wedge \langle a(k) \rangle \square$  and  $y_{k\downarrow} \wedge \langle a(k) \rangle$  are separated by  $a$ . Therefore, by definition of separation we obtain that  $\Delta(x_{k+1\downarrow} \wedge x_{k+1\uparrow}) \neq \Delta(y_{k+1\downarrow} \wedge y_{k+1\uparrow})$ .

Finally, suppose that both  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are very strongly separated by  $a$ . We have that  $\text{vss}(m) = m$  for every  $m \ll a$ . Moreover we have chosen  $k$  maximal such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ . Therefore, by symmetry we must have without loss of generality that either  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \square$  and  $y_{k+1\downarrow} = y_{k\downarrow}$  or that  $x_{k+1\downarrow} = x_{k\downarrow} \wedge \langle a(k) \rangle \uparrow$  and  $y_{k+1\downarrow} = y_{k\downarrow} \wedge \langle a(k) \rangle$ . This implies that  $x_{k\uparrow}(0) \neq y_{k\uparrow}(0)$ . Thus, by Lemma 2.21 we obtain that  $\Delta(x_{k\downarrow} \wedge x_{k\uparrow}) \neq \Delta(y_{k\downarrow} \wedge y_{k\uparrow})$ , because  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ .

Altogether, by symmetry we can conclude that in every case  $\Delta(x) \neq \Delta(y)$ . This completes the proof.  $\dashv$

Both the following definition and Lemma 2.35 are necessary to guarantee that property (b) of Theorem 0.3.8 follows from our Main Theorem.

DEFINITION. Let  $x, y \ll a$ . We say that  $\Gamma_\gamma(x)$  is a *proper initial segment* of  $\Gamma_\gamma(y)$  iff there exists  $k \in \omega$  such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ ,  $x_{j\downarrow}$  is strongly mixed by  $a$  for every  $j \geq k$  and there exists  $l \geq k$  such that  $y_{l\downarrow}$  is separated in some sense by  $a$ .

LEMMA 2.35. *There are no  $x, y \ll a$  such that  $\Gamma_\gamma(x)$  is a proper initial segment of  $\Gamma_\gamma(y)$ .*

PROOF. Assume to the contrary that there exist  $x, y \ll a$  such that  $\Gamma_\gamma(x)$  is a proper initial segment of  $\Gamma_\gamma(y)$ . According to the definition above there exists  $k \in \omega$  such that  $\Gamma_\gamma(x)$  corresponds with  $\Gamma_\gamma(y)$  up to  $k$ ,  $x_{j\downarrow}$  is strongly mixed by  $a$  for every  $j \geq k$  and there exists  $l \geq k$  such that  $y_{l\downarrow}$  is separated in some sense by  $a$ . By Lemma 2.32 we get that  $x_{k\downarrow}$  and  $y_{k\downarrow}$  are mixed by  $a$ . Hence by definition of being mixed there exist  $x_0, y_0 \ll a$  such that  $\Delta(x_{k\downarrow} \hat{\ } x_0) = \Delta(y_{k\downarrow} \hat{\ } y_0)$ . Since  $x_{j\downarrow}$  is strongly mixed by  $a$  for all  $j \geq k$ , by (e) of canonical we have that  $(x_{k\downarrow} \hat{\ } x_0)_{j\downarrow}$  is also strongly mixed by  $a$  for every  $j \geq k$ . Equally, since there exists  $l \geq k$  such that  $y_{l\downarrow}$  is separated in some sense by  $a$ , by (e) of canonical there exists  $i \geq k$  such that  $(y_{k\downarrow} \hat{\ } y_0)_{i\downarrow}$  is also separated in some sense by  $a$ . Hence by definition of  $\Gamma_\gamma$  we have  $\Gamma_\gamma(x_{k\downarrow} \hat{\ } x_0) \neq \Gamma_\gamma(y_{k\downarrow} \hat{\ } y_0)$ . Since  $\Delta(x_{k\downarrow} \hat{\ } x_0) = \Delta(y_{k\downarrow} \hat{\ } y_0)$ , we get a contradiction to Lemma 2.34.  $\dashv$

DEFINITION. Let  $x \ll a$ . We call  $\Gamma_\gamma(x)$  *finite* iff there exists  $k \in \omega$  such that  $x_{j\downarrow}$  is strongly mixed by  $a$  for every  $j \geq k$ . Otherwise we call  $\Gamma_\gamma(x)$  *infinite*.

LEMMA 2.36. *If  $\Delta(\Omega^\omega)$  is countable, then  $\Gamma_\gamma(x)$  is finite for every  $x \ll a$ .*

PROOF. By (f) of canonical we have that either  $\Gamma_\gamma(x)$  is finite for all  $x \ll a$  or that  $\Gamma_\gamma(x)$  is infinite for every  $x \ll a$ . Assume to the contrary that  $\Gamma_\gamma(x)$  is infinite for all  $x \ll a$ .

Inductively, we construct  $s_f \ll a$  for every  $f \in \omega^{<\omega}$ , where  $\omega^{<\omega}$  denotes the set of all finite sequences of nonnegative integers. Moreover for every  $i \in \omega$  let  $\omega^i$  be the set of all sequences of  $i$  nonnegative integers. Put  $s_\emptyset = \emptyset$  and suppose that for some  $j \in \omega$  and all  $i \leq j$  the sets  $s_f$  have already been constructed for every  $f \in \omega^i$ . First, for every  $f \in \omega^j$  let  $l_f \in \omega$  be minimal such that  $s_f < a \upharpoonright l_f$ . Therewith we put  $a_f = a \upharpoonright l_f$ . Moreover for all  $f \in \omega^j$  let  $k_f \in \omega$  be minimal such that  $s_f \hat{\ } (a_f \upharpoonright k_f)$  is separated in some sense by  $a$ . These  $k_f$  exist, as  $\Gamma_\gamma(x)$  is infinite for each  $x \ll a$ . Now put  $s_{f \hat{\ } \langle i \rangle} = s_f \hat{\ } (a_f \upharpoonright k_f) \hat{\ } \langle a(k_f + i) \rangle$  for every  $i \in \omega$ . This completes the construction.



Finally, let  $x_F = \bigcup_{i \in \omega} s_{F \wedge i}$  for every  $F \in \omega^\omega$ . Clearly, we have  $x_F \ll a$  for each such  $F$ . Moreover by definition of  $\Gamma_\gamma$  it holds that  $\Gamma_\gamma(x_F) \neq \Gamma_\gamma(x_G)$  for all  $F, G \in \omega^\omega$  with  $F \neq G$ . By Lemma 2.34 we get a contradiction to the fact that  $\Delta(\Omega^\omega)$  is countable.

This completes the proof of the lemma and with it the proof of the Main Theorem. ◻

# A. COROLLARIES

First, we prove that a finite-dimensional version of Theorem 0.3.5 of Taylor follows from our Main Theorem.

**COROLLARY.** *Let  $k > 0$ . For every mapping  $f: \Omega^k \rightarrow \omega$  there exist  $\gamma: \Omega^{<\omega} \rightarrow \{sm, \text{min-sep}, \text{max-sep}, \text{min-max}, \text{sss}, \text{vss}\}$  and  $a \in \Omega^\omega$  such that the following properties hold.*

(a) *For no  $s \in (a)^{k-1}$  we have that  $\gamma(s) = \text{sss}$ .*

(b) *If  $i \geq k$ , then  $\gamma(x \upharpoonright i) = sm$ .*

(c) *For every  $x, y \ll a$  it follows that  $f(x \upharpoonright k) = f(y \upharpoonright k)$  iff  $\Gamma_\gamma(x) = \Gamma_\gamma(y)$ .*

**PROOF.** We define  $\Delta: \Omega^\omega \rightarrow \omega$  by  $x \mapsto f(x \upharpoonright k)$ . First, we show that  $\Delta$  is a Borel measurable mapping. By construction of  $\Delta$  we have that  $\Delta^{-1}(i) = \cup \{(s, \omega_{\text{max}})^\omega: s \in \Omega^k \wedge f(s) = i\}$  for every  $i \in \omega$ . Hence  $\Delta$  is even continuous.

Thus, we can apply our Main Theorem to  $\Delta$  and get  $a \in \Omega^\omega$  such that  $\Delta(x) = \Delta(y)$  iff  $\Gamma_\gamma(x) = \Gamma_\gamma(y)$  for all  $x, y \ll a$ . Again, by construction of  $\Delta$  we have that  $x \upharpoonright i$  is strongly mixed by  $a$  for every  $i \geq k$ . Therefore it remains to show that property (a) of our corollary holds.

For that purpose assume to the contrary that  $s$  is still strongly separated by  $a$  for some  $s \in (a)^{k-1}$ . Hence we have that  $s \wedge m$  and  $s \wedge m \uparrow$  are mixed by  $a$  for every  $m \ll a$ . This implies that for all  $m \ll a$  there exist  $x, y \ll a$  such that  $\Delta(s \wedge m \wedge x) = \Delta(s \wedge m \uparrow \wedge y)$ . By Lemma 2.20 we get that  $s \wedge m \uparrow$  is still strongly separated by  $a$  for every  $m \ll a$ . Moreover we have that  $s \wedge m \in (a)^k$ . Thus, by property (b) we get a contradiction to Lemma 2.35.  $\dashv$

We conclude our work by showing that our Main Theorem implies Theorem 0.3.8 of Prömel and Voigt.

**PROOF.** Assuming a Borel function  $\Delta: [\omega]^\omega \rightarrow \mathbb{R}$  we construct the mapping  $\Delta^*: \Omega^\omega \rightarrow \mathbb{R}$  with  $x \mapsto \Delta(\{\min(x(i)): i \in \omega\})$ . In order to apply our Main Theorem we have to prove that  $\Delta^*$  is Borel. Let  $g: \Omega^\omega \rightarrow [\omega]^\omega$  with  $g(x) = \{\min(x(i)): i \in \omega\}$ . Since  $\Delta^* = \Delta \circ g$  and  $\Delta$  is Borel, it is enough to show that  $g$  is Borel. Let  $\wp(\omega)$  denote the set of all subsets of  $\omega$ .  $\wp(\omega)$  can be identified with the Cantor space  $2^\omega$  as a topological space endowed with the product topology. Since  $[\omega]^\omega \subseteq \wp(\omega)$ , for every  $I, J \in [\omega]^{<\omega}$  with  $I \cap J = \emptyset$  the sets  $U_{I, J} = \{X \in [\omega]^\omega: \forall i \in I \forall j \in J i \in X \wedge j \notin X\}$  form a basis for the topology on  $[\omega]^\omega$ . It is obvious that the (sub)basis is countable, so it suffices to show that  $g^{-1}(U_{I, J})$  is Borel for each  $I, J$ . We have that  $g^{-1}(U_{I, J}) = \{x \in \Omega^\omega: \{\min(x(i)): i \in \omega\} \in U_{I, J}\}$ . The sets  $\{\prod_{i \in \omega} V_i: \forall i \in \omega V_i \subseteq [\omega]^{<\omega} \wedge V_i = [\omega]^{<\omega}$  for all but finitely many  $i\} \cap \Omega^\omega$  form a basis for  $\Omega^\omega$ . Since only a finite number of pieces  $x(i)$  consider the sets  $I, J$ ,  $g^{-1}(U_{I, J})$  is a

union of open sets of  $\Omega^\omega$ . Hence  $g^{-1}(U_{i, j})$  is open, too. Therewith  $g$  is continuous. Since continuous mappings are Borel,  $g$  is also Borel.

For given  $a$  let  $x \ll a$ . Assume that for some  $k$  the set  $x \upharpoonright k$  is max-separated, min-max-separated or strongly separated by  $a$ . We have that  $x \upharpoonright k \wedge \langle x(k) \rangle$  and  $x \upharpoonright k \wedge \langle x(k) \cup x(k+1) \rangle$  are mixed by  $a$ , since  $\langle x(k) \rangle$  and  $\langle x(k) \cup x(k+1) \rangle$  have the same minimum and hence  $\Delta^*(x \upharpoonright k \wedge \langle x(k) \rangle \wedge y) = \Delta^*(x \upharpoonright k \wedge \langle x(k) \cup x(k+1) \rangle \wedge y)$  for all  $y \ll a$ . But this contradicts the cases (b) - (d) of Lemma 2.10. Thus, for all  $k$  we neither have that  $x \upharpoonright k$  is max-separated, min-max-separated, still strongly separated nor very strongly separated by  $a$ .

Therewith our Main Theorem yields  $\gamma^*: \Omega^{<\omega} \rightarrow \{sm, min-sep\}$  and  $a \in \Omega^\omega$  such that for all  $x, y \ll a$  it holds that  $\Delta^*(x) = \Delta^*(y)$  iff  $\Gamma_{\gamma^*}(x) = \Gamma_{\gamma^*}(y)$ . Let  $A = \{min(a(i)): i \in \omega\}$  and define for every  $x \ll a$  the mapping  $\Gamma: [A]^\omega \rightarrow [A]^{\leq\omega}$  by  $\Gamma(\{min(x(i)): i \in \omega\}) := \Gamma_{\gamma^*}(x)$ .

By definition of  $\Gamma_{\gamma^*}$  we get that  $\Gamma(X) \subseteq X$  for all  $X \in [A]^\omega$ .

Additionally, Theorem 0.3.8 requires that no  $\Gamma(X)$  is a proper initial segment of some  $\Gamma(Y)$ . This property directly follows from Lemma 2.35.

Finally, since both  $\Delta^*$  and  $\Gamma_{\gamma^*}$  only depend on the minima of all pieces, for all  $X, Y \in [A]^\omega$  it follows that  $\Delta(X) = \Delta(Y)$  iff  $\Gamma(X) = \Gamma(Y)$ .  $\dashv$

## B. ERKLÄRUNG

Hiermit erkläre ich, daß die von mir eingereichte Dissertation mit dem Titel „Canonical Forms of Borel Functions on the Milliken Space“ – abgesehen von der Beratung durch Herrn Prof. Dr. Otmar Spinas – nach Inhalt und Form meine eigene Arbeit ist. Sie hat weder ganz noch zum Teil einer anderen Stelle im Rahmen eines Prüfungsverfahrens vorgelegen.

Kiel, den 18.04.02

Olaf Klein

# C. LEBENS LAUF

## **Persönliche Daten**

*Name:* *Olaf Klein*  
*Geburtsdatum, -ort:* *25.01.1974 in Hamburg*  
*Anschrift:* *Johann-Fleck-Straße 41*  
*24106 Kiel*  
*Familienstand:* *ledig*  
*Staatsangehörigkeit:* *deutsch*  
*Namen der Eltern:* *Johannes Klein*  
*Bärbel Klein, geb. Zimmermann*

## **Schulbildung**

*1979-1984* *Besuch der Grundschule in Hamburg-Ochsenwerder*  
*1984-1993* *Besuch des Gymnasiums „Klosterschule“*  
*in Hamburg-Berliner Tor*  
*Abiturnote: 1,3*

## **Studium**

*1993-1999* *Studium an der Universität Hamburg*  
*Studienziel: Diplom Mathematik*  
*Abschluß: Diplom-Mathematiker*  
*Studiendauer: 12 Semester*  
*Abschlußnote: „gut“*  
*2000* *Beginn der Promotion*  
*an der Christian-Albrechts-Universität zu Kiel*  
*unter der Betreuung von Prof. Dr. Otmar Spinas*

## D. BIBLIOGRAPHY

- [Ra30] RAMSEY, F. P.: *On a problem of formal logic*, *Proceedings of the London Mathematical Society* (2), vol. 30 (1930), pp. 264-286
- [ErRa50] ERDÖS, P.; RADO, R.: *A combinatorial theorem*, *J. London Math. Soc.*, vol. 25 (1950), pp. 249-255
- [Ku66] KURATOWSKI, K.: *Topology*, *Academic Press New York*, vol. 1 (1966)
- [Ma68] MATHIAS, A. R. D.: *On a generalization of Ramsey's theorems*, *Notices of the American Mathematical Society*, vol. 15 (1968), p. 931
- [Si70] SILVER, J.: *Every analytic set is Ramsey*, *The Journal of Symbolic Logic*, vol. 35 (1970), nr. 1, pp. 60-64
- [GrRo71] GRAHAM, R. L.; ROTHSCHILD, B. L.: *Ramsey's theorem for  $n$ -parameter sets*, *Trans. Amer. Math. Soc.*, vol. 159 (1971), pp. 413-432
- [GaPr73] GALVIN, F.; PRIKRY, K.: *Borel Sets and Ramsey's Theorem*, *The Journal of Symbolic Logic*, vol. 38 (1973), nr. 2, pp. 193-198
- [Ba74] BAUMGARTNER, J. E.: *A short proof of Hindman's Theorem*, *Journal of Combinatorial Theory* (A), vol. 17 (1974), pp. 384-386
- [El74] ELLENTUCK, E.: *A new proof that analytic sets are Ramsey*, *The Journal of Symbolic Logic*, vol. 39 (1974), nr. 1, pp. 163-165
- [Hi74] HINDMAN, N.: *Finite sums from sequences within cells of a partition of  $\mathbb{N}$* , *Journal of Combinatorial Theory* (A), vol. 17 (1974), pp. 1-11
- [Mi75] MILLIKEN, K. R.: *Ramsey's Theorem with Sums or Unions*, *Journal of Combinatorial Theory* (A), vol. 18 (1975), pp. 276-290
- [Ta76] TAYLOR, A. D.: *A Canonical Partition Relation for Finite Subsets of  $\omega$* , *Journal of Combinatorial Theory* (A), vol. 21 (1976), pp. 137-146
- [PuRö82] PUDLAK, P.; RÖDL, V.: *Partition theorems for systems of finite subsets of integers*, *Discrete Math.*, vol. 39 (1982), pp. 67-73
- [PrVo85] PRÖMEL, H. J.; VOIGT, B.: *Canonical Forms of Borel-Measurable Mappings  $\Delta: [\omega]^\omega \rightarrow \mathbb{R}$* , *Journal of Combinatorial Theory* (A), vol. 40 (1985), pp. 409-417
- [Ke95] KECHRIS, A. S.: *Classical Descriptive Set Theory*, *Springer-Verlag Berlin*, 1995

- [To98] TODORCEVIC, S.: *Infinite-Dimensional Ramsey Theory, Preprint*, (1998)

# E. SYMBOLS

$\emptyset$	<i>empty set</i>
$\mathbb{R}$	<i>real numbers</i>
$\Delta$	<i>Borel measurable <math>\Delta: \Omega^\omega \rightarrow \mathbb{R}</math></i> <b>5</b>
$\omega$	<i>natural numbers</i> <b>5</b>
$[X]^\kappa$	<i>set of all subsets of <math>X</math> with the cardinality <math>\kappa</math></i> <b>5</b>
$[X]^{<\omega}$	<i>set of all finite subsets of <math>X</math></i> <b>5</b>
$[X]^{\leq\omega}$	<i>set of all countable subsets of <math>X</math></i> <b>5</b>
$A \triangleleft B$	<b>8</b>
$[a, A]^\omega$	<b>8</b>
$\Omega^\omega$	<i>Milliken space</i> <b>10</b>
$\Omega^k, \Omega^{<k}, \Omega^{\leq k}$	<b>10</b>
$\omega_{max}$	<i>finest mapping on the Milliken space</i> <b>10</b>
$dom(s)$	<i>domain of <math>s</math></i> <b>10</b>
$ran(s)$	<i>range of <math>s</math></i> <b>10</b>
$(a)^k$	<b>10</b>
$(a)^{<k}, (a)^{\leq k}$	<b>10</b>
$(s, a)^\omega$	<i>basic open sets of the Milliken space</i> <b>10</b>
$s \ll t, s \ll b, a \ll b$	<i><math>s, a</math> coarser than <math>t, b</math></i> <b>10</b>
$s < t, s < b$	<b>10</b>
$s \hat{\ } t, s \hat{\ } a$	<i>concatenation of <math>s</math> and <math>t, a</math></i> <b>10</b>
$a \upharpoonright k, a \upharpoonright k$	<i>restriction of <math>a</math> to <math>k</math></i> <b>10</b>
$s \uparrow \hat{\ } t, s \uparrow \hat{\ } x$	<b>15</b>
$s \uparrow$	<b>15</b>
$s \square$	<i>variable for <math>s</math> or <math>s \uparrow</math></i> <b>15</b>
$sm, sm(m)$	<i>strongly mixed</i> <b>5, 34</b>
$min-sep, min-sep(m)$	<i>min-separated</i> <b>5, 34</b>
$max-sep, max-sep(m)$	<i>max-separated</i> <b>5, 34</b>
$min-max, min-max(m)$	<i>min-max-separated</i> <b>5, 34</b>
$sss, sss(m)$	<i>still strongly separated</i> <b>5, 34</b>
$vss, vss(m)$	<i>very strongly separated</i> <b>5, 34</b>
$\gamma$	<b>5, 34</b>
$\Gamma_\gamma(x)$	<i>canonical mapping</i> <b>5, 34</b>
$x_{k\downarrow}$	<b>35</b>
$x_{k\uparrow}$	<b>35</b>



# F. INDEX

*accepts* 11

*Baire property* 13

*canonical* 22

*Cantor space* 7

*closure* 12

*compatible* 18

*completely decided* 20

*completely H-Ramsey* 11

*completely Ramsey* 8

*corresponds* 35

*decided* 15

*decides* 11

*Ellentuck topology* 8

*finite* 40

*H-Ellentuck topology* 10

*H-Ramsey* 11

*H-Ramsey null* 12

*infinite* 40

*interior* 13

*max-separated* 20

*meager* 12

*metric topology* 7, 14

*Milliken space* 10

*min-max-separated* 20

*min-separated* 20

*mixed* 15

*nowhere dense* 12

*pigeon-hole principle* 6

*proper initial segment* 40

*Ramsey* 7

*rejects* 11

*separated* 15

*separated in some sense* 20

*Souslin operation* 14

*Souslin scheme* 14

*still strongly separated* 29

*strongly mixed* 20

*strongly separated* 20

*very compatible* 19

*very strongly separated* 29