

Leibniz-Institut für Meereswissenschaften

N₂O
in the Atlantic Ocean and the Baltic Sea

Dissertation
zur Erlangung des Doktorgrades
der Mathematisch-Naturwissenschaftlichen Fakultät
der Christian-Albrechts-Universität zu Kiel

vorgelegt von
Sylvia Walter

Kiel, 2005

Hiermit erkläre ich, dass ich die vorliegende Doktorarbeit selbständig und ohne unerlaubte Hilfen erstellt habe. Ferner habe ich weder diese noch eine ähnliche Arbeit an einer anderen Abteilung oder Hochschule im Rahmen eines Prüfungsverfahrens vorgelegt, veröffentlicht oder zur Veröffentlichung vorgelegt.

(Sylvia Walter)

Referentin: Prof. Dr. Karin Lochte
Korreferent: Prof. Dr. Douglas Wallace

Tag der mündlichen Prüfung : 27.01.2006
Zum Druck genehmigt : 08.02.2006

This thesis is based on the following manuscripts:

Walter, S., H.W. Bange, U. Breitenbach, and D.W.R. Wallace, Nitrous oxide in the North Atlantic Ocean, 2005.

(manuscript)

Walter, S., H.W. Bange, and D.W.R. Wallace, Nitrous oxide in the surface layer of the tropical North Atlantic Ocean along a West to East transect, *Geophysical Research Letters*, 31, L23S07, doi:10.1029/2004GL019937, 2004.

Walter, S., I. Peeken, K. Lochte, and H.W. Bange, Nitrous oxide measurements during EIFEX, the European Iron Fertilisation Experiment in the subpolar South Atlantic Ocean, 2005. *(accepted, Geophysical Research Letters)*

Walter, S., U. Breitenbach, H.W. Bange, G. Nausch, and D.W.R. Wallace, Nitrous oxide water column distribution during the transition from anoxic to oxic conditions in the Baltic Sea, 2005.

(manuscript)

Walter, S., J. Suling, M. Tank, H.W. Bange, and J.F. Imhoff, Analysis of bacterial communities in the subtropical North Atlantic by 16S rRNA genes, 2005.

(manuscript)

Walter, S. and H.W. Bange, Influence of HgCl₂-poisoning and temperature on N₂O-concentrations during storage, 2005.

(manuscript)

Contribution of authors to the manuscripts

Nitrous oxide in the North Atlantic Ocean, 2005.

Sylvia Walter took the samples and made most of the measurements, evaluated the data and wrote the manuscript. Ulrich Breitenbach supported the measurements. Hermann W. Bange and Douglas W.R. Wallace assisted with input to the manuscript and revision.

Nitrous oxide in the surface layer of the tropical North Atlantic Ocean along a West to East transect, 2004.

Sylvia Walter made the measurements and evaluated the data. Hermann W. Bange and Sylvia Walter wrote the paper, reviewed by Douglas W.R. Wallace.

Nitrous oxide measurements during EIFEX, the European Iron Fertilisation Experiment in the subpolar South Atlantic Ocean, 2005.

Sylvia Walter made most of the measurements and evaluated the data. Hermann W. Bange and Sylvia Walter wrote the paper, supported by Ilka Peeken and Karin Lochte.

Nitrous oxide water column distribution during the transition from anoxic to oxic conditions in the Baltic Sea, 2005.

Sylvia Walter took the samples and made the measurements, evaluated the data and wrote the paper. Ulrich Breitenbach supported the measurements. Hermann W. Bange and Douglas W.R. Wallace assisted with input to the manuscript and revision. Günther Nausch made the participation to the cruise into the Baltic Sea possible and provided the physical and chemical data.

Analysis of bacterial communities in the subtropical North Atlantic by 16S rRNA genes, 2005.

Sylvia Walter took the samples, made the experimental work, and wrote the manuscript. Jörg Süling and Marcus Tank assisted with the experimental work and input to the manuscript.

Influence of HgCl₂-poisoning and temperature on N₂O-concentrations during storage, 2005.

Sylvia Walter designed the experiments, made the measurements, evaluated the data and wrote the manuscript, reviewed by Hermann W. Bange.

Table of Contents

<u>Summary</u>	11
<u>Zusammenfassung</u>	12
<u>Chapter 1</u>	13
Introduction	
1. Nitrous oxide (N₂O)	
1.1. Climatic relevance of N₂O	
1.2. Biological production of N₂O	
1.2.1. Nitrification	
1.2.2. Denitrification	
1.3. Sources and sinks of N₂O	
1.4. N₂O and Oceans	
2. Study areas and short compilation of N₂O data	
2.1. The Atlantic Ocean	
2.2. The Baltic Sea	
3. Thesis outline	
References	
<u>Chapter 2</u>	31
Nitrous oxide in the North Atlantic Ocean	
<u>Chapter 3</u>	57
Nitrous oxide in the surface layer of the tropical North Atlantic Ocean along a West to East transect	
<u>Chapter 4</u>	71
Nitrous oxide measurements during EIFEX, the European Iron Fertilization Experiment in the subpolar South Atlantic Ocean	
<u>Chapter 5</u>	83
Nitrous oxide water column distribution during the transition from anoxic to oxic conditions in the Baltic Sea	
<u>Chapter 6</u>	113
Analysis of bacterial communities in the subtropical North Atlantic by 16S rRNA genes	
<u>Chapter 7</u>	143
Influence of HgCl₂-poisoning and temperature on N₂O-concentrations during storage	
<u>Acknowledgements</u>	151

Für meine Oma, meine Mutter, Kerstin und Roland!
Die vier Menschen, die ich am meisten liebe und denen
ich alles zu verdanken habe!

Aim of this study was the investigation of the vertical distribution of N_2O and the factors influencing this distribution, particularly in the North Atlantic Ocean and the Baltic Sea. Dissolved and atmospheric N_2O was measured during several cruises. These data, in combination with physical and chemical parameters, were used to determine sources and sinks of N_2O . Furthermore, possible production pathways of N_2O were investigated using the relationship between N_2O and relevant production parameters, and their relation to physical processes in the oceans. The hypothesis whether N_2O concentrations increased by oceanic iron fertilization via enhanced substrate supply for microbial respiration, was tested during the European Iron Fertilization Experiment (EIFEX) in the sub-polar South Atlantic. First investigations about the vertical structure of bacterial communities in the subtropical North Atlantic were started, with the intention to find correlations between the community structure of bacteria and the distribution of N_2O . This work presents a detailed picture of the distribution of N_2O in the North Atlantic Ocean and the Baltic Sea. In summary, this study adds 80 profiles (1482 data points) of the Atlantic Ocean, and 26 profiles (231 data points) of the Baltic Sea to the database.

Atlantic Ocean

We could show that the production of N_2O in the subpolar North Atlantic is negligible, whereas the tropical North Atlantic is a net source of atmospheric N_2O , with a pronounced regional variability. The data indicate a production of N_2O by nitrification. However, the contribution of autotrophic nitrifiers to the bacterial community seems to be low, therefore a production of N_2O by others than autotrophic nitrifiers should be taken into account. In contrast to other authors the hypothesis of additional N_2O production triggered during Fe fertilization could not be confirmed.

Baltic Sea

N_2O in the Baltic Sea was measured after a major inflow of North Sea water. We could show that the inflow event has had a strong influence on the production and conversion of N_2O . The inflow led to a stimulation of the production of N_2O mainly by nitrification, but due to the clear water column stratification the Baltic Sea is not expected to be a source of atmospheric N_2O .

Ziel dieser Arbeit war die Untersuchung der vertikalen Verteilung von N_2O und den Faktoren, die diese Verteilung beeinflussen, insbesondere im Atlantik und der Ostsee. Auf mehreren Fahrten wurden sowohl atmosphärische als auch gelöste N_2O -Konzentrationen gemessen. Mit Hilfe dieser Daten, unterstützt durch physikalische und chemische Messdaten, wurden Quellen und Senken für N_2O bestimmt. Weiterhin wurden mögliche Produktionswege von N_2O mittels Korrelationsanalysen zwischen N_2O und produktionsrelevanten Parametern sowie deren Beziehung zu physikalischen Prozessen untersucht. Die Hypothese, ob eine erhöhte Biomasse-Produktion während einer Fe-Düngung zu einer erhöhten N_2O -Produktion führt, war Ziel der Untersuchungen während des European Iron Fertilization Experiments (EIFEX) im subpolaren Südatlantik. Außerdem wurden erste Untersuchungen zur vertikalen Struktur von Bakteriengemeinschaften im subtropischen Atlantik durchgeführt, um mögliche Korrelationen zwischen der Gemeinschaftsstruktur und den gemessenen N_2O -Konzentrationen zu finden. Die im Rahmen dieser Arbeit durchgeführten Studien liefern ein detailliertes Bild über die Vertikalstruktur der N_2O -Verteilung im Nordatlantik und der Ostsee. Im Atlantik sind somit 80 weitere N_2O -Tiefenprofile (1482 Datenpunkte) verfügbar, in der Ostsee weitere 26 (231 Datenpunkte).

Atlantischer Ozean

Wir konnten zeigen, dass die Produktion von N_2O im subpolaren Nordatlantik zu vernachlässigen ist, wohingegen der tropische Nordatlantik eine Quelle für atmosphärisches N_2O mit einer ausgeprägten regionalen Variabilität darstellt. Die Daten deuten auf eine N_2O -Produktion durch nitrifizierende Prozesse. Allerdings scheint der Anteil an autotrophen Nitrifizierern innerhalb der untersuchten Bakteriengemeinschaften gering zu sein, daher sollte eine N_2O -Produktion durch andere Organismen in Betracht gezogen werden. Die Hypothese einer erhöhten N_2O -Produktion ausgelöst durch Fe-Düngung konnten wir im Gegensatz zu anderen Autoren nicht bestätigen.

Ostsee

N_2O -Konzentrationen in der Ostsee wurden nach einem starken Nordseewassereinstrom gemessen. Wir konnten zeigen, dass dieser Einstrom einen starken Einfluss auf die Produktion und Umsetzung von N_2O hatte. Der Einstrom führte zu einer Stimulierung der N_2O -Produktion, wobei hauptsächlich nitrifizierende Prozesse eine Rolle spielen dürften. Aufgrund der deutlichen Stratifizierung der Ostsee wird diese nicht als Quelle für atmosphärisches N_2O angenommen.

Chapter **1**

Introduction and thesis outline

Nitrogen and its derivatives are essential for life on earth; the nitrogen cycle is one of the major biogeochemical cycles. Due to the plethora of involved compounds, organisms, and different ecosystems the nitrogen cycle is extremely complex. The increasing anthropogenic influence on the nitrogen cycle, especially due to intensifying agriculture and its negative consequences by eutrophication and over-fertilization [Prather *et al.*, 2001], results in an increasing scientific interest. Especially the production and conversion of climatic relevant trace gases such as nitrous oxide (N₂O) receives more and more interest. Since the industrial revolution the amounts of these trace gases in the atmosphere increased continuously [Prather *et al.*, 2001]. Particularly agriculture activities contribute to enhanced N₂O emissions directly by emissions from agriculture soils and animal production and indirectly from agriculture nitrogen input resulting in the stimulation of N₂O producing processes in both terrestrial and aquatic ecosystems [Mosier *et al.*, 1998].

1. Nitrous oxide (N₂O)

1.1 Climatic relevance of N₂O

N₂O is the most abundant combined nitrogen compound in the atmosphere [Jacobson *et al.*, 2000]. The first evidence of N₂O in the atmosphere was shown in 1938, but not until 1970 the importance of N₂O for the chemistry of the atmosphere was discovered by Paul Crutzen [Crutzen, 1970; McElroy and McConnell, 1971]. N₂O influences directly and indirectly the Earth's climate: In the troposphere (0 – 12 km) it acts as a greenhouse gas like carbon dioxide (CO₂), directly contributing to global warming. Moreover, N₂O can reach up to the stratosphere (12 – 80 km) due to a relatively long atmospheric lifetime of 114 years [Prather *et al.*, 2001]. In the stratosphere it acts as the major source for nitric oxide radicals (NO) which are involved in one of the main ozone (O₃) reaction cycles [Crutzen and Schmailzl, 1983; WMO, 2003]. Since the industrial revolution at the end of the 19th century, the concentration of N₂O in the atmosphere has increased rapidly by about 17 %, from 270 ppb to 318 ppb in 2002 [Sowers, 2001; Walter *et al.*, 2004]. At present an annual increase of N₂O of 0.8 ppb is observed [Prather *et al.*, 2001].

1.2 Biological production of N₂O

Within the nitrogen cycle N₂O is an intermediary or a by-product of several biological processes. Nitrification and denitrification are commonly accepted to be the main N₂O producing processes [Bonin *et al.*, 2002; Bouwman *et al.*, 1995; Kroeze *et al.*, 1999] (see Fig. 1). Many organisms like bacteria, archaea, and even some fungi are able to produce N₂O via these two processes [Cabello *et al.*, 2004; Könneke *et al.*, 2005; Zumft, 1997; Bleakley and Tiedje, 1982]. Plants were also shown to be N₂O producers [Smart and Bloom, 2001; Pihlatie *et al.*, 2005]. However, nitrifiers are also able to denitrify, methanotrophic bacteria can nitrify, and even denitrification reactions are not strictly oxygen inhibited [Wrage *et al.*, 2001; Sorokin *et al.*, 2001]. Nitrification and denitrification are therefore no longer considered as processes specific only for single species or groups that are traditionally called nitrifiers and denitrifiers, thus the classical distinction between nitrifiers and denitrifiers begins to fade [Wrage *et al.*, 2001; Zehr and Ward, 2002]. Although nitrification and denitrification are assumed to be the most important production pathways, up to now the contribution of these processes to the global N₂O budget remains unclear and is discussed controversially [Kroeze and Seitzinger, 1998; Codispoti *et al.*, 2001; Popp *et al.*, 2002; Yamagishi *et al.*, 2005].

Fig.1 shows possible production pathways of N₂O during nitrification and denitrification.




Fig. 1: Possible production pathways of N₂O during nitrification and denitrification

Although the processes have identical educts and products, they differ by the structure of used enzymes [Moreno-Vivián *et al.*, 1999; Richardson and Watmough, 1999]. Many organisms are able to use different enzymes for similar reactions or to switch between these pathways depending on the environmental conditions [Moir *et al.*, 1995; Richardson, 2000; Wrage *et al.*, 2001]. The yield of N₂O depends on the biosynthetic pathway,

the involved species, and the environmental conditions, e.g. the concentration of dissolved oxygen [Abou Seada and Ottow, 1985; Jiang and Bakken, 2000; Brettar and Rheinheimer, 1991; Goreau et al., 1980; Vollack and Zumft, 2001; Wetzel, 1983]. The next sections give a short overview of the main N₂O producing processes.

1.2.1 Nitrification

In general, nitrification is an aerobic process in which organic or inorganic nitrogen compounds are oxidized to nitrate (NO₃⁻) via hydroxylamine (NH₂OH) and nitrite (NO₂⁻) [Papen et al., 1989; Schmidt et al., 1999] (see Fig. 1). In this process N₂O is considered as a by-product, produced either by a reduction of NO₂⁻ with NH₂OH [Chalk and Smith, 1983; Ritchie and Nicholas, 1972] or incomplete oxidation of NH₂OH [Hooper and Terry, 1979]. However, the exact metabolism is still under discussion [Hynes and Knowles, 1984; Ostrom et al., 2000].

Nitrification can be conducted by both autotrophic and heterotrophic organisms. Autotrophic nitrifiers gain energy for CO₂ fixation by oxidation of ammonium (NH₄⁺). Typically two groups of microorganisms are involved: NH₄⁺-oxidizing bacteria (primary nitrifiers) and NO₂⁻-oxidizing bacteria (secondary nitrifiers) [Bock et al., 1986]. Both groups of organisms were formerly classified as nitrifying bacteria in the family Nitrobacteraceae [Buchanan, 1917], but phylogenetically they are not closely related to each other [Woese et al., 1984a; Woese et al., 1984b; Woese et al., 1985]. Several autotrophic organisms, namely NH₄⁺-oxidizers are able to combine the nitrification pathway with denitrifying reactions under aerobic conditions, the so called nitrifier denitrification. Nitrifier denitrification includes the oxidation of NH₄⁺ to NO₂⁻, followed by the reduction of NO₂⁻ via N₂O to N₂ [Wrage et al., 2001].

Besides autotrophic nitrifiers also heterotrophic nitrifiers are known. In contrast to the former group heterotrophic nitrifiers gain no energy during this process [Robertson et al., 1988; Schmidt et al., 1999; Prosser, 1989]. A possible advantage of this pathway might be the conservation of NO₃⁻ or NO₂⁻ for use during oxygen depleted conditions [Castignetti and Hollocher, 1984]. Similar to autotrophic nitrifiers they are also able to perform denitrifying reactions under aerobic conditions, with N₂O as an intermediate (see Fig. 1). Performed by heterotrophic nitrifiers this process is called aerobic denitrification [Wrage et al., 2001]. Commonly the relevance of heterotrophic nitrifiers within the nitrogen cycle was considered to be low, due to their lower nitrification rates in compari-

son to autotrophic nitrifiers [Knowles, 1985; Jetten *et al.*, 1997]. However, in recent years much attention has been paid to heterotrophic nitrifiers because many of them are found to denitrify their nitrification products simultaneously [Robertson *et al.*, 1988; Nishio *et al.*, 1998]. This might lead to a significant underestimation of their nitrification ability and the release of N₂O by heterotrophic nitrifiers. Papen *et al.* [1989] found heterotrophic nitrifiers producing much more N₂O per cell than autotrophic nitrifiers. Thus, since heterotrophic nitrifiers are more widespread and include a variety of species [Mevel and Prieur, 2000], they may contribute significantly to the global N₂O budget [Jetten *et al.*, 1997].

1.2.2 Denitrification

Denitrification is the reduction of NO₃⁻ to N₂, with N₂O as a regular intermediate product [Naqvi *et al.*, 2000]. These reactions are characteristic for various subclasses of the Proteobacteria, but also organisms of other eubacterial and archaean genera are able to denitrify [Chen *et al.*, 2002; Shapleigh, 2000; Cabello *et al.*, 2004]. Denitrifiers are facultative anaerobic [Schlegel, 1992], and the change between aerobic and anaerobic metabolism is probably controlled particularly by the O₂ concentration, although little is known about the detailed mechanisms and signals [Baumann *et al.*, 1996; John, 1977; Soerensen, 1987]. Denitrifying enzymes are known to be O₂ sensitive, e.g., the O₂ induced inhibition of a N₂O reductase is stronger than that of other reductases [Knowles, 1982]. In suboxic habitats the inhibition of N₂O reductases therefore results in an accumulation of N₂O. In anoxic habitats, N₂O is used as an electron acceptor and is thereby reduced to N₂ [Elkins *et al.*, 1978; Cohen and Gordon, 1978].

1.3 Sources and sinks of N₂O

An estimation of the quantity of nitrous oxide sources was published by the Intergovernmental Panel on Climate Change in 2001 [Prather *et al.* 2001] (see Fig. 2). N₂O is released by both anthropogenic and natural sources. Anthropogenic sources are particularly agricultural ac-




Fig. 2: Sources of atmospheric N₂O, modified from the IPCC report of 2001

tivities such as crop and stock farming, but also industrial production e.g. of nylon or picric acid. The dominating natural sources for N₂O are soils and oceans.

N₂O is relatively stable. In the atmosphere degradation of N₂O takes place by photolysis and oxidative reactions [McElroy *et al.*, 1976; Crutzen, 1974]. Sinks of dissolved N₂O in the oceans are regions with depleted O₂ concentrations, where N₂O is used as an electron acceptor instead of O₂ in microbial processes [Elkins *et al.*, 1978; Cohen and Gordon, 1978]. Oceanic sinks for atmospheric N₂O are regions of deep water formation [Bange and Andreae, 1999].

1.4 N₂O and Oceans

Oceans cover more than 70 % of the earth surface. They play an important role in biogeochemical cycles, particularly in conversion of compounds, their transport and the exchange with the atmosphere. One important function of the oceans is that they can be sinks or sources of atmospheric gases, e.g. trace gases such as CO₂ or N₂O. More than 25 % of naturally produced N₂O are emitted by the oceans, including continental shelves and estuaries [Prather *et al.*, 2001; Seitzinger *et al.*, 2000] (see Fig. 3). However, the contribution of different oceanic areas to the global N₂O budget is extremely variable. Particularly coastal regions, including estuarine and upwelling regions make a significant contribution to this budget [Bange *et al.*, 1996].




Fig. 3: Natural sources of atmospheric N₂O, modified from Seitzinger *et al.* 2000

Measurements of nitrification and denitrification rates in the ocean are sparse. To estimate the production of N_2O a linear regression between the excess of N_2O (ΔN_2O) and the apparent oxygen utilization (AOU) is commonly used [Yoshinari, 1976; Cohen and Gordon, 1978; Butler *et al.*, 1989; Yoshida *et al.*, 1989]. A positive correlation between these two parameters indicates nitrification, whereas in anoxic habitats (N_2O is used instead of O_2 as an electron acceptor) linear correlations between ΔN_2O and AOU are not existent [Elkins *et al.*, 1978; Cohen and Gordon, 1978].

2. Study areas and short compilation of N_2O data

2.1 The Atlantic Ocean

The Atlantic Ocean (Fig. 4) is the second largest world's ocean, and extends into both the Arctic and Antarctic region. The Atlantic Ocean is divided by the Mid-Atlantic Ridge into a series of eastern and western basins. Together with several sills the Mid-Atlantic Ridge has a strong impact on the circulation of the deeper water layers. The Atlantic Ocean contains five mediterranean seas; amongst these the European Mediterranean Sea and the Baltic Sea. The most important feature is the formation of Deep Water in the northern part, which drives the oceanic water circulation, the so called "conveyor belt".

Due to this constant renewal of water, driving also the input and the transport of atmospheric oxygen to the deep ocean, the Atlantic Ocean is well ventilated and permanently oxygenated. Therefore no permanent suboxic zones comparable to those found in the Arabian Sea or the eastern tropical Pacific Ocean exist [Tomczak and Godfrey, 2001]. Despite the fact that the Atlantic Ocean is by far the best investigated part of the world's ocean, data of the vertical N_2O distribution are com-




Fig. 4: Recent measurements of vertical N_2O profiles of the last 25 years.

parably sparse. In the last decades only a few vertical profiles of N_2O were published. Fig. 4 shows the locations of three cruises in the Atlantic Ocean. The cruises CITHER 1-1 and 1-2 and BLAST II measured N_2O concentrations down to the bottom, whereas data from AMT 12 and 13 provide N_2O concentrations down to approximately 300 m. A meridional transect along $30^\circ W$ and single stations in the subtropical and tropical north-western Atlantic additionally have been measured. Table 1 gives an overview of the stations and data points measured in the Atlantic Ocean prior to our investigations.

Table 1: Overview of available N_2O data in the Atlantic Ocean, data points of *Junge and Hahn* [1971] were not available

region	cruise (year)	max. depth	number of stations / data points	reference
NE – SW Atlantic	BLAST II (1994)	5389 m	40 / 582	<i>J. Butler, pers. communication</i> [2005]
NE – SW Atlantic	AMT 12 (2003)	304 m	25 / 186	<i>G. Forster, pers. communication</i> [2005]
NE – SW Atlantic	AMT 13 (2003)	305 m	22 / 134	<i>G. Forster, pers. communication</i> [2005]
W – E tropical Atlantic	CITHER 1-1 (1993)	5678 m	21 / 440	[<i>Oudot et al.</i> , 2002]
W – E tropical Atlantic	CITHER 1-2 (1993)	3994 m	26 / 540	[<i>Oudot et al.</i> , 2002]
N - S Atlantic	1969/1970	3000 m	13 / n.a.	[<i>Junge and Hahn</i> , 1971]
NE Atlantic	1977/1978	2195 m	1 / 11	[<i>Cohen and Gordon</i> , 1979]
NW subtropical Atlantic	1976	~4800 m	14 / 248	[<i>Yoshinari</i> , 1976]
NW tropical Atlantic	1976	~1500 m	2 / 30	[<i>Yoshinari</i> , 1976]

In the present work data from four cruises in the Atlantic Ocean are reported. The first cruise (May / June 2002), started in Hamburg, Germany and followed the WOCE-A2 transect to Halifax, Canada. Depth profiles of N_2O in the subpolar North Atlantic were measured at 16 stations. The subtropical North Atlantic was investigated during March / April 2004. The cruise started in Fort de France, Martinique (French Antilles) in the western part of the Atlantic and ended in Lisbon (Portugal). Samples were taken at 37 stations. The tropical North Atlantic samples were taken during the M55-SOLAS cruise in October/November 2002. This cruise started in the western tropical North Atlantic in Willemstad, Curaçao (Netherlands Antilles) and followed a cruise track along $10 - 11^\circ N$ to Douala (Cameroon). The track included a transect to the equator between 26 and $23.5^\circ W$. N_2O profiles were taken at 20 stations. The last cruise was during the European Iron Fertilisation Experiment (EIFEX) in the subpolar South Atlantic Ocean in February/March 2004. Here seven vertical N_2O profiles were measured, inside and outside of the iron fertilized patch.

2.2 The Baltic Sea

The Baltic Sea is a mediterranean sea of the Atlantic Ocean and part of the European continental shelf consisting of a series of basins (Arkona, Bornholm, and Gotland Basin; see Fig. 5). The basins show only a restricted horizontal and vertical water exchange due to shallow sills and a clear salinity stratification of water masses. A permanent halocline effectively prevents the vertical exchange and is the reason for the development of stagnant deep waters with oxygen depletion up to anoxia accompanied by accumulation of hydrogen sulphide (H_2S). Occasional inflow of oxygenated North Sea Water terminates these stagnation periods, however, during the last two decades these inflow events became less frequent and intense, thus the periods of anoxic conditions are increasing [Tomczak and Godfrey, 2001; Nausch et al., 2003; Feistel et al., 2003]. The most important recent inflow events during the last years were in 1993 and 2003 [Feistel et al., 2003]. N_2O measurements in the Baltic Sea are as sparse as in the Atlantic Ocean. Only a few vertical profiles exist, mainly from three cruises Rönner made in spring and summer 1977 and 1980 [Rönner, 1983; Rönner and Sörensson, 1985]. He investigated the distribution of N_2O at 20 stations in the Baltic Proper, the Bothnian Sea and the Bothnian Bay. Additionally, two single profiles in the Baltic Proper (Gotland Deep and north of it) are available, measured in summer 1986 and 1987 by Brettar and Rheinheimer [Brettar and Rheinheimer, 1992]. In the present thesis one cruise in the Baltic Sea is reported, after the major inflow event in January 2003. In October 2003 N_2O concentrations were measured at 26 stations in the southern and central Baltic Sea.




Fig. 5: The Baltic Sea; AB: Arkona Basin, BB: Bornholm Basin, eGB: eastern Gotland Basin, wGB: western Gotland Basin

3. Thesis outline

Since the industrial revolution the N_2O concentration in the atmosphere has increased due to human activities [*Kroeze and Seitzinger, 1998*], and because of the annual increasing rate N_2O might influence the climate of the next decades significantly [*Lashof and Ahuja, 1990*]. Oceans play an important role for the global budget of atmospheric N_2O emissions contributing more than a quarter of naturally produced atmospheric N_2O [*Seitzinger et al., 2000; Prather et al., 2001*].

Due to the insufficient number of vertical profiles of N_2O in the oceans, particularly the North Atlantic Ocean and the Baltic Sea, one aim of this study was to provide a more detailed and high resolution picture of the distribution of N_2O . These data should be used to investigate sources and sinks of N_2O , and to determine the role of the North Atlantic Ocean and the Baltic Sea as sources of atmospheric N_2O . Furthermore possible production pathways of N_2O should be investigated using the relationship between N_2O and relevant production parameters such as oxygen and nitrate, and their relation to physical processes in the oceans. During this thesis the possibility arose to start first investigations about the vertical structure of bacterial communities. 16S rRNA genes were used to obtain information of the vertical distribution of bacteria in the subtropical North Atlantic, with special regard to potential N_2O producers.

Chapter 2 gives an overview of the distribution of N_2O in the North Atlantic, whereas in chapter 3 the N_2O air-sea exchange in the tropical North Atlantic is presented. The possibility of stimulating N_2O production processes and counteracting the climatic benefits of a drawdown of atmospheric CO_2 during iron fertilization experiments in the South Atlantic is investigated in chapter 4. In chapter 5 the distribution of N_2O with regard to changed environmental conditions by a North Sea Water inflow in the Baltic Sea were studied. Since oceanic N_2O is mainly produced via microbial pathways we additionally investigated the bacterial community structures by 16S rRNA genes in the subtropical North Atlantic to obtain more information about the distribution of potential N_2O producers. This is content of chapter 6. Finally a short storage experiment is presented. Measurements are often not possible directly after sampling. For a validation of obtained data it is necessary to know, whether concentrations are affected over time between sample collections to measurements.

References

- About Seada, M.N.I., and J.C.G. Ottow, Effect of increasing oxygen concentration on total denitrification and nitrous oxide release from soil by different bacteria, *Biology and Fertility of Soils*, 1 (1), 31-38, 1985.
- Bange, H.W., S. Rapsomanikis, and M.O. Andreae, Nitrous oxide in coastal waters, *Global Biogeochemical Cycles*, 10 (1), 197-207, 1996.
- Bange, H.W., and M.O. Andreae, Nitrous oxide in the deep waters of the world's oceans, *Global Biogeochemical Cycles*, 13 (4), 1127-1135, 1999.
- Baumann, B., M. Snozzi, A.J.B. Zehnder, and J.R. van der Meer, Dynamics of denitrification activity of *Paracoccus denitrificans* in continuous culture during aerobic-anaerobic changes, *Journal of Bacteriology*, 178 (15), 4367-4374, 1996.
- Bleakley, B.H., and J.M. Tiedje, Nitrous oxide production by organisms other than nitrifiers or denitrifiers, *Applied and Environmental Microbiology*, 44 (6), 1342-1348, 1982.
- Bock, E., H.-P. Koops, and H. Harms, Cell biology of nitrifying bacteria, in: *Prosser, J.I. (Ed.), Nitrification. Special Publications of the Society for General Microbiology*, 20, 17-38, 1986.
- Bonin, P., C. Tamburini, and V. Michotey, Determination of the bacterial processes which are sources of nitrous oxide production in marine samples, *Water Research*, 36 (3), 722-732, 2002.
- Bouwman, A.F., K.W. Van der Hoek, and J.G.J. Olivier, Uncertainties in the global source distribution of nitrous oxide, *Journal of Geophysical Research*, 100 (D2), 2785-2800, 1995.
- Brettar, I., and G. Rheinheimer, Denitrification in the central Baltic: evidence for hydrogen sulfide oxidation as motor of denitrification at the oxic-anoxic interface, *Marine Ecology Progress Series* 77 (2-3), 157-169, 1991.
- Brettar, I., and G. Rheinheimer, Influence of carbon availability on denitrification in the central Baltic Sea, *Limnology and Oceanography*, 37 (6), 1146-1163, 1992.
- Buchanan, R.E., Studies on the nomenclature and classification of the bacteria III. The families of the Eubacteriales, *Journal of Bacteriology*, 2, 347-350, 1917.
- Butler, J.H., J.W. Elkins, T.M. Thompson, and K.B. Egan, Tropospheric and dissolved N₂O of the West Pacific and East Indian Oceans during the El-Nino Southern Oscillation event of 1987, *Journal of Geophysical Research-Atmospheres*, 94 (D12), 14865-14877, 1989.
- Cabello, P., M.D. Roldan, and C. Moreno-Vivian, Nitrate reduction and the nitrogen cycle in archaea, *Microbiology-Sgm*, 150, 3527-3546, 2004.

- Castignetti, D., and T.C. Hollocher, Heterotrophic nitrification among denitrifiers, *Applied and Environmental Microbiology*, 47, 620-623, 1984.
- Chalk, P.M., and C.J. Smith, Chemodenitrification, *Developments in Plant and Soil Sciences*, 9, 65-89, 1983.
- Chen, M.Y., S.S. Tsay, K.Y. Chen, Y.C. Shi, Y.T. Lin, and G.H. Lin, *Pseudoxanthomonas taiwanensis* sp nov., a novel thermophilic, N₂O producing species isolated from hot springs, *International Journal of Systematic and Evolutionary Microbiology*, 52, 2155-2161, 2002.
- Codispoti, L.A., J.A. Brandes, J.P. Christensen, A.H. Devol, S.W.A. Naqvi, H.W. Paerl, and T. Yoshinari, The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the anthropocene?, *Scientia Marina*, 65, 85-105, 2001.
- Cohen, Y., and L.I. Gordon, Nitrous oxide in the oxygen minimum of the eastern tropical North Pacific: evidence for its consumption during denitrification and possible mechanisms for its production, *Deep-Sea Research*, 25 (6), 509-524, 1978.
- Cohen, Y., and L.I. Gordon, Nitrous oxide production in the ocean, *Journal of Geophysical Research-Oceans and Atmospheres*, 84 (C1), 347-353, 1979.
- Crutzen, P.J., The influence of nitrogen oxides on the atmospheric ozone content, *Quarterly Journal of the Royal Meteorological Society*, 96, 320-325, 1970.
- Crutzen, P.J., Estimates of possible variations in total ozone due to natural causes and human activities, *Ambio*, 3, 201-210, 1974.
- Crutzen, P.J., and U. Schmailzl, Chemical budgets of the stratosphere, *Planetary and Space Science*, 31 (9), 1009-1032, 1983.
- Elkins, J.W., S.C. Wofsy, M.B. McElroy, C.E. Kolb, and W.A. Kaplan, Aquatic sources and sinks for nitrous oxide, *Nature*, 275 (5681), 602-606, 1978.
- Feistel, R., G. Nausch, W. Matthäus, and E. Hagen, Temporal and spatial evolution of the Baltic deep water renewal in spring 2003, *Oceanologia*, 45 (4), 623-642, 2003.
- Goreau, T.J., W.A. Kaplan, S.C. Wofsy, M.B. McElroy, F.W. Valois, and S.W. Watson, Production of nitrite and nitrous oxide by nitrifying bacteria at reduced concentrations of oxygen, *Applied and Environmental Microbiology* 40 (3), 526-532, 1980.
- Hooper, A.B., and K.R. Terry, Hydroxylamine oxidoreductase of *Nitrosomonas*: production of nitric oxide from hydroxylamine, *Biochimica et Biophysica Acta*, 571, 12-20, 1979.
- Hynes, R.K., and R. Knowles, Production of nitrous oxide by *Nitrosomonas europaea*: effects of acetylene, pH, and oxygen, *Canadian Journal of Microbiology*, 30, 1397-1404, 1984.

- Jacobson, M.C., edited by R.J. Charlson, H. Rodhe, and G.H. Orians, *Earth System Science - From biogeochemical cycles to global change*, Academic Press, San Diego, 2000.
- Jetten, M.S.M., S. Logemann, G. Muyzer, L.A. Robertson, S. deVries, M.C.M. van Loosdrecht, and J.G. Kuenen, Novel principles in the microbial conversion of nitrogen compounds, *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology*, 71 (1-2), 75-93, 1997.
- Jiang, Q.Q., and L.R. Bakken, Nitrous oxide production and methane oxidation by different ammonia-oxidizing bacteria, *Applied and Environmental Microbiology*, 65, 2679-2684, 2000.
- John, P., Aerobic and anaerobic bacterial respiration monitored by electrodes, *Journal of General Microbiology*, 98, 231-238, 1977.
- Junge, C., and J. Hahn, N₂O measurements in the North Atlantic, *Journal of Geophysical Research*, 76 (33), 8143-8146, 1971.
- Knowles, R., Denitrification, *Microbiological Reviews*, 46, 43-70, 1982.
- Knowles, R., Microbial transformations as sources and sinks for nitrogen oxides, in J. A. Brierley (ed.), *Plantary ecology*, Van Nostrand Reinhold Co., Inc., New York, 411-426, 1985.
- Könneke, M., A.E. Bernhard, J.R. de la Torre, C.B. Walker, J.B. Waterbury, and D.A. Stahl, Isolation of an autotrophic ammonia-oxidizing marine archaeon, *Nature*, 437 (22), 543-546, 2005.
- Kroeze, C., and S.P. Seitzinger, The impact of land use on N₂O emissions from watersheds draining into the northeastern Atlantic Ocean and European Seas, *Environmental Pollution*, 102, 149-158, 1998.
- Kroeze, C., A. Mosier, L. Bouwman, Closing the global N₂O budget: A retrospective analysis 1500- 1994, *Global Biogeochemical Cycles*, 13(1), 1-8, 1999.
- Lashof J. and D. Ahuja, Relative contributions of greenhouse gas emissions to the global warming, *Nature*, 344, 529-531, 1990.
- McElroy, M.B. and J.C. McConnell (1971), Nitrous oxide: A natural source of stratospheric NO, *Journal of Atmospheric Sciences*, 28, 1085.
- Mc Elroy, M.B., J.W. Elkins, S.C. Wofsy and Y.L. Yung, Sources and sinks for atmospheric N₂O, *Reviews of Geophysics and Space Physics*, 14, 143-150, 1976.
- Mével, G., and D. Prieur, Heterotrophic nitrification by a thermophilic *Bacillus* species as influenced by different culture conditions, *Canadian Journal of Microbiology*, 46, 465-473, 2000.

- Moir, J.W.B., D.J. Richardson, and S.J. Ferguson, The expression of redox proteins of denitrification in *Thiosphaera pantotropha* grown with oxygen, nitrate, and nitrous oxide as electron acceptors, *Archives of Microbiology*, 164 (1), 43-49, 1995.
- Moreno-Vivián, C., P. Cabello, M. Martínez-Luque, R. Blasco, and F. Castillo, Prokaryotic nitrate reduction: Molecular properties and functional distinction among bacterial nitrate reductases, *Journal of Bacteriology*, 181 (21), 6573-6584, 1999.
- Mosier, A.R., C. Kroeze, C. Nevison, O. Oenema, S. Seitzinger, and O. van Cleemput, Closing the global N₂O budget: Nitrous oxide emissions through the agricultural nitrogen cycle, *Nutrient cycling in agroecosystems*, 52, 225-248, 1998.
- Naqvi, S.W.A., D.A. Jayakumar, P.V. Narvekar, H. Naik, V. Sarma, W. D'Souza, S. Joseph, and M.D. George, Increased marine production of N₂O due to intensifying anoxia on the Indian continental shelf, *Nature*, 408 (6810), 346-349, 2000.
- Nausch, G., W. Matthäus, and R. Feistel, Hydrographic and hydrochemical conditions in the Gotland Deep area between 1992 and 2003, *Oceanologia*, 45 (4), 557-569, 2003.
- Nishio, T., T. Yoshikura, H. Mishima, Z. Inouye, and H. Itoh, Conditions for nitrification and denitrification by an immobilized heterotrophic nitrifying bacterium *Alcaligenes faecalis* OKK17, *Journal of Fermentation and Bioengineering*, 86 (4), 351-356, 1998.
- Ostrom, N.E., M.E. Russ, B. Popp, T.M. Rust, and D.M. Karl, Mechanisms of nitrous oxide production in the subtropical North Pacific based on determinations of the isotopic abundances of nitrous oxide and di-nitrogen, *Chemosphere: Global Change Science*, 2 (3-4), 281-290, 2000.
- Oudot, C., P. Jean-Baptiste, E. Fourre, C. Mormiche, M. Guevel, J.F. TERNON, and P. Le Corre, Transatlantic equatorial distribution of nitrous oxide and methane, *Deep-Sea Research Part I-Oceanographic Research Papers*, 49 (7), 1175-1193, 2002.
- Papen, H., R. Vonberg, I. Hinkel, B. Thoene, and H. Rennenberg, Heterotrophic nitrification by *Alcaligenes faecalis* – NO₂⁻, NO₃⁻, N₂O, and NO production in exponentially growing cultures, *Applied and Environmental Microbiology*, 55 (8), 2068-2072, 1989.
- Pihlatie, M., P. Ambus, J. Rinne, K. Pilegaard, and T. Vesala, Plant-mediated nitrous oxide emissions from beech (*Fagus sylvatica*) leaves, *New Phytologist*, 168, 93-98, 2005.
- Popp, B.N., M.B. Westley, S. Toyoda, T. Miwa, J.E. Dore, N. Yoshida, T.M. Rust, F.J. Sansone, M.E. Russ, N.E. Ostrom, and P.H. Ostrom, Nitrogen and oxygen isotopic constraints on the origins and sea-to-air flux of N₂O in the oligotrophic subtropical North Pacific gyre, *Global Biogeochemical Cycles*, 16 (4), doi: 10.1029/2001GB001806, 2002.
- Prather, M., D. Ehhalt, F. Dentener, R. Derwent, E. Dlugokencky, E. Holland, I. Isaksen, J. Katima, V. Kirchhoff, P. Matson, P. Midgley, and M. Wang, Atmospheric chemistry and greenhouse gases, in *Climate Change 2001: The Scientific Basis. Contribution*

- of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by J.T. Houghton, Y. Ding, D.J. Griggs, M. Noguer, P.J. Van der Linden, X. Dai, K. Maskell, and C.A. Johnson, pp. 239-287, Cambridge University Press, Cambridge, UK, 2001.
- Prosser, J.I., Autotrophic nitrification in bacteria, *Advances in Microbial Physiology*, 30, 125-181, 1989.
- Richardson, D.J., Bacterial respiration: a flexible process for a changing environment, *Microbiology*, 146, 551-571, 2000.
- Richardson, D.J., and N.J. Watmough, Inorganic nitrogen metabolism in bacteria, *Current Opinion in Chemical Biology*, 3, 207-219, 1999.
- Ritchie, G.A.F., and D.J.D. Nicholas, Identification of nitrous oxide produced by oxidative and reductive processes in *Nitrosomonas europaea*, *Biochemical Journal*, 126, 1181-1191, 1972.
- Robertson, L.A., E.W.J. van Niel, R.A.M. Torremans, and J.G. Kuenen, Simultaneous nitrification and denitrification in aerobic chemostat cultures of *Thiosphaera pantotropha*, *Applied and Environmental Microbiology*, 54 (11), 2812-2818, 1988.
- Rönner, U., Distribution, production and consumption of nitrous oxide in the Baltic Sea, *Geochimica et Cosmochimica Acta*, 47, 2179-2188, 1983.
- Rönner, U., and F. Sörensson, Denitrification rates in the low oxygen waters of the stratified Baltic Proper, *Applied and Environmental Microbiology* 50 (4), 801-806, 1985.
- Schlegel, H.G., *Allgemeine Mikrobiologie*, 7th ed. Thieme Verlag, Stuttgart, New York, 1992.
- Schmidt, I., T. Gries, and T. Willuweit, Nitrification - Fundamentals of the metabolism and problems at the use of ammonia oxidizers, *Acta Hydrochimica et Hydrobiologica*, 27 (3), 121-135, 1999.
- Seitzinger, S.P., C. Kroeze, and R.V. Styles, Global distribution of N₂O emissions from aquatic systems: Natural emissions and anthropogenic effects, *Chemosphere: Global Science Change*, 2, 267-279, 2000.
- Shapleigh, J.P., The denitrifying procaryotes, in *The Prokaryotes: An Evolving Electronic Resource for the Microbiological Community*, 3rd edition, release 3.1, Springer Verlag, New York, 2001.
<http://141.150.157.117:8080/prokPUB/index.htm>
- Smart D.R, and A.J. Bloom, Wheat leaves emit nitrous oxide during nitrate assimilation, *Proceedings of the National Academy of Sciences of the United States of America*, 98 (14), pp 7875-7878, 2001.
- Soerensen, J., Nitrate reduction in marine sediment: Pathways and interactions with iron and sulfur cycling, *Geomicrobiology Journal*, 5 (3-4), 401-422, 1987.

- Sorokin, D.Y., J.G. Kuenen, and M.S.M. Jetten, Denitrification at extremely high pH values by the alkaliphilic, obligately chemolithoautotrophic, sulfur-oxidizing bacterium *Thioalkalivibrio denitrificans* strain ALJD, *Archives of Microbiology*, 175 (2), 94-101, 2001.
- Sowers, T., N₂O record spanning the penultimate deglaciation from the Vostok ice core, *Journal of Geophysical Research-Atmospheres*, 106 (D23), 31903-31914, 2001.
- Tomczak, M., and J.S. Godfrey, *Regional Oceanography: An Introduction*, 2001.
<http://www.es.flinders.edu.au/~mattom/regoc/pdfversion.html>
- Vollack, K.U., and W.G. Zumft, Nitric oxide signaling and transcriptional control of denitrification genes in *Pseudomonas stutzeri*, *Journal of Bacteriology*, 183 (8), 2516-2526, 2001.
- Walter, S., H.W. Bange, and D.W.R. Wallace, Nitrous oxide in the surface layer of the tropical North Atlantic Ocean along a west to east transect, *Geophysical Research Letters*, 31 (23), 2004.
- Wetzel, R.G., *Limnology*, Saunders College Publishing, Philadelphia, Pa., 1983.
- WMO, Scientific assessment of ozone depletion: 2002, pp. 498, WMO (World Meteorological Organization), Geneva, 2003.
- Woese, C.R., E. Stackebrandt, W.G. Weisburg, B.J. Paster, M.T. Madigan, V.J. Fowler, C.M. Hahn, P. Blanz, R. Gupta, Nealson, and G.E. Fox, The phylogeny of purple bacteria: the alpha subdivision, *Systematic and Applied Microbiology*, 5, 315-326, 1984a.
- Woese, C.R., J.B. Weisburg, B.J. Paster, C.M. Hahn, R.S. Tanner, N.R. Krieg, H.-P. Koops, H. Harms, and E. Stackebrandt, The phylogeny of purple bacteria: the beta subdivision, *Systematic and Applied Microbiology*, 5, 327-336, 1984b.
- Woese, C.R., W.G. Weisburg, C.M. Hahn, B.J. Paster, L.B. Zablen, B.J. Lewis, T.J. Macke, W. Ludwig, and E. Stackbrandt, The phylogeny of purple bacteria: the gamma subdivision, *Systematic and Applied Microbiology*, 6, 25-33, 1985.
- Wrage, N., G.L. Velthof, M.L. van Beusichem, and O. Oenema, Role of nitrifier denitrification in the production of nitrous oxide, *Soil Biology and Biochemistry*, 33 (12-13), 1723-1732, 2001.
- Yamagishi, H., N. Yoshida, S. Toyoda, B.N. Popp, M.B. Westley, and S. Watanabe, Contributions of denitrification and mixing on the distribution of nitrous oxide in the North Pacific, *Geophysical Research Letters*, 32 (4), 2005.
- Yoshida, N., H. Morimoto, M. Hirano, I. Koike, S. Matsuo, E. Wada, T. Saino, and A. Hattori, Nitrification rates and ¹⁵N abundances of N₂O and NO₃⁻ in the western North Pacific, *Nature*, 342, 895-897, 1989.
- Yoshinari, T., Nitrous oxide in the sea, *Marine Chemistry*, 4, 189-202, 1976.

Zehr, J.P., and B.B. Ward, Nitrogen cycling in the ocean: New perspectives on processes and paradigms, *Applied and environmental microbiology*, 68 (3), 1015-1024, 2002.

Zumft, W.G., Cell biology and molecular basis of denitrification, *Microbiology and Molecular Biology Reviews*, 61 (4), 533-&, 1997.

Chapter 2

Nitrous oxide in the North Atlantic Ocean

**Sylvia Walter, Hermann W. Bange, Ulrich Breitenbach,
Douglas W.R. Wallace**

Abstract

In order to investigate the role of the North Atlantic Ocean as a source of atmospheric nitrous oxide and to decipher the major formation pathways of nitrous oxide, measurements of dissolved nitrous oxide were made during three cruises in the tropical, subtropical and subpolar North Atlantic in Oct. /Nov. 2002, Mar. /Apr. 2004, and May 2002, respectively. Nitrous oxide was close to equilibrium or slightly supersaturated in the surface layers suggesting that the North Atlantic acts as a weak source of nitrous oxide to the atmosphere. Depth profiles showed supersaturation throughout the water column with a distinct increasing trend from the subpolar to the tropical region. Lowest nitrous oxide concentrations, near equilibrium and with an average of $11.0 \pm 1.7 \text{ nmol L}^{-1}$, were found in the subpolar North Atlantic where the profiles showed no clear maxima. Highest values up to 37.3 nmol L^{-1} occurred in the tropical North Atlantic with clear maxima at approximately 400 m. A positive correlation of nitrous oxide with nitrate, as well as excess nitrous oxide with AOU, was only observed in the subtropical and tropical regions. Therefore, we conclude that the formation of nitrous oxide occurs in the tropical region rather than in the subpolar region of the North Atlantic and suggest nitrification is the dominant formation pathway in the subtropical and tropical regions.

1. Introduction

Nitrous oxide (N_2O) is an important atmospheric trace gas due to its influence on the Earth's climate. In the troposphere N_2O acts as a greenhouse gas whereas in the stratosphere it is involved in the depletion of ozone by providing NO-radicals [Prather *et al.*, 2001]. Since the beginning of the industrial revolution the global mean tropospheric N_2O mole fraction has risen rapidly from 270 ppb up to 314 ppb in 1998 [Prather *et al.*, 2001]. About 24 % of the natural sources of atmospheric N_2O are contributed by the oceans [Prather *et al.*, 2001; Seitzinger *et al.*, 2000]. Nitrous oxide is an important component of the oceanic nitrogen cycle, mainly formed by the microbial processes of nitrification and denitrification [Codispoti *et al.*, 2001; Goreau *et al.*, 1980]: Nitrification is an aerobic two-step process in which ammonium is oxidized to nitrate ($\text{NH}_4^+ \rightarrow \text{NH}_2\text{OH} \rightarrow \text{NO}_2^- \rightarrow \text{NO}_3^-$) by two different groups of bacteria. In this process nitrous oxide is assumed to be a by-product, however until now the exact pathway for N_2O production remains unclear. In suboxic habitats nitrate can be reduced by denitrification to molecular nitrogen ($\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO} \rightarrow \text{N}_2\text{O} \rightarrow \text{N}_2$), here nitrous oxide is an intermediate product. Especially at oxic/suboxic boundaries N_2O is produced by coupled nitrification and denitrification, due to the transfer of common intermediates [Yoshinari *et al.*, 1997]. Another possibility is aerobic denitrification, whereby under fully aerobic conditions organisms convert ammonia into nitrogen gas without the intermediary accumulation of nitrite [Robertson *et al.*, 1988]. All processes depend on oxygen concentrations, as well as the availability of substrates such as ammonium and nitrate. Many organisms are able to switch between different pathways depending on environmental conditions, and also the yield of N_2O during a process depends on environmental conditions [Goreau *et al.*, 1980; Poth and Focht, 1985; Richardson, 2000]. Positive correlations of N_2O with apparent oxygen utilization (AOU) or nitrate are interpreted as production of nitrous oxide by nitrification [Yoshinari, 1976; Cohen and Gordon, 1978; Yoshida *et al.*, 1989]. However, up to now the dominant production pathway for N_2O on the global scale and the contribution of different pathways still remains unclear [Codispoti *et al.*, 2001; Popp *et al.*, 2002].

Information on the vertical N_2O distribution in the North Atlantic is sparse, only a few profiles are available. The first vertical profiles for the North Atlantic were published by Junge and Hahn [1971] and Yoshinari [1976], additional data were collected by Butler *et*

al. [1995], and recently data from a transect at 7°30' N were reported by *Oudot et al.* [2002]. In this paper we present a comprehensive set of 73 vertical profiles of nitrous oxide from three trans-Atlantic cruises, covering the subpolar North Atlantic, the subtropical and the tropical North Atlantic. Based on these new data, we examine the regional differences of the N₂O distribution and its formation pathways.

2. Study area

2.1 Research cruises

Samples from the three cruises were collected over the period from May 2002 to April 2004 (see Fig. 1).




Fig. 1: Cruise tracks for 'Gauss 384-1' (subpolar North Atlantic, May 16th to June 14th 2002), 'Meteor 60-5' (subtropical North Atlantic, March 9th to April 14th 2004) and 'Meteor 55' (tropical North Atlantic, October 13th to November 16th 2002). Numbers are given for stations where N₂O profiles were measured.

The first cruise (May / June 2002), started in Hamburg, Germany with the German research vessel 'Gauss'. The cruise track followed the WOCE-A2 transect to Halifax, Canada. Depth profiles of N₂O were measured at 16 stations. The WOCE-A2 transect is located between 42 °N and 49 °N.

The subtropical North Atlantic was investigated during March / April 2004 onboard the research vessel 'Meteor'. The cruise started in Fort de France, Martinique (French Antilles) in the western part of the Atlantic and ended in Lisbon (Portugal). Samples were taken at 37 stations. Most stations were co-located with stations where samples were taken during the Transient Tracers in the Ocean Program (TTO) in 1982.

The tropical North Atlantic samples were taken during the M55-SOLAS cruise [Wallace and Bange, 2004] in October/November 2002, again with the German research vessel 'Meteor'. This cruise started in the western tropical North Atlantic in Willemstad, Curaçao (Netherlands Antilles) and followed a cruise track along 10 – 11 °N to Douala (Cameroon). The track included a transect to the equator between 26 °W and 23.5 °W. N₂O profiles were taken at 20 stations.

2.2 Hydrography

Several water masses in the North Atlantic can be identified in the T-S-diagram based on data from the three cruises (see Fig. 2). The main Atlantic water masses were identified according to commonly used classification schemes [Tomczak, 1999; Alvarez et al., 2004; Aiken et al., 2000; Joyce et al., 2001; Poole and Tomczak, 1999].




Fig. 2: T-S-diagram of the North Atlantic; 2a) T-S-diagram with data from all three cruises; 2b) T-S-diagram in Fig. 2a framed by box.; AW: Amazon Water; STMW: Subtropical Mode Water; MW: Mediterranean Water; SACW: South Atlantic Central Water; NACW: North Atlantic Central Water; ;AAIW: Antarctic Intermediate Water, AABW: Antarctic Bottom Water; INADW: lower North Atlantic Deep Water; unADW: upper North Atlantic Deep Water; LSW: Labrador Sea Water

The WOCE A2 transect (Gauss 384-1 cruise), is located at the boundary region between the subpolar gyre [*Gordon, 1986*] and the subtropical gyre [*Krauss, 1996*]. This region is highly variable, characterized by the exchange of upper-ocean water between the gyres mainly via the North Atlantic Current, and the Labrador Current. One of the most important water masses here is the Labrador Sea Water (LSW). These water masses provide the major part of the North Atlantic Intermediate Water in combination with the outflow of Mediterranean Sea Water (MW), which is detected in the eastern basin of the subtropical Atlantic Ocean near the Strait of Gibraltar [*Richardson et al., 2000*] and the Antarctic Intermediate Water (AAIW) from the south [*Lorbacher, 2000*]. Additional water masses of the southern hemisphere that penetrate into the North Atlantic are the South Atlantic Central Water (SACW) and the Antarctic Bottom Water (AABW). SACW flows northwards, and mixes with the North Atlantic Central Water (NACW) at approximately 15 °N in the western and 20 °N in the eastern basin [*Poole and Tomczak, 1999; Aiken et al., 2000*].

A typical freshwater influence was found during the Meteor 55 cruise in the western tropical North Atlantic. Water of the Amazon was detected in the surface water, identified by high temperatures and low salinity. These plumes of freshwater are transported northwards by the North Brazil Current and eastwards by the equatorial current system [*Fratantoni and Glickson, 2002*].

3. Material and methods

Water samples for N₂O analysis were collected in triplicate from various depths, taken with a 24-Niskin-bottle rosette, equipped with a CTD-sensor. The analytical method applied is a modification of the method described by [Bange *et al.*, 2001]. Bubble free samples were taken immediately following oxygen sampling in 24 mL glass vials, sealed directly with butyl rubber stoppers and crimped with aluminium caps. To prevent microbial activity, samples were poisoned with 500 µL of 2 mM mercury chloride solution. Then 10 mL of sample was replaced with a helium headspace for each vial, and the samples were allowed to equilibrate for at least two hours at room temperature (temperature was recorded continuously). A 9 mL subsample from the headspace was used to flush a 2 mL sample loop after passing through a moisture trap (filled with Sicapent[®], Merck Germany). Gaschromatographic separation was performed at 190 °C on a packed molecular sieve column (6ft x 1/8"SS, 5A, mesh 80/100, Alltech GmbH, Germany). The N₂O was detected with an electron capture detector. A mixture of argon with 5 % by volume methane was used as carrier gas with a flow of 21 mL min⁻¹. For the two-point calibration procedure we used standard gas mixtures with 311.8 ± 0.2 ppb and 346.5 ± 0.2 ppb N₂O in synthetic air (Deuste Steininger GmbH, Mühlhausen Germany). The standard mixtures have been calibrated against the NOAA (National Oceanic and Atmospheric Administration, Boulder, Co.) standard scale in the laboratories of the Air Chemistry Division of the Max Planck Institute for Chemistry, Mainz, Germany.

Calculations

N₂O water concentrations (C_{N₂O}) were calculated as follows:

$$C_{N_2O} \left[\text{nmol L}^{-1} \right] = \left(\beta \times P V_{wp} + \frac{x P}{R T} V_{hs} \right) / V_{wp} \quad (1)$$

where β stands for the Bunsen solubility in nmol L⁻¹ atm⁻¹ [Weiss and Price, 1980], x is the dry gas mole fraction of N₂O in the headspace in ppb, P is the atmospheric pressure in atm, V_{wp} and V_{hs} stand for the volumes of the water and headspace phases, respectively. R is the gas constant (8.2054 10⁻² L atm mol⁻¹ K⁻¹) and T is the temperature during equilibration. The salinity was measured by the CTD-Sensor during water sample collection. The overall relative mean analytical error was estimated to be ± 1.8 %.

The excess N_2O (ΔN_2O) was calculated as the difference between the calculated N_2O equilibrium concentration and the measured concentration of N_2O as follows

$$\Delta N_2O \text{ [nmol L}^{-1}\text{]} = N_2O \text{ (observed)} - N_2O \text{ (equilibrium)}. \quad (2)$$

To calculate the N_2O equilibrium concentration we used three different atmospheric mole fractions. Between the mixed layer and the atmosphere, N_2O exchanges in about three weeks [Najjar, 1992], thus we calculated ΔN_2O in the mixed layer using the actual atmospheric N_2O value of 318 ppb measured during the ‘Meteor 55’ cruise [Walter *et al.*, 2004]. Below the thermocline, exchange with the atmosphere is unlikely, thus, calculated N_2O equilibrium concentrations depend on the atmospheric N_2O mole fraction at the time of deep-water formation. However, the exact atmospheric mole fraction of N_2O during deep-water formation is unknown because of uncertainty in age determination of water masses. Generally, tropical Atlantic deep waters below 2000 m seem to be older than 200 years [Broecker and Peng, 2000]. Therefore for depths > 2000 m ΔN_2O was calculated with the tropospheric preindustrial value of 270 ppb [Flückiger *et al.*, 1999]. An average of the actual and the preindustrial atmospheric value (i.e., 294 ppb) was used for the depth range between the upper thermocline and 2000 m. The thermocline was defined as the depth where the temperature differs from the surface temperature by more than 0.5 °C [Tomczak and Godfrey, 2001]. For the subtropical and subpolar region we calculated ΔN_2O with these same mole fractions, although the age of water masses is different to the tropical Atlantic and therefore some values may be underestimated, whereas others may be overestimated. The resulting uncertainties of ΔN_2O are about 10-15 %; however, our conclusions are not significantly affected by this uncertainty. The equilibrium values of dissolved oxygen (O_2) were calculated with the equation given by [Weiss, 1970].

The apparent oxygen utilization (AOU) was calculated as followed:

$$AOU \text{ [}\mu\text{mol L}^{-1}\text{]} = O_2 \text{ (equilibrium)} - O_2 \text{ (observed)}. \quad (3)$$

4. Results

4.1 Distribution of nitrous oxide in the North Atlantic

4.1.1 N₂O distribution along isopycnal levels




Fig. 3: Distribution of N₂O in the North Atlantic along isopycnal levels. Dots indicate stations with available data for the isopycnal levels. 3a) surface – thermocline; 3b) thermocline – 26.0; 3c) 26.1 – 26.5; 3d) 26.6 – 27.0; 3e) 27.1 – 27.5; 3f) 27.6 – 27.93

In the surface layer of the North Atlantic (Fig. 3a) N_2O concentrations were relatively uniform with $8.5 \pm 1.2 \text{ nmol L}^{-1}$. In the region of the Labrador Current N_2O concentrations were enhanced with an average of $11.6 \pm 0.9 \text{ nmol L}^{-1}$. During the ‘Meteor 55’ cruise, a plume of Amazon Water had been identified in the western basin of the tropical North Atlantic [Körtzinger, 2003]. In contrast to [Oudot *et al.*, 2002], who reported enhanced values in the plume of the Amazon River, we found no influence on N_2O concentrations [Walter *et al.*, 2004].

Below the thermocline, N_2O concentrations were variable with respect to depths and regions. We found highest concentrations in the eastern basin of the tropical North Atlantic throughout the water column, with maximum concentrations on σ_θ surfaces between 26.3 and 27.1 (Fig. 3b-e). At the Midatlantic Ridge, located at approximately 40°W , a distinct boundary between the western and eastern Atlantic basins was observed (Fig. 3d-e). In the eastern subtropical North Atlantic, at approximately 1000 m (σ_θ 27.6-27.7), a tongue of outflow water from the Mediterranean Sea was detected by higher values of salinity and temperature [Richardson *et al.*, 2000]. However, we found no apparent influence of the Mediterranean water on N_2O concentrations.

Like the surface layer, deep waters (Fig. 3f) showed nearly uniform N_2O concentrations, though with higher values of $13.3 \pm 1.6 \text{ nmol L}^{-1}$. However, a weak but distinct trend of decreasing concentrations from the tropics ($13.1 \pm 1.3 \text{ nmol L}^{-1}$) to the subpolar North Atlantic ($11.1 \pm 1.4 \text{ nmol L}^{-1}$) could be observed.

4.1.2 Vertical N₂O distribution

The vertical distribution of N₂O showed characteristically different profiles in different regions of the North Atlantic (Fig. 4a-c), and between the western and eastern basins of these regions (Fig. 5a-c).




Fig. 4: N₂O concentration in the North Atlantic plotted against depth. 4a) subpolar, 4b) subtropical, 4c) tropical North Atlantic




Fig. 5: Selected vertical N₂O profiles in the western basin (filled symbols) and the eastern basin (open symbols) in the North Atlantic. Stations were indicated by numbers. 5a) subpolar, 5b) subtropical, 5c) tropical North Atlantic

In the subpolar North Atlantic (Fig. 4a) vertical gradients of nitrous oxide were weak over the complete cruise track, with no clear or only a very weakly pronounced subsurface maximum. N₂O concentrations were near equilibrium ($11.0 \pm 1.3 \text{ nmol L}^{-1}$) throughout the water column, with average concentrations of $8.6 \pm 1.4 \text{ nmol L}^{-1}$ in the surface layer ($\sigma_{\theta} 25.3\text{-}27.0$) and $11.3 \pm 1.5 \text{ nmol L}^{-1}$ below the thermocline down to the bottom ($\sigma_{\theta} 26.2\text{-}27.7$). No differences between the western and the eastern basin were found (Fig. 5a).

In contrast, N₂O distributions and profiles in both the subtropical and tropical North Atlantic showed strong variations with water depth (Fig. 4b-c, Fig. 5b-c). In both regions, the profiles generally had one distinct maximum. In the surface layer (σ_θ 19.3-26.8) concentrations were uniform, increasing below the thermocline up to a maximum and decreasing down to approximately 2000 m (σ_θ 22.1-27.8). Below 2000 m (σ_θ 27.8-27.9) N₂O concentrations were nearly constant with depth in both basins.

In the subtropical North Atlantic (Fig. 4b) N₂O surface concentrations were 8.7 ± 0.7 nmol L⁻¹, comparable to those in the subpolar North Atlantic. Maximum values were found at depths between 600 to 1000 m (σ_θ 26.7 – 27.7); values ranged from 14.0 in the eastern basin (#195) to 21.3 nmol L⁻¹ in the western basin (#156). Below 2000 m ($\sigma_\theta > 27.8$), concentrations were nearly constant at 13.1 ± 0.9 nmol L⁻¹. Profiles in the western subtropical North Atlantic showed distinct maxima, while in the eastern basin no clear maximum was expressed (Fig. 5b). From the western to the eastern basin maximum concentrations decreased slightly from 17.7 ± 1.4 nmol L⁻¹ to 15.1 ± 0.7 nmol L⁻¹. East of the Midatlantic Ridge maxima were not clearly expressed and were broader. Additionally, maximum Δ N₂O values were lower in the eastern (5.5 ± 0.6 nmol L⁻¹) than in the western basin (7.9 ± 1.3 nmol L⁻¹).

In the tropical North Atlantic (Fig. 4c) surface concentrations, with an average of 7.4 ± 1.1 nmol L⁻¹, were slightly lower than in the subtropical and subpolar North Atlantic. In contrast to the subtropical North Atlantic, maxima of N₂O concentrations were found at shallower depths of approximately 400m (σ_θ 26.8 – 27.1). The maximum values were higher in general, and ranged from 23.8 nmol L⁻¹ in the western basin (#4) to 32.1 nmol L⁻¹ in the eastern basin (#47). At station #36, located in the Guinea Dome area [Siedler *et al.*, 1992; Snowden and Molinari, 2003], we observed the highest N₂O values of about 37.3 nmol L⁻¹ at 400 m (σ_θ 27.0) (Fig. 4c). At the equatorial stations the N₂O maxima were found at shallower water depths (240 m to 280 m, σ_θ 26.6 – 27.0). Maximum values ranged from 22.3 nmol L⁻¹ (#26) to 24.9 nmol L⁻¹ (#24). Below 2000 m ($\sigma_\theta > 27.8$) concentrations in the tropical North Atlantic were similar to those in the subtropics with an average of 13.2 ± 1.3 nmol L⁻¹. In both basins of the tropical North Atlantic profiles looked similar with sharp and clear maxima, however, concentrations throughout the water column increased from west to east (Fig. 5c). Below 2000 m ($\sigma_\theta > 27.8$) N₂O concentrations were about 2 nmol L⁻¹ higher in the eastern than in the

western basin, whereas the difference of the maximum values was even higher (approximately 8 nmol L^{-1}).

N_2O profiles of the subtropical and tropical North Atlantic are in good agreement, both in absolute concentrations and shape of profiles, with those measured during the Bromine Latitudinal Air/Sea Transect II (BLAST II) cruise in Oct./Nov. 1994 [Butler *et al.*, 1995] <http://www.cmdl.noaa.gov/hats/ocean/blast2/blastii.html>. Below 1500 m in the North Atlantic as far as 20°S , the mean N_2O concentration observed by Butler *et al.* was about $13.5 \pm 1.0 \text{ nmol L}^{-1}$ ($n = 18$) which is in good agreement with our measurements ($12.6 \pm 1.5 \text{ nmol L}^{-1}$, $n = 449$).

4.2 Comparison of nitrous oxide with other parameters

Parameters most relevant for comparison with nitrous oxide are those assumed to be directly in connection with production pathways of N_2O , like oxygen or the apparent oxygen utilization (AOU), and nitrate. In general, we found $\Delta\text{N}_2\text{O}$ positively correlated with AOU and nitrate (Fig. 6). However, in view of differences between the basins, these correlations might not be sufficient and need higher resolution. Therefore data were divided as shown in Figure 7.




Fig. 6: $\Delta\text{N}_2\text{O}$ in comparison with AOU (6a) and nitrate (6b) for the North Atlantic, sigma σ_θ is colour coded in kg m^{-3} .

$$y (\Delta\text{N}_2\text{O}) = 0.089x (\text{AOU}) - 3.245 \quad R^2 = 0.71$$

$$y (\Delta\text{N}_2\text{O}) = 0.331x (\text{NO}_3^-) - 1.263 \quad R^2 = 0.32$$

In Fig. 7a-f correlations between the excess of N_2O (ΔN_2O) with AOU (Fig. 7a-c) and with NO_3^- (Fig. 7d-f) are divided for all data of the respective regions.




Fig. 7: ΔN_2O in comparison with AOU (7a-c) and nitrate (7d-f), sigma σ_θ is colour coded in $kg\ m^{-3}$.

7a and 7d) subpolar North Atlantic

7b and 7e) subtropical North Atlantic;

$$y (\Delta N_2O) = 0.0473x (\text{AOU}) + 1.1409, R^2 = 0.86 \text{ for } \sigma_\theta < 27.7 (\sim <1000\text{m});$$

$$y (\Delta N_2O) = 0.2497x (NO_3^-) + 1.1776, R^2 = 0.80 \text{ for } \sigma_\theta < 27.7 (\sim <1000\text{m});$$

circled data represent $\sigma_\theta > 27.7 (\sim >1000\text{m})$

7c and 7f) tropical North Atlantic;

$$\text{line a: } y (\Delta N_2O) = 0.0785x (\text{AOU}) + 2.4381, R^2 = 0.87 \text{ for } \sigma_\theta < 27.1 (\sim <500\text{m});$$

$$\text{line b: } y (\Delta N_2O) = 0.0942x (\text{AOU}) - 6.6675, R^2 = 0.81 \text{ for } \sigma_\theta > 27.1 (\sim >500\text{m});$$

$$\text{line a: } y (\Delta N_2O) = -0.4848x (NO_3^-) + 3.1756, R^2 = 0.79 \text{ for } \sigma < 27.1 (\sim <500\text{m});$$

$$\text{line b: } y (\Delta N_2O) = 0.7379x (NO_3^-) - 14.665, R^2 = 0.79 \text{ for } \sigma > 27.1 (\sim >500\text{m})$$

Since a multiple regression analysis turned out to be not applicable due to the co-linearity between the independent variables nitrate and AOU, we applied simple regression analysis for the isopycnal levels below the thermocline (see Tab. 2). Above the thermocline in the surface layer no correlations were found.

In the subpolar North Atlantic (Fig. 7a, d) ΔN_2O is low, with values near zero. There were no significant correlations found with AOU (Fig. 7a) or nitrate (Fig. 7d; see Tab. 2).

In the subtropical North Atlantic (Fig. 7b, e) $\Delta\text{N}_2\text{O}$ was also low with values ranging from 0 to 10 nmol L⁻¹. In contrast to the subpolar North Atlantic we found significant correlations between $\Delta\text{N}_2\text{O}$, AOU (Fig. 7b) and nitrate (Fig. 7e; see Tab. 2), especially at depths down to the N_2O maxima (ca. 1000 m; $\sigma_\theta < 27.7$; see regression lines in Fig. 7b and 7e). Below 1000 m ($\sigma_\theta > 27.7$) $\Delta\text{N}_2\text{O}$ did not correlate with AOU or NO_3^- (circled data in Fig. 7b and 7e).

In the tropical North Atlantic (Fig. 7c, f) correlations between $\Delta\text{N}_2\text{O}$, AOU (Fig. 7c) and nitrate (Fig. 7f) were more pronounced. We observed different $\Delta\text{N}_2\text{O}/\text{AOU}$ ratios at depths down to $\Delta\text{N}_2\text{O}$ maxima and below the $\Delta\text{N}_2\text{O}$ maxima down to the bottom. From the surface layer down to 500 m ($\sigma_\theta < 27.1$; see regression line a) the slope of the regression line ($\Delta\text{N}_2\text{O}/\text{AOU}$) was approximately 20 % lower than at depths below 500 m ($\sigma_\theta > 27.1$; see regression line b), what implies that the yield of N_2O at equal AOU is lower at shallower depths.

Table 2: Regression analyses between $\Delta\text{N}_2\text{O}$ and AOU, and $\Delta\text{N}_2\text{O}$ and NO_3^- at different isopycnal levels. Bold numbers mean relationships with significance levels of $p < 0.001$ and $R^2 > 0.5$. The coefficients a and b mean slope and intercept.

region	sigma	n	$\Delta\text{N}_2\text{O} / \text{AOU}$			$\Delta\text{N}_2\text{O} / \text{NO}_3^-$		
			a	b	R^2	a	b	R^2
subpolar	thermocline - 26.0	-	-	-	-	-	-	-
	26.1 - 26.5	-	-	-	-	-	-	-
	26.6 - 27.0	13	0.045	-0.209	0.21	0.205	-0.882	0.09
	27.1 - 27.5	49	0.028	-0.452	0.23	0.159	-1.087	0.14
	27.6 - 27.9	179	0.022	-1.714	0.07	0.164	-3.240	0.03
subtropical	thermocline - 26.0	-	-	-	-	-	-	-
	26.1 - 26.5	45	0.038	1.307	0.58	0.297	1.337	0.51
	26.6 - 27.0	106	0.065	0.397	0.75	0.390	0.186	0.68
	27.1 - 27.5	121	0.055	0.426	0.67	0.312	-0.052	0.60
	27.6 - 27.9	355	0.069	-3.083	0.38	-0.126	4.579	0.02
tropical	thermocline - 26.0	34	0.080	3.783	0.72	0.599	4.681	0.61
	26.1 - 26.5	39	0.099	-1.421	0.83	0.777	-3.212	0.81
	26.6 - 27.0	98	0.107	-3.112	0.49	0.376	6.068	0.20
	27.1 - 27.5	63	0.114	-10.638	0.43	1.289	-35.05	0.42
	27.6 - 27.9	69	0.075	-4.853	0.67	0.619	-11.83	0.66

5. Discussion

Based on our results, we were able to assign measured N_2O concentrations to the water masses as shown in Figure 2a (Fig. 8). In the following we discuss distributions and possible origins of nitrous oxide at different depths with regard to these water masses.




Fig. 8: N_2O concentration (a) and ΔN_2O (b) distributed in a T-S-diagram, the N_2O and ΔN_2O concentrations are colour coded in $nmol L^{-1}$. AW: Amazon Water; STMW: Subtropical Mode Water; MW: Mediterranean Water; SACW: South Atlantic Central Water; NACW: North Atlantic Central Water; AAIW: Antarctic Intermediate Water; LSW: Labrador Sea Water

5.1 N_2O in the surface layer of the North Atlantic

In the surface layer of the North Atlantic the distribution of N_2O was relatively uniform, with concentrations near equilibrium. This is in line with the assumption that denitrification and nitrification as sources of nitrous oxide in the surface layer seem to be negligible due to the high oxygen concentrations and light inhibition of nitrification [Horrigan *et al.*, 1981]. Thus, correlations between ΔN_2O , AOU and nitrate were nonexistent. Accordingly, we suggest that the N_2O distribution in the surface layer is most likely driven by solubility and mixing effects. This is also applicable for the enhanced N_2O concentrations found in the Labrador Current. The ΔN_2O concentrations, which are corrected for temperature, showed no enhanced values in this region. Thus, higher N_2O concentrations in the Labrador Current are likely caused by the solubility effect as well. In the warmer surface layer of the tropical North Atlantic ΔN_2O values were up to $4 nmol L^{-1}$, indicating the tropical North Atlantic acts as a weak source for atmospheric N_2O [Walter *et al.*, 2004].

5.2 N₂O below the surface layer down to 2000 m

Variations of N₂O vertical profiles reflected effects of water mass ventilation and subsurface N₂O production history. In the subpolar North Atlantic we assume that the hydrographic setting (such as convection processes during deep water formation and vertical mixing) is responsible for the observed concentrations and distributions of N₂O and Δ N₂O. The most important feature in the subpolar North Atlantic is the formation of deep water in winter in the Labrador and Irminger Seas as part of the North Atlantic circulation in 500 to 2000 m [Rhein, 2000], which carries the atmospheric N₂O imprint to depth. Labrador Sea Water spreads rapidly east- and southwards [Rhein, 2000] and thus causes the uniform distribution of N₂O within the subpolar North Atlantic. Although the productivity of phytoplankton is relatively high in the subpolar North Atlantic, we assume low biological production of N₂O because of two factors: 1) high oxygen concentrations and 2) low temperatures. The yield of N₂O depends on oxygen concentration [Goreau *et al.*, 1980; Poth and Focht, 1985; Richardson, 2000; Codispoti *et al.*, 1992], whereas high oxygen concentrations weaken the production of N₂O. Furthermore the low temperatures of the North Atlantic might have been crucial as well. The temperature dependence of both nitrification rates and enzyme activities is controversially discussed [Berounsky and Nixon, 1990; Vouve *et al.*, 2000; Barnard *et al.*, 2005; Herbert, 1999; Rheinheimer, 1964; Hansen *et al.*, 1981; Rysgaard *et al.*, 1996], however growth rates and biological production of bacteria clearly depend on the prevailing temperatures [Bock and Wagner, 2001; Hoppe *et al.*, 2002]. Thus, N₂O production might not be limited directly by temperature but indirectly by the limited abundance of N₂O producing bacteria.

In the subtropical North Atlantic concentrations of N₂O and Δ N₂O were distinctly higher compared to the subpolar North Atlantic. Profiles differed clearly between the western and eastern basin. N₂O profiles in the western basin showed clearly expressed N₂O maxima between 600 to 1000 m. This pattern was not observable east of the Midatlantic Ridge where N₂O and Δ N₂O concentrations were lower than in the western basin, and no peak maxima were observed. Hydrographic processes likely explain the shape of profiles, especially the advection of Labrador Sea Water (LSW) into the eastern basin. LSW with low N₂O concentrations is transported either along the eastern continental slope of America or across the Charlie-Gibbs-Fracture-Zone [Bower *et al.*, 2002]. It flows into the eastern subtropical basin at 500-2000 m [Rhein, 2000; Alvarez *et al.*, 2004], and

spreads north- and southwards [Bower *et al.*, 2002; Rhein, 2000]. In the western basin N₂O concentrations and profiles are in agreement with profiles published by [Yoshinari, 1976], who also found maximum values in water masses with lower oxygen concentrations. These were identified as Antarctic Intermediate Water (AAIW), which flows northwards. We assume the AAIW transports N₂O from the south to the subpolar North Atlantic. At depths shallower than 1000 m ($\sigma_0 < 27.7$) Δ N₂O was significantly correlated with oxygen utilization and nitrate concentrations (Fig. 7b, 7e; Tab. 2), indicating nitrification has contributed to measured N₂O concentrations.

In the tropical North Atlantic the N₂O profiles and the observed trend along the West-East transect are in overall agreement with recently published data from a transect along 7.5 °N [Oudot *et al.*, 2002] and a previous study in the Guinea Dome area [Oudot *et al.*, 1990]. Although the overall pattern is the same, we observed generally lower N₂O concentrations than Oudot *et al.* [2002]. This might be a result of a calibration disagreement, supported by measured atmospheric N₂O values of 316 ppb in 1993. For example, we found a peak N₂O concentration of up to 37.3 nmol L⁻¹ in South Atlantic Central Water (SACW) of the eastern basin, whereas Oudot *et al.* [2002] reported values of up to 60 nmol kg⁻¹. Oudot *et al.* [2002] assumed enhanced biological activity and remineralization of organic matter in upwelling ecosystems to be responsible for these higher values in the east. However, upwelling in this area is a temporary event [Voituriez *et al.*, 1982; Siedler *et al.*, 1992], and during our cruise no upwelling was observed.

Despite the fact that upwelling might have a long-term large-scale effect, we suppose additional reasons for the higher N₂O concentrations in the eastern basin. The productivity in the eastern basin is fueled not only by coastal upwelling [Signorini *et al.*, 1999] but also by dust deposition off the West African coast [Mills *et al.*, 2004]. Moreover, nutrient input by major tropical rivers such as the Senegal, Gambia and Niger [Perry *et al.*, 1996] contribute to enhanced production off the West African coast, indicated by enhanced chlorophyll a concentrations (for the 2002 seasonal cycle of chlorophyll a see monthly data set of Sea-viewing Wide Field-of-view Sensor (SeaWiFS): <http://earthobservatory.nasa.gov/Observatory/Datasets/chlor.seawifs.html>).

Enhanced productivity leads to a high export production [Antia *et al.*, 2001]. Subsequently lowered O₂ concentrations in the eastern intermediate layers due to the demineralization of organic matter support the production of N₂O. Therefore, enhanced N₂O

concentrations in the eastern basin at this time of year (Oct./Nov. 2002) might be a residual signal of past high production episode [Signorini *et al.*, 1999].

Upwelling events, indicated by lower sea surface temperatures, were only found at the equator. Surface N₂O concentration and sea surface temperature were positively correlated [Walter *et al.*, 2004], and the comparably shallow N₂O maxima along the equator were caused by upwelling.

Due to the occurrence of linear relationships between $\Delta\text{N}_2\text{O}$ and AOU and between N₂O and nitrate we conclude that nitrification might be the major pathway of N₂O formation in the tropical Atlantic Ocean [Yoshida *et al.*, 1989]. Like N₂O, $\Delta\text{N}_2\text{O}$ showed an increasing trend from West to East indicating that nitrification is more pronounced in the eastern than the western basin of the tropical Atlantic.

5.3 N₂O in deep waters > 2000 m ($\sigma_\theta > 27.8$)

Because the deep ocean contains high nitrate concentrations, nitrification was assumed to be responsible for N₂O production [Zehr and Ward, 2002; Bange and Andreae, 1999]. Due to the low $\Delta\text{N}_2\text{O}$ in deep waters and insufficient correlations with nitrate and AOU, we assume N₂O at these depths probably originates from deep water formation and mixing processes of southern and northern hemisphere water masses. N₂O profiles from a cruise into the Antarctic circumpolar current (2 °E / 49.5 °) [Walter *et al.*, in press] and BLAST II data east of Patagonia reveal distinctly higher N₂O concentrations in the deep waters of the southern hemisphere, with values of approximately 17 nmol L⁻¹. Northwards transport within Antarctic Bottom Water could lead to enhanced N₂O concentrations in the deep water of the North Atlantic by mixing and diffusion process.

6. Conclusions

N₂O concentrations in the North Atlantic showed characteristic variations in the vertical and horizontal distributions. In general, distribution of N₂O can be explained by a combination of biological and hydrographic reasons. The main conclusions of the present study are

- ④ Production of N₂O by nitrification occurs mainly in the tropical North Atlantic, especially in the eastern basin. Maximum values were found in the Antarctic Intermediate Water (AAIW) in the western basin, and in the South Atlantic Central Water (SACW) in the eastern basin.
- ④ Vertical N₂O distribution and shape of profiles in the subtropical North Atlantic originate from production by nitrification and advection of AAIW from the south into the western subtropical North Atlantic, respectively advection of LSW from the north in the eastern subtropical North Atlantic.
- ④ In the subpolar North Atlantic mainly mixing processes may control the distribution of N₂O, particularly the deep water formation in the Labrador Sea. Production seems to be negligible.
- ④ Tropical and subtropical regions showed supersaturation throughout the water column, thus the tropical and subtropical North Atlantic act as a source of atmospheric N₂O.
- ④ Outflow water of the Amazon or the Mediterranean Sea does not affect the N₂O concentration.

Acknowledgements

We thank the captains and crews of FS Meteor and FS Gauss for their help during sampling. We especially thank Peter Fritsche, Hans Peter Hansen, Frank Malien and Jens Schafstall for nutrient and oxygen measurements and CTD handling during the Meteor cruises. Thanks to Klaus Peter Koltermann and his colleagues of the Bundesamt für Seeschifffahrt und Hydrographie in Hamburg for the opportunity to participate on the cruise 'Gauss 384-1'. Thanks to R. Hoffmann (Max Planck Institute for Chemistry, Mainz, Germany) for calibration of our standard gas mixtures. This study was financially supported by the Deutsche Forschungsgemeinschaft (DFG) by grants no WA1434/1, WA1434/3, and WA 1434/5.

References

- Aiken, J., N. Rees, S. Hooker, P. Holligan, A. Bale, D. Robins, G. Moore, R. Harris, and D. Pilgrim, The Atlantic Meridional Transect: overview and synthesis of data, *Progress in Oceanography*, 45 (3-4), 257-312, 2000.
- Alvarez, M., F.F. Perez, H. Bryden, and A.F. Rios, Physical and biogeochemical transports structure in the North Atlantic subpolar gyre, *Journal of Geophysical Research-Oceans*, 109 (C3), 2004.
- Antia, A.N., W. Koeve, G. Fischer, T. Blanz, D. Schulz-Bull, J. Scholten, S. Neuer, K. Kremling, J. Kuss, R. Peinert, D. Hebbeln, U. Bathmann, M. Conte, U. Fehner, and B. Zeitzschel, Basin-wide particulate carbon flux in the Atlantic Ocean: Regional export patterns and potential for atmospheric CO₂ sequestration, *Global Biogeochemical Cycles*, 15 (4), 845-862, 2001.
- Bange, H.W., and M.O. Andreae, Nitrous oxide in the deep waters of the world's oceans, *Global Biogeochemical Cycles*, 13 (4), 1127-1135, 1999.
- Bange, H.W., S. Rapsomanikis, and M.O. Andreae, Nitrous oxide cycling in the Arabian Sea, *Journal of Geophysical Research-Oceans*, 106 (C1), 1053-1065, 2001.
- Barnard, R., P.W. Leadley, and B.A. Hungate, Global change, nitrification, and denitrification: A review, *Global Biogeochemical Cycles*, 19 (1), doi: 10.1029/2004GB002282, 2005.
- Berounsky, V.M., and S.W. Nixon, Temperature and the annual cycle of nitrification in waters of Narragansett Bay, *Limnology and Oceanography*, 35 (7), 1610-1617, 1990.
- Bock, E., and M. Wagner, Oxidation of inorganic nitrogen compounds as an energy source, in *The Prokaryotes: An Evolving Electronic Resource for the Microbiological Community*, 3rd edition, release 3.7, Springer Verlag, New York, 2001
<http://141.150.157.117:8080/prokPUB/index.htm>
- Bower, A.S., B. Le Cann, T. Rossby, W. Zenk, J. Gould, K. Speer, P.L. Richardson, M.D. Prater, and H.M. Zhang, Directly measured mid-depth circulation in the north-eastern North Atlantic Ocean, *Nature*, 419 (6907), 603-607, 2002.
- Broecker, W.S., and T.-H. Peng, Comparison of ³⁹Ar and ¹⁴C ages for waters in the deep ocean, *Nuclear Instruments and Methods in Physics Research B*, 172, 473-478, 2000.
- Butler, J.H., J.M. Lobert, S.A. Yvon, and L.S. Geller, The distribution and cycling of halogenated trace gases, in *Reports on Polar Research No. 168 - The expedition ANTARKTIS XII of RV "Polarstern" in 1994/95: Reports of legs ANT XII/1 and 2*, edited by G. Kattner, and D.K. Fütterer, pp. 33-40, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 1995.

- Codispoti, L.A., J.A. Brandes, J.P. Christensen, A.H. Devol, S.W.A. Naqvi, H.W. Paerl, and T. Yoshinari, The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the anthropocene?, *Scientia Marina*, 65 (Suppl. 2), 85-105, 2001.
- Codispoti, L.A., J.W. Elkins, T. Yoshinari, G.E. Friederich, C.M. Sakamoto, and T.T. Packard, *On the nitrous oxide flux from productive regions that contain low oxygen waters*, 271-284 pp., Oxford and IBH, New Delhi (India), 1992.
- Cohen, Y., and L.I. Gordon, Nitrous oxide in the oxygen minimum of the eastern tropical North Pacific: evidence for its consumption during denitrification and possible mechanisms for its production, *Deep-Sea Research*, 25 (6), 509-524, 1978.
- Flückiger, J., A. Dallenbach, T. Blunier, B. Stauffer, T.F. Stocker, D. Raynaud, and J.M. Barnola, Variations in atmospheric N₂O concentration during abrupt climatic changes, *Science*, 285 (5425), 227-230, 1999.
- Fratantoni, D.M., and D.A. Glickson, North Brazil current ring generation and evolution observed with SeaWiFS, *Journal of Physical Oceanography*, 32 (3), 1058-1074, 2002.
- Gordon, A.L., Interocean exchange of thermohaline water, *Journal of Geophysical Research*, 91, 5037-5047, 1986.
- Goreau, T.J., W.A. Kaplan, S.C. Wofsy, M.B. McElroy, F.W. Valois, and S.W. Watson, Production of NO₂⁻ and N₂O by nitrifying bacteria at reduced concentrations of oxygen, *Applied and Environmental Microbiology*, 40 (3), 526-532, 1980.
- Hansen, J.I., T.H. Blackburn, and K. Henriksen, Seasonal distribution of nitrifying bacteria and rates of nitrification in coastal marine sediments, *Microbial Ecology*, 7 (4), 297-304, 1981.
- Herbert, R.A., Nitrogen cycling in coastal marine ecosystems, *FEMS Microbiology Reviews*, 23, 563-590, 1999.
- Hoppe, H.G., K. Gocke, R. Koppe, and C. Begler, Bacterial growth and primary production along a north-south transect of the Atlantic Ocean, *Nature*, 416 (6877), 168-171, 2002.
- Horrigan, S.G., A.F. Carlucci, and P.M. Williams, Light inhibition of nitrification in sea-surface films, *Journal of Marine Research*, 39 (3), 557-565, 1981.
- Joyce, T.M., A. Hernandez-Guerra, and W.M. Smethie, Zonal circulation in the NW Atlantic and Caribbean from a meridional World Ocean Circulation Experiment hydrographic section at 66°W, *Journal of Geophysical Research-Oceans*, 106 (C10), 22095-22113, 2001.
- Junge, C., and J. Hahn, N₂O measurements in the North Atlantic, *Journal of Geophysical Research*, 76 (33), 8143-8146, 1971.

- Körtzinger, A., A significant CO₂ sink in the tropical Atlantic Ocean associated with the Amazon River plume, *Geophysical Research Letters*, 30 (24), 2287, doi: 10.1029/2003GL018841, 2003.
- Krauss, W., Comments on the development of our knowledge of the general circulation of the North Atlantic Ocean, *The Warmwatersphere of the North Atlantic Ocean*, W. Krauss (ed.), pp 1-31, Gebrüder Borntraeger, Berlin-Stuttgart, 1996.
- Lorbacher, K., Niederfrequente Variabilität meridionaler Transporte in der Divergenzzone des nordatlantischen Subtropen- und Subpolarwirbels - Der WOCE-Schnitt A2, *Bericht des Bundesamtes für Seeschifffahrt und Hydrographie*, 22, 156, 2000.
- Mills, M.M., C. Ridame, M. Davey, J. La Roche, and R.J. Geider, Iron and phosphorus co-limit nitrogen fixation in the eastern North Atlantic, *Nature*, 429, 292-294, 2004.
- Najjar, R.G., Marine biogeochemistry, in *Climate System Modeling*, edited by K.E. Trenberth, pp. 241-280, Cambridge University Press, Cambridge, 1992.
- Oudot, C., C. Andrie, and Y. Montel, Nitrous oxide production in the tropical Atlantic Ocean, *Deep-Sea Research Part I*, 37 (2), 183-202, 1990.
- Oudot, C., P. Jean-Baptiste, E. Fourre, C. Mormiche, M. Guevel, J.-F. TERNON, and P. Le Corre, Transatlantic equatorial distribution of nitrous oxide and methane, *Deep-Sea Research Part I*, 49, 1175-1193, 2002.
- Perry, G.D., P.B. Duffy, and N.L. Miller, An extended data set of river discharges for validation of general circulation models, *Journal of Geophysical Research*, 101 (D16), 21,339-21,349, 1996.
- Poole, R., and M. Tomczak, Optimum multiparameter analysis of the water mass structure in the Atlantic Ocean thermocline, *Deep-Sea Research Part I-Oceanographic Research Papers*, 46 (11), 1895-1921, 1999.
- Popp, B.N., M.B. Westley, S. Toyoda, T. Miwa, J.E. Dore, N. Yoshida, T.M. Rust, F.J. Sansone, M.E. Russ, N.E. Ostrom, and P.H. Ostrom, Nitrogen and oxygen isotopic constraints on the origins and sea-to-air flux of N₂O in the oligotrophic subtropical North Pacific gyre, *Global Biogeochemical Cycles*, 16 (4), doi: 10.1029/2001GB001806, 2002.
- Poth, M., and D.D. Focht, ¹⁵N kinetic analysis of N₂O production by *Nitrosomonas europaea* - an examination of nitrifier denitrification, *Applied and Environmental Microbiology*, 49 (5), 1134-1141, 1985.
- Prather, M., D. Ehhalt, F. Dentener, R. Derwent, E. Dlugokencky, E. Holland, I. Isaksen, J. Katima, V. Kirchhoff, P. Matson, P. Midgley, and M. Wang, Atmospheric chemistry and greenhouse gases, in *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by J.T. Houghton, Y. Ding, D.J. Griggs, M. Noguer, P.J. Van der Linden, X. Dai, K. Maskell, and C.A. Johnson, pp. 239-287, Cambridge University Press, Cambridge, UK, 2001.

- Rhein, M., Oceanography - Drifters reveal deep circulation, *Nature*, 407 (6800), 30-31, 2000.
- Rheinheimer, G., Untersuchungen über den Einfluß der Temperatur auf die Nitrifikation im Elbe-Aestuar, *Archiv für Mikrobiologie*, 49, 283-290, 1964.
- Richardson, D.J., Bacterial respiration: a flexible process for a changing environment, *Microbiology*, 146, 551-571, 2000.
- Richardson, P.L., A.S. Bower, and W. Zenk, A census of Meddies tracked by floats, *Progress in Oceanography*, 45 (2), 209-250, 2000.
- Robertson, L.A., E.W.J. van Niel, R.A.M. Torremans, and J.G. Kuenen, Simultaneous nitrification and denitrification in aerobic chemostat cultures of *Thiosphaera pantotropa*, *Applied and Environmental Microbiology*, 54 (11), 2812-2818, 1988.
- Rysgaard, S., K. Finster, and H. Dahlgard, Primary production, nutrient dynamics and mineralisation in a northeastern Greenland fjord during the summer thaw, *Polar Biology*, 16 (7), 497-506, 1996.
- Seitzinger, S.P., C. Kroeze, and R.V. Styles, Global distribution of N₂O emissions from aquatic systems: Natural emissions and anthropogenic effects, *Chemosphere: Global Science Change*, 2 (3), 267-279, 2000.
- Siedler, G., N. Zangenberg, and R. Onken, Seasonal changes in the tropical Atlantic circulation - Observation and simulation of the Guinea Dome, *Journal of Geophysical Research-Oceans*, 97 (C1), 703-715, 1992.
- Signorini, S.R., R.G. Murtugudde, C.R. McClain, J.R. Christian, J. Picaut, and A.J. Busalacchi, Biological and physical signatures in the tropical and subtropical Atlantic, *Journal of Geophysical Research-Oceans*, 104 (C8), 18367-18382, 1999.
- Snowden, D.P., and R.L. Molinari, Subtropical cells in the Atlantic Ocean: An observational summary, in *Interhemispheric Water Exchange in the Atlantic Ocean*, edited by G.J. Goni, and P. Malanotte-Rizzoli, pp. 287-312, Elsevier, New York, 2003.
- Tomczak, M., Some historical, theoretical and applied aspects of qualitative water mass analysis, *Journal of Marine Research*, 57, 275-303, 1999.
- Tomczak, M., and J.S. Godfrey, *Regional Oceanography: An Introduction*, 2001.
<http://www.es.flinders.edu.au/~mattom/regoc/pdfversion.html>
- Voituriez, B., A. Herbland, and R. Leborgne, Equatorial upwelling in the eastern Atlantic during FGGE, *Oceanologica Acta*, 5 (3), 301-314, 1982.
- Vouve, F., G. Guiraud, C. Marol, M. Girard, P. Richard, and M.J.C. Laima, NH₄⁺ turnover in intertidal sediments of Marennes-Oleron Bay (France): effect of sediment temperature, *Oceanologica Acta*, 23 (5), 575-584, 2000.

- Wallace, D.W.R., and H.W. Bange, Introduction to special section: Results of the *Meteor 55*: Tropical SOLAS Expedition, *Geophysical Research Letters*, 31 (L23S01), 1-4, 2004.
- Walter, S., H.W. Bange, and D.W.R. Wallace, Nitrous oxide in the surface layer of the tropical North Atlantic Ocean along a west to east transect, *Geophysical Research Letters*, 31 (23), 2004.
- Walter, S., I. Peeken, K. Lochte, and H.W. Bange, Nitrous oxide measurements during EIFEX, the European Iron Fertilisation Experiment in the subpolar South Atlantic Ocean, *Geophysical Research Letters* (in press.), 2005.
- Weiss, R.F., The solubility of nitrogen, oxygen and argon in water and seawater, *Deep-Sea Research*, 17, 721-735, 1970.
- Weiss, R.F., and B.A. Price, Nitrous oxide solubility in water and seawater, *Marine Chemistry*, 8, 347-359, 1980.
- Yoshida, N., H. Morimoto, M. Hirano, I. Koike, S. Matsuo, E. Wada, T. Saino, and A. Hattori, Nitrification rates and ^{15}N abundances of N_2O and NO_3^- in the western North Pacific, *Nature*, 342, 895-897, 1989.
- Yoshinari, T., Nitrous oxide in the sea, *Marine Chemistry*, 4, 189-202, 1976.
- Yoshinari, T., M.A. Altabet, S.W.A. Naqvi, L. Codispoti, A. Jayakumar, M. Kuhland, and A. Devol, Nitrogen and oxygen isotopic composition of N_2O from suboxic waters of the eastern tropical North Pacific and the Arabian Sea - Measurement by continuous-flow isotope-ratio monitoring, *Marine Chemistry*, 56 (3-4), 253-264, 1997.
- Zehr, J.P., and B.B. Ward, Nitrogen cycling in the ocean: New perspectives on processes and paradigms, *Applied and Environmental Microbiology*, 68 (3), 1015-1024, 2002.

Chapter 3

Nitrous oxide in the surface layer of the tropical North Atlantic Ocean along a West to East transect

Sylvia Walter, Hermann W. Bange, Douglas W. R. Wallace

Abstract

Nitrous oxide (N₂O) was measured during the first German SOLAS (Surface Ocean – Lower Atmosphere Study) cruise in the tropical North Atlantic Ocean onboard R/V *Meteor* during October/November 2002. About 900 atmospheric and dissolved N₂O measurements were performed with a semi-continuous GC-ECD system equipped with a seawater-gas equilibrator. Surface waters along the main transect at 10°N showed no distinct longitudinal gradient. Instead, N₂O saturations were highly variable ranging from 97% to 118% (in the Guinea Dome Area, 11°N, 24°W). When approaching the continental shelf of West Africa, N₂O surface saturations went up to 113%. N₂O saturations in the region of the equatorial upwelling (at 0–1.5 °N, 23.5–26 ° W) were correlated with decreasing sea surface temperatures and showed saturations up to 109%. The overall mean N₂O saturation was 104 ± 4 % indicating that the tropical North Atlantic Ocean is a net source of atmospheric N₂O.

Introduction

Nitrous oxide (N_2O) is an important atmospheric trace gas because it influences, directly and indirectly, the Earth's climate to a significant degree: In the troposphere, it acts as a greenhouse gas with a relatively long atmospheric lifetime [IPCC, 2001] whereas in the stratosphere it is the major source for nitric oxide radicals, which are involved in one of the main ozone reaction cycles [WMO, 2003]. Published source estimates indicate that the world's oceans play a major role in the global budget of atmospheric nitrous oxide [IPCC, 2001]. Generally, oligotrophic areas seem to be near equilibrium with the atmosphere, whereas coastal and equatorial upwelling areas show enhanced N_2O concentrations [Nevison *et al.*, 1995; Suntharalingam and Sarmiento, 2000]. Here we present about 900 measurements of dissolved and atmospheric N_2O during the first German SO-LAS (Surface Ocean – Lower Atmosphere Study) cruise. It is the first high-resolution data set of N_2O in the tropical North Atlantic Ocean along a West to East transect and it is complementary to previous N_2O measurements of Oudot *et al.* [1990; 2002] and Weiss *et al.* [1992].

The cruise took place onboard R/V *Meteor* (expedition no. M55) from Willemstad (Curaçao, Netherl. Antilles) to Douala (Cameroon) from 12 October to 17 November 2002. The cruise track consisted of two main transects: (i) The West to East transect along 10-12°N covering the oligotrophic tropical North Atlantic Ocean and the continental shelf area of the West African coast off Guinea Bissau and (ii) a shorter West to East transect along the equatorial upwelling (Fig. 1).




Fig. 1: Cruise track of M55 in October–November 2002. N_2O measurements were started 17 October and were finished 14 November. Areas of special interest discussed in the text are marked.

1. Method

N_2O was determined with a gas chromatograph equipped with an electron capture detector. Further details of the analysis system are described in *Bange et al. [1996]*. A series of measurements of atmospheric N_2O and N_2O in seawater-equilibrated air followed by two standards was repeated every 50 min. Mixtures of N_2O in synthetic air were used to obtain two-point calibration curves. The mixtures used contained 311.7 ± 0.1 and 346.5 ± 0.2 ppb N_2O , respectively. These are gravimetrically prepared gas mixtures (Deuste Steinger GmbH, Mühlhausen, Germany) and have been calibrated against the NOAA (National Oceanic and Atmospheric Administration, Boulder, Co.) standard scale in the laboratories of the Air Chemistry Division of Max Planck Institute for Chemistry Mainz, Germany. The precision, calculated as the ratio of the standard deviation of the atmospheric measurements and the mean atmospheric mixing ratio, was 0.8%.

Seawater was pumped continuously from a depth of 4 m into a shower-type equilibrator developed by R. F. Weiss (Scripps Institution of Oceanography, La Jolla, Ca.). N_2O con-

centrations (C , in nmol L^{-1}) were calculated by applying the solubility equation of *Weiss and Price* [1980]:

$$C = \beta(T, S) x' P,$$

where x' is the measured N_2O dry mole fraction, P is the atmospheric pressure, and β is the solubility coefficient, which is a function of the water temperature (T) and salinity (S). Time series of surface seawater temperature (SST), salinity, wind speed, and atmospheric pressure were obtained from the ship's records. Differences between the seawater temperature at the seawater intake and the continuously recorded water temperature in the equilibrator were corrected:

$$C_w = C \beta(T_{eq}) / \beta(SST)$$

with $\beta(SST)$ and $\beta(T_{eq})$ representing the N_2O solubility at seawater temperature and water temperature inside the equilibrator at the time of the measurement, respectively. N_2O saturations (Sat) in % (i.e., 100% = equilibrium) were calculated as follows:

$$Sat = 100 C_w / C_a$$

where C_a is the equilibrium concentration of dissolved N_2O based on the actual measurement of ambient air (see above). The mean relative errors of the N_2O concentrations and saturations were calculated to be 1.2 % and 1.6 %, respectively (details of the error propagation computation are given in *Bange et al.* [2001]).

2. Results and Discussion

The mean atmospheric N₂O dry mole fraction was 318 ± 3 ppb. Due to the seasonal northward shift of the Intertropical Convergence Zone to about 10°N, the origin of the air masses sampled during the cruise were from both the northern and the southern hemisphere. 4-days air mass back trajectories (provided by the German Weather Service, Offenbach, Germany) indicated that air masses sampled at latitudes south of 7°N originated from the southern hemisphere. Based on this classification we computed mean N₂O values for northern and southern hemisphere air masses of 319 ± 3 ppb and 317 ± 2 ppb, respectively. The observed atmospheric values are in agreement with N₂O measurements at the baseline monitoring stations Ragged Point, Barbados and Cape Grim, Tasmania. Monthly mean values were 317 ppb (Cape Grim) and 318 ppb (Ragged Point) for October/November 2002. These values were taken from the Advanced Global Atmospheric Gases Experiment (AGAGE) data set (updated version from November 2003) [Prinn *et al.*, 2000]. AGAGE data are available from the anonymous ftp site: cdiac.esd.ornl.edu (subdirectory /pub/ale_gage_Agage/Agage/gc-md/monthly) at the Carbon Dioxide Information Analysis Center in Oak Ridge, Tennessee.

N₂O saturations along the main cruise track ranged from 97% to 118% and the SST was generally between 27 and 30 °C (Fig. 2). Since the main cruise track was located between the eastward flowing North Equatorial Countercurrent (NECC) and the westward flowing North Equatorial Current (NEC) [Stramma and Schott, 1999], we crossed several times meandering waters of different origins causing a high variability of the N₂O saturation: Low N₂O saturations of about 100% observed around 24 Oct., 27-28 Oct., and 2 Nov. were generally associated with decreases in salinity (Fig. 2).




Fig. 2: Salinity, seasurface temperature (SST), N₂O saturation, wind speed in 10 m height (u10), and N₂O flux density during M55. Areas of special interest discussed in the text are marked (see Fig. 1): A, equatorial upwelling; B, Guinea Dome; C, shelf of West Africa (water depths <200m).

This results from the retroflection of the North Brazil Current, which advects Amazon plume waters (with low N₂O, see below) eastward into the NECC [Fratantoni and Glickson, 2002]. Freshwater influences were observed twice: First, at around 50°E (19 Oct., Fig. 2) when we crossed the northern boundary of the Amazon River plume (minimum salinity 32.14) and second, on the continental shelf off West Africa where we measured a drop in salinity down to 31.30 (5-6 Nov., Fig. 2). N₂O saturations were not enhanced in the Amazon River plume, whereas an increase in N₂O saturations up to 113% was observed on the West African shelf. The low N₂O saturations in the Amazon River plume were attributed to the fact that N₂O-rich waters from the Amazon River are N₂O-depleted because of outgasing to the atmosphere and mixing with near-equilibrium oceanic waters while distributed to the North [Oudot et al., 2002]. The high N₂O saturations on the continental African shelf might result from N₂O-rich riverine waters or groundwater seepage, but not from coastal upwelling as indicated by the uniform SSTs. N₂O saturations up to 118% were observed in the area of the Guinea Dome at 11°N, 24°W (3-4 Nov., Fig. 2) which is well-known for pronounced Ekman upwelling [Siedler et al., 1992; Signorini et al., 1999]. In the equatorial region (0-1.5°N, 28-30 Oct., Fig. 2) SSTs dropped well below 27°C and were associated with enhanced N₂O saturations (up to 109%). We found a good correlation between N₂O concentrations and SST in the equatorial upwelling region (Fig. 3) indicating that the enhanced surface N₂O saturations were resulting from upwelling of N₂O-enriched subsurface waters.




Fig. 3: Correlation ($r^2 = 0.94$, $n = 46$) of SST and N_2O concentration in the equatorial upwelling area ($0-2^\circ N$, $23.5-26^\circ W$). Open circles stand for in-situ measurements and crosses stand for the corresponding equilibrium concentrations calculated as a function of the actual SST, salinity, atmospheric dry mole fraction, and ambient pressure.

In order to account for the NECC/NEC system and the observed N_2O features we defined two latitudinal aligned open ocean regions and a shelf region: 1) the tropical North Atlantic ranging from $1.5-12^\circ N$ with $SST > 27^\circ C$, 2) the equatorial upwelling from $0-1.5^\circ N$ with $SST < 27^\circ C$, and 3) the shelf area off the West African coast (water depth $< 200m$). An overview of the regional mean concentrations and saturations is given in Table 1.

Table 1: Mean N_2O concentrations, saturations, and flux densities during M55. Values are given as mean \pm 1sd. Number of measurements is given in parenthesis.

	Overall mean ($n = 451$)	$0 - 1.5^\circ N$ ($n = 27$)	$1.5 - 12^\circ N$ ($n = 416$)	Shelf ($n = 8$)
Concentration, $nmol L^{-1}$	6.00 ± 0.24	6.49 ± 0.07	5.27 ± 0.20	6.31 ± 0.11
Saturation, %	104 ± 4	107 ± 1	103 ± 3	110 ± 2
Flux density, $nmol m^{-2} s^{-1}$	0.007 ± 0.011	0.018 ± 0.006	0.006 ± 0.011	0.002 ± 0.002

The mean N₂O concentrations and saturations of the shelf and equatorial regions are significantly enhanced compared to the 1.5-12°N region. The high variability calculated for the tropical North Atlantic region is biased by the complex hydrography, which is influenced by the Amazon plume, the NECC/NEC system and the Guinea Dome upwelling with highly variable N₂O concentrations. However, a more detailed regional analysis is hampered by the limited data set.

Our data from the tropical North Atlantic are in agreement with previously published data. Recently, *Oudot et al.* [2002] reported a mean N₂O saturation of 108 ± 3 %, mainly measured along two transects at 7.5°N and 4.5°S during January–March 1993. They also observed a trend towards enhanced values when approaching the West African coast (up to 118%). In a previous study in the Guinea Dome area during June–August 1986, *Oudot et al.* [1990] observed mean N₂O saturations in the range from 126 ± 5 to 132 ± 6 % which are considerably higher than our results (Table 1). In the period from 1979 to 1989, *Weiss et al.* [1992] took part in several measurement campaigns with cruise tracks across the tropical North Atlantic Ocean. Their N₂O measurements are in good agreement with the results presented here. For example, the mean N₂O saturation during the first part of the TTO/TAS leg 3 in February 1983, which covered two transects along 9.5°N (from 20.25 to 28 °W) and 28°W (from 9.5 to the equator), was about 105 %. Enhanced values during TTO/TAS leg 3 were observed on the coast off Guinea-Bissau (up to 179%) and in the equatorial upwelling (up to 111%). In contrast to our measurements, the high N₂O values observed off Guinea Bissau were caused by coastal upwelling (SST <27°) [*Weiss et al.*, 1992]. Summarizing the results from various N₂O measurements in the open tropical North Atlantic, we found only slight differences (with the exception of the data from the Guinea Dome area by *Oudot et al.* [1990]). Significant differences as found for the Guinea Dome might be caused by seasonal variability of the circulation patterns [*Stramma and Schott*, 1999] in connection with different spatial data coverage. Since coastal upwelling was absent during our cruise, N₂O saturations on the shelf off West Africa were comparably low.

2.1 N₂O air-sea exchange

The air–sea exchange flux density (F) was parameterized as

$$F = k_w(u) (C_w - C_a),$$

where k_w (in m s^{-1}) is the gas transfer coefficient as a function of wind speed (u in 10 m height), C_w is the measured N₂O seawater concentration, and C_a is the equilibrium N₂O concentration in seawater based on the measured atmospheric value (for calculation of C_w and C_a see Methods section). To calculate k_w , we used the combined linear and quadratic k_w – u relationship (N00) from *Nightingale et al. [2000]*:

$$k_w = 9.25 \cdot 10^{-5} u + 6.17 \cdot 10^{-7} u^2.$$

The N00 relationship shows a dependence on wind speeds intermediate between the commonly used relationships of *Liss and Merlivat [1986]* and *Wanninkhof [1992]*. The measured wind speeds were normalized to 10 m height by using the relationship of *Garratt [1977]*. k_w was adjusted by multiplying with $(Sc/600)^{-0.5}$, where Sc is the *Schmidt* number for N₂O. Sc was calculated using empirical equations for the kinematic viscosity of seawater [*Siedler and Peters, 1986*] and the diffusion coefficient of N₂O in water. N₂O diffusion coefficients (D_{N_2O} in $\text{m}^2 \text{s}^{-1}$) were calculated with the equation derived from a compilation of actual measurements [*Rhee, 2000*]:

$$D_{N_2O} = 3.16 \times 10^{-6} \exp(-18370 / RT),$$

where T is the water temperature in K and R is the universal gas constant. The commonly used equation for D_{N_2O} by *Broecker and Peng [1974]* was replaced since *Rhee's [2000]* equation provides a more reasonable fit with a considerably reduced uncertainty of less than 10% [*Rhee, 2000*]. Flux densities calculated with the above equation are lower by about 10% when compared to computations with *Broecker and Peng's [1974]* equation [*Bange et al., 2001*]. We did not apply a correction of D_{N_2O} for seawater since the effect of seawater on the diffusion of dissolved gases is not uniform [*King et al., 1995*] and, to our knowledge, no measurements of the N₂O diffusion in seawater have been published. The regional mean flux densities clearly reflect the interplay of saturation and wind speeds (Fig. 2, Tab. 1). In the equatorial region enhanced N₂O saturations and compara-

bly high wind speeds result in high flux densities, whereas over the shelf enhanced N₂O saturations were associated with very low wind speed resulting in low flux densities (Fig. 2). The mean flux density of the tropical North Atlantic region is biased by the high variability of both N₂O saturations and wind speeds. The overall mean N₂O flux density was $0.007 \pm 0.011 \text{ nmol m}^{-2} \text{ s}^{-1}$ which is at the lower end of previously published flux densities: *Oudot et al.* [1990; 2002] computed overall mean flux densities of 0.013–0.021 $\text{nmol m}^{-2} \text{ s}^{-1}$ and $0.026 \pm 0.032 \text{ nmol m}^{-2} \text{ s}^{-1}$ for the tropical North and South Atlantic and the Guinea Dome area, respectively. The obvious discrepancy might be caused by different spatial data coverage, seasonal variability of the N₂O concentrations and wind speeds, and the use of different approaches for the transfer coefficient k_w .

3. Summary

N₂O saturations in the tropical North Atlantic Ocean during October–November 2002 were highly variable and range from 97 to 118 %. The mean overall saturation was 104 ± 4 %. Enhanced saturations were observed in the Guinea Dome area (up to 118%), in the equatorial upwelling (up to 109%), and the shallow continental shelf area off the West African Coast (up to 113%). Our results are in agreement with previously published data sets. We found a good correlation of seawater temperature with N₂O concentrations in the equatorial upwelling area. We conclude that the tropical North Atlantic Ocean is a net source of N₂O to the atmosphere with a pronounced regional variability.

Acknowledgements

We thank A. Körtzinger and M.O. Andreae for technical support and acknowledge the help of the other M55 participants and the officers and crew of R/V *Meteor*. We especially thank R. Hoffmann for the calibration of our standards and two anonymous reviewers for their valuable comments. We are indebted to the authorities of Guinea-Bissau for permission to work in their territorial waters. The investigations were financially supported by the *Deutsche Forschungsgemeinschaft* through grants WA1434/1 and WA1434/3.

References

- Bange, H.W., M.O. Andreae, S. Lal, C.S. Law, S.W.A. Naqvi, P.K. Patra, T. Rixen, and R.C. Upstill-Goddard, Nitrous oxide emissions from the Arabian Sea: A synthesis, *Atmospheric Chemistry and Physics*, *1*, 61–71, 2001.
- Bange, H.W., S. Rapsomanikis, and M.O. Andreae, The Aegean Sea as a source of atmospheric nitrous oxide and methane, *Marine Chemistry*, *53*, 41–49, 1996.
- Broecker, W.S., and T.-H. Peng, Gas exchange rates between air and sea, *Tellus*, *26*, 21–35, 1974.
- Fratantoni, D.M., and D.A. Glickson, North Brazil Current ring generation and evolution observed with SeaWiFS, *Journal of Physical Oceanography*, *32*, 1058–1074, 2002.
- Garratt, J.R., Review of the drag coefficients over oceans and continents, *Monthly Weather Review*, *105*, 915–929, 1977.
- IPCC, *Climate change 2001: The scientific basis. Contribution of working group I to the third assessment report of the Intergovernmental Panel on Climate Change (IPCC)*, 881 pp., Cambridge University Press, Cambridge (UK) and New York (NY, USA), 2001.
- King, D.B., W.J. De Bryun, M. Zheng, and E.S. Saltzman, Uncertainties in the molecular diffusion coefficient of gases in water for the estimation of air-sea exchange, in *Air-water gas transfer*, edited by B. Jähne, and E.C. Monahan, pp. 13–22, AEON Verlag & Studio, Hanau, Germany, 1995.
- Liss, P.S., and L. Merlivat, Air-sea exchange rates: introduction and synthesis, in *The role of air-sea exchange in geochemical cycling*, edited by P. Buat-Ménard, pp. 113–127, D. Reidel Publishing Company, Dordrecht, 1986.
- Nevison, C.D., R.F. Weiss, and D.J. Erickson III, Global oceanic emissions of nitrous oxide, *Journal of Geophysical Research*, *100*, 15,809–15,820, 1995.
- Nightingale, P., G. Malin, C.S. Law, A.J. Watson, P.S. Liss, M.I. Liddicoat, J. Boutin, and R.C. Upstill-Goddard, In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochemical Cycles*, *14*, 373–387, 2000.
- Oudot, C., C. Andrieu, and Y. Montel, Nitrous oxide production in the tropical Atlantic Ocean, *Deep-Sea Research*, *37*, 183–202, 1990.
- Oudot, C., P. Jean-Baptiste, E. Fourré, C. Mormiche, M. Guevel, J.-F. TERNON, and P. Le Corre, Transatlantic equatorial distribution of nitrous oxide and methane, *Deep-Sea Research I*, *49*, 1175–1193, 2002.

- Prinn, R.G., R.F. Weiss, P.J. Fraser, P.G. Simmonds, D.M. Cunnold, F.N. Alyea, S.O. O'Doherty, P. Salameh, B.R. Miller, J. Huang, R.H.J. Wang, D.E. Hartley, C. Harth, L.P. Steele, G. Sturrock, P.M. Midgley, and A. McCulloch, A history of chemically and radiatively important gases in air deduced from ALE/GAGE/AGAGE, *Journal of Geophysical Research*, 105, 17,751–17,792, 2000.
- Rhee, T.S., The process of air-water gas exchange and its application, PhD thesis, Texas A&M University, College Station, 2000.
- Siedler, G., and H. Peters, Properties of sea water, in *Oceanography*, Landolt-Börnstein New Ser., Group V, vol. 3a, edited by J. Sündermann, pp. 233–264, Springer Verlag, New York, 1986.
- Siedler, G., N. Zangenberg, and R. Onken, Seasonal changes in the tropical Atlantic circulation: Observation and simulation of the Guinea Dome, *Journal of Geophysical Research*, 97, 703–715, 1992.
- Signorini, S.R., R.G. Murtugudde, C.R. McClain, J.R. Christian, J. Picaut, and A.J. Busalacchi, Biological and physical signatures in the tropical and equatorial Atlantic, *Journal of Geophysical Research*, 104, 18,367–18,382, 1999.
- Stramma, L., and F. Schott, The mean flow field of the tropical Atlantic Ocean, *Deep-Sea Research II*, 46, 279–303, 1999.
- Suntharalingam, P., and J.L. Sarmiento, Factors governing the oceanic nitrous oxide distribution: Simulations with an ocean general circulation model, *Global Biogeochemical Cycles*, 14, 429–454, 2000.
- Wanninkhof, R., Relationship between wind speed and gas exchange over the ocean, *Journal of Geophysical Research*, 97, 7373–7382, 1992.
- Weiss, R.F., and B.A. Price, Nitrous oxide solubility in water and seawater, *Marine Chemistry*, 8, 347–359, 1980.
- Weiss, R. F., F. A. Van Woy, and P. K. Salameh, Surface water and atmospheric carbon dioxide and nitrous oxide observations by shipboard automated gas chromatography: Results from expeditions between 1977 and 1990, *Rep. NDP-044*, Carbon Dioxide Info. Anal. Cent., Oak Ridge Natl. Lab., Oak Ridge, TN, 1992.
- WMO, *Scientific assessment of ozone depletion: 2002*, 498 pp., WMO (World Meteorological Organization), Geneva, 2003.

Chapter 4

Nitrous oxide measurements during EIFEX, the European Iron Fertilisation Experiment in the subpolar South Atlantic Ocean

Sylvia Walter, Ilka Peeken, Karin Lochte, Hermann W. Bange

Abstract

We measured the vertical water column distribution of nitrous oxide (N_2O) during the European Iron Fertilization Experiment (EIFEX) in the subpolar South Atlantic Ocean during February/March 2004 (R/V *Polarstern* cruise ANT XXI/3). Despite a huge build-up and sedimentation of a phytoplankton bloom, a comparison of the N_2O concentrations within the fertilized patch with concentrations measured outside the fertilized patch revealed no N_2O accumulation within 33 days. This is in contrast to a previous study in the Southern Ocean, where enhanced N_2O accumulation occurred in the pycnocline. Thus, we conclude that Fe fertilization does not necessarily trigger additional N_2O formation and we caution that a predicted radiative offset due to a Fe-induced additional release of oceanic N_2O might be overestimated. Rapid sedimentation events during EIFEX might have hindered the build-up of N_2O and suggest, that not only the production of phytoplankton biomass but also its pathway in the water column needs to be considered if N_2O radiative offset is modeled.

1. Introduction

Inspired by the iron (Fe) limitation hypothesis [*Martin et al.*, 1991], several Fe fertilization experiments have been performed in high nutrient-low chlorophyll (HNLC) regions such as the Southern Ocean, and the subarctic and equatorial Pacific Ocean [see e.g., *Boyd*, 2004; *Boyd*, 2002]. *Fuhrman and Capone* [1991] pointed out that stimulating ocean productivity by Fe addition might result in an enhanced formation of nitrous oxide (N₂O). This point is especially important in view of the fact that N₂O is an atmospheric trace gas with a high global warming potential [*Jain et al.*, 2001]. Thus, enhanced N₂O formation by Fe addition might counteract the climatic benefits of a drawdown of atmospheric carbon dioxide (CO₂).

Fuhrman and Capone [1991] argued that enhanced productivity will lead to an enhanced nitrogen export from the euphotic zone, which in turn would result in additional N₂O formation via enhanced nitrification (NH₄⁺ → NH₂OH → NO₂⁻ → NO₃⁻). N₂O formed via nitrification is thought to be dominating in the oxic part of the world's oceans [see e.g., *Nevison et al.*, 2003]. The idea of a link between Fe fertilization and enhanced N₂O formation was supported by the study of *Law and Ling* [2001] who found a small but significant N₂O accumulation in the pycnocline during the Southern Ocean Iron Enrichment Experiment (SOIREE) in the Australasian sector of the Southern Ocean (61°S, 140°E) in February 1999. Recently, *Jin and Gruber* [2003] predicted the long-term effect of Fe fertilization on oceanic N₂O emissions on a global scale with a coupled physical-biogeochemical model. Based on their model results they concluded that Fe fertilization-induced N₂O emissions could offset the radiative benefits of the CO₂ drawdown [*Jin and Gruber*, 2003].

Here we present our measurements of N₂O during the European Iron Fertilization Experiment (EIFEX; R/V *Polarstern* cruise ANT XXI/3) in the subpolar South Atlantic Ocean from 9 February to 21 March 2004 [*Smetacek and cruise participants*, 2005].

1.1 The EIFEX setting

A mesoscale cyclonic eddy, embedded in a meander of the Antarctic Polar Front, was identified as suitable for the EIFEX study [*Strass et al.*, 2005]. The eddy was centered at 49.4°S 2.25°E and extended over an area of 60x100 km. First fertilization was performed on 12-13 February by releasing 6000 kg iron sulfate (FeSO₄) into the mixed layer over

an area of 150 km². Since iron concentrations had been decreasing [P. Croot, personal communication], fertilization was repeated on 26-27 February by releasing 7000 kg FeSO₄ over an area of 400 km². All sampled stations were located inside the eddy; the stations within fertilized waters will be called in-stations and those from unfertilized waters out-stations (Table 1). Inside and outside the fertilized patch was determined by photosynthetic activity (Fv/Fm) performed by Fast-Repetition-Rate-Fluorescence (Fast-Tracka, Chelsea, UK) [Röttgers *et al.*, 2005]. Fv/Fm is known to be a very sensitive parameter, which increases immediately after iron fertilization.

The hydrographic settings of the sampling stations were not uniform: The in-stations' hydrographic properties did not show any variability. However, the out-station 514 showed, in comparison with the in-stations, enhanced potential water temperatures in the density (σ_t) range from 27.25 to 27.7 kg m⁻³ (corresponding to a approximate depth range from 200 to 400 m). The hydrographic properties of the out-stations 546 and 587 were almost identical to the in-stations. This implies that station 514 is not a representative out-station and was therefore excluded from the comparison (see also discussion of N₂O data below).

2. Methods

Triplicate water samples from various depths were taken with from a 24 x 12 L-bottle rosette, equipped with a CTD-sensor. The analytical method applied is a modification of the method described by Bange *et al.* [2001]: Bubble free samples were taken immediately following oxygen (O₂) sampling in 24 mL glass vials, sealed directly with butyl rubber stoppers and crimped with aluminium caps. To prevent microbial activity, samples were poisoned with 500 μ L of a saturated aqueous mercury chloride (HgCl₂) solution. The samples were stored in the dark at 4 °C until analysis in our home laboratory from June to August 2004. In a time series experiment we found that N₂O concentrations in samples treated as described above did not change significantly over 10 months [Walter, 2005].

N₂O water concentrations (C_w) were calculated as follows:

$$C_w \left[\text{nmol L}^{-1} \right] = \left(\beta x' P V_{wp} + \frac{x' P}{R T} V_{hs} \right) / V_{wp}$$

where β stands for the Bunsen solubility in $\text{nmol L}^{-1} \text{ atm}^{-1}$ [Weiss and Price, 1980], x' is the dry gas mole fraction of N₂O in the headspace in ppb, P is the atmospheric pressure in atm (set to 1 atm), V_{wp} and V_{hs} stand for the volumes of the water (14 mL) and headspace phases (10 mL), respectively. R is the gas constant ($8.2054 \cdot 10^{-2} \text{ L atm mol}^{-1} \text{ K}^{-1}$) and T is the temperature during equilibration.

For calibration we used standard gas mixtures with 311.8 ± 0.2 ppb and 346.5 ± 0.2 ppb N₂O in synthetic air (DEUSTE Steininger GmbH, Mühlhausen, Germany). The standard mixtures have been calibrated against the NOAA (National Oceanic and Atmospheric Administration, Boulder, Co.) standard scale in the laboratories of the Air Chemistry Division of the Max Planck Institute for Chemistry, Mainz, Germany). The standard deviation of the N₂O concentration (C_w) was approximated with $(C_{w\text{max}} - C_{w\text{min}}) / 1.91$, where $C_{w\text{min}}$ and $C_{w\text{max}}$ stand for the minimal and maximal N₂O concentrations of the triplicate samples, respectively. The factor 1.91 is derived from the statistical method by David [1951]. The overall mean analytical error was $\pm 2.7\%$ ($\pm 0.5 \text{ nmol L}^{-1}$).

N₂O saturations (*sat*) in % (i.e., 100% = equilibrium) were calculated as $sat = 100 C_w / C_a$, where C_a is the equilibrium concentration of dissolved N₂O based on the N₂O atmospheric dry mole fraction, water temperature, and salinity [Weiss and Price, 1980]. For calculating C_a in the mixed layer an ambient air mole fraction of 317.8 ppb was applied, which is the average of the monthly mean N₂O dry mole fractions measured at the AGAGE (Advanced Global Atmospheric Gases Experiment, see Prinn et al. [2000]) baseline monitoring station Cape Grim (Tasmania) during February and March 2004. AGAGE data are available from the anonymous ftp site: [cdiac.esd.ornl.edu \(subdirectory/pub/ale_gage_agage/agage/gc-md/monthly\)](http://cdiac.esd.ornl.edu/subdirectory/pub/ale_gage_agage/agage/gc-md/monthly) at the Carbon Dioxide Information Analysis Center in Oak Ridge, Tennessee.

Dissolved O₂, nitrate, and CTD data were provided by the participating working groups. Further details can be found in the cruise report by Smetacek and cruise participants [2005].

3. Results and Discussion

An overview of the N₂O measurements during EIFEX is given in Table 1 and in Figure 1. Mixed layer N₂O saturations were comparable to surface saturations (~ 103%) from the same region measured during the Ajax cruise leg 2 in Jan-Feb 1984 [Weiss *et al.*, 1992]. Moreover, the overall mean N₂O deep water (> 2000m) concentration of $17.5 \pm 0.2 \text{ nmol L}^{-1}$ is in good agreement with the N₂O deep water-water age relationship by Bange and Andreae [1999]. Both, the observed surface saturation and deep-water concentration support the view that the N₂O samples were not affected by the time lag between sampling and measurements.

In agreement with the results from SOIREE [Law and Ling, 2001], we did not observe a difference in N₂O mixed layer saturations between in-stations and out-stations (Table 1), which implies that N₂O emissions were not significantly different either.

Table 1: N₂O measurements during EIFEX. Class. stands for classification and indicates whether a profile was inside or outside of the fertilized patch. ML stands for mixed layer; here defined as the depth where the temperature differs from the surface temperature by more than 0.5 °C. Conc. and sat. stand for concentration and saturation, respectively.

Station no.	Latitude [°S]	Longitude [°E]	Date	Days after 1 st / 2 nd fertilization	Patch class.	N ₂ O ML conc. ^a [nmol L ⁻¹]	N ₂ O ML sat. ^a [%]
513	49.59	2.05	28 Feb 04	16 / 2	In	13.3 ± 0.1 (5)	102 ± 1 (5)
514	49.31	2.34	29 Feb 04	17 / 3	Out	13.5 ± 0.3 (3)	104 ± 2 (3)
544	49.36	1.87	07 Mar 04	24 / 10	In	13.8 ± 0.5 (3)	106 ± 4 (3)
546	49.47	2.09	10 Mar 04	27 / 13	Out	13.1 (2)	102 (2)
570	49.43	2.05	14 Mar 04	31 / 17	In	13.1 ± 0.3 (5)	102 ± 3 (5)
580	49.12	2.38	16 Mar 04	33 / 19	In	12.5 ± 0.2 (3)	97 ± 1 (3)
586	49.50	2.10	18 Mar 04	35 / 21	Out	13.1 ± 0.5 (4)	102 ± 4 (4)

^a Given as average \pm standard deviation. Number of depths used for averaging is given in parenthesis.

The N₂O profiles showed a pronounced maximum between 500 and 750 m which was associated with the O₂ minimum and the nitrate maximum (Figure 1) indicating that nitrification was the main N₂O formation process. Our N₂O concentrations are comparable to N₂O measurements from the South Atlantic and Southern Oceans [Butler *et al.*, 1995; Law and Ling, 2001; Rees *et al.*, 1997].




Fig. 1: N_2O (open circles), water temperature (solid lines), NO_3^- (dashed lines), and O_2 (dashed dotted line) at the EIFEX stations listed in Table 1. O_2 data are only available for station 570 in the depth range from 0-1500m. Please note that O_2 is given in $\mu\text{mol L}^{-1}$ divided by 10.

Following the approach by *Law and Ling [2001]*, we fitted a polynomial to the $\text{N}_2\text{O}-\sigma_t$ data of stations 546 and 587 (Figure 2). Out-station 514 was excluded because it obviously was not representative as indicated by the data in Figure 2 (see also section EIFEX setting). A comparison of the N_2O concentrations of the in-stations with the polynomial fit based on the out stations revealed no significant differences (Figure 2).




Fig. 2: N_2O concentrations vs. density (σ_t) during EIFEX. (A) Out-stations: Triangles stand for stations 546 and 587 and crosses stand for station 514. The bold solid line represents a 3rd order polynomial fit based on stations 546 and 587 (see text for statistical details). The thin solid line represents a 3rd order polynomial fit based on station 514. (B) In-stations: 513, 544, 570, and 580 (symbols) compared with the polynomial fit based on out-stations 546 and 587 (bold line, see panel A). The dashed lines indicate the standard error of the predicted N_2O . Depths intervals are indicated.

A 3rd order polynomial fit to the in-stations ($-52.766x^3 + 4320.7x^2 - 117,915x + 1,072,529$, $r^2 = 0.95$, $n = 67$, standard error of predicted $N_2O = \pm 0.63 \text{ nmol L}^{-1}$) was almost identical to the out-stations' fit ($-48.474x^3 + 3967.8x^2 - 108,241x + 984,148$, $r^2 = 0.96$, $n = 30$, standard error of predicted $N_2O = \pm 0.56 \text{ nmol L}^{-1}$). Thus, we conclude that no significant changes in the N_2O concentrations occurred during EIFEX.

Our conclusion is in contrast to the observation by *Law and Ling [2001]*. They found an accumulation of N_2O up to $0.9 - 1 \text{ nmol L}^{-1}$ in the pycnocline (60 – 80 m water depth) within 13 days during SOIREE. Adapting a N_2O accumulation rate of $0.08 \text{ nmol L}^{-1} \text{ d}^{-1}$ ($= 1 \text{ nmol L}^{-1} / 13 \text{ days}$), an increase of 2.6 nmol L^{-1} ($= 0.08 \text{ nmol L}^{-1} \times 33 \text{ days}$) would have been expected for a N_2O accumulation in the pycnocline in 100 – 200 m during EIFEX. This was not the case (Figure 2). It is possible that N_2O accumulation in the pycnocline was not detected because of insufficient analytical precision and/or coarse sampling of the depths profiles: A possible N_2O accumulation must have been low ($< 0.5 \text{ nmol L}^{-1}$ over the duration of the experiment as implied by our mean analytical error) or must have taken place in a narrow depth range of less than 40 m (i.e., the mean depth spacing of sampling from the surface to the pycnocline in about 200 m). Moreover, in contrast to EIFEX, Fe addition during SOIREE was performed four times within a week over a much smaller area (50 km^2 , *Law and Ling [2001]*). Therefore, the observed N_2O accumulation in the pycnocline during SOIREE may have been a fast short-term response to the intensive short-term Fe fertilization. Because we started N_2O sampling 16 days after the first Fe addition (i.e., 2 days after the second Fe addition) we might have missed this short-term signal during EIFEX.

During EIFEX chlorophyll *a* (chl *a*) standing stocks increased 3 fold until day 26, but remarkably decreased thereafter [*Peeken et al., 2005*]. The main beneficiaries of the iron fertilization were diatoms in all size classes [*Hoffmann et al., 2005*]. Towards the end of the experiment, the diatom marker fucoxanthin and chl *a* could be followed down the water column to 4000 m and a low ratio of phaeopigments to chl *a* indicated the export of fresh material most likely originating from the iron fertilized patch [*Peeken et al., 2005*]. An explanation for the absence of an increase of N_2O in the deep (e.g., in the O_2 minimum zone) might be the very rapid export of the fresh phytoplankton material to the deep ocean during EIFEX [*Peeken et al., 2005*], which started about 23 days after the second Fe addition. Thus, we can argue that the rapid export of organic material during EIFEX might have been too rapid for the nitrifying bacteria in the deep ocean to adapt to

and, thus, an additional build-up of N₂O in the deep could not take place. Nitrifying bacteria, especially ammonium-oxidizing bacteria (AOB), are known for lag phases up to several weeks after periods of low metabolic activities [Schmidt *et al.*, 1999].

The responsible process for the N₂O accumulation during SOIREE [Law and Ling, 2001] and the proposed further increase of N₂O in prolonged iron fertilization experiments could not be identified. Thus, a possible link between N₂O accumulation and Fe fertilization remains to be not a simple cause-and-effect mechanisms and the magnitude of a possible radiative offset still needs to be proven.

4. Conclusions

We did not observe a N₂O accumulation during the *in situ* iron fertilization experiment EIFEX in the subpolar South Atlantic Ocean in February/March 2004. This is in contrast to previous measurements by Law and Ling [2001] in the Australasian sector of the Southern Ocean. We conclude that Fe fertilization does not necessarily trigger additional N₂O formation, which might depend on differences of the environmental conditions (e.g., the fate of the Fe-induced phytoplankton bloom). We caution, therefore, that predictions of a radiative offset caused by a Fe-induced additional release of oceanic N₂O [Jin and Gruber, 2003; Law and Ling, 2001] might be overestimated. In order to solve this problem further long-term experiments with particular emphasis on sedimentation processes are necessary to prove a link between Fe addition and enhancement of N₂O formation and the subsequent release of N₂O to the atmosphere.

Acknowledgements

We thank Katrin Bluhm, Linn Hoffmann, and Dieter Wolf-Gladrow for their help during sampling. We are grateful to B. Cisewski, H. Leach, H. Prandke, and V. Strass (“the CTD group”) for making the CTD data available for us. We acknowledge the help of the chief scientist Victor Smetacek, the ANT XXI/3 participants, and the officers and crew of R/V *Polarstern* for conducting a successful experiment. We especially thank R. Hoffmann for the calibration of our standards. Two anonymous reviewers provided valuable comments. The investigations were financially supported by the IFM-GEOMAR *Forschungsbereich Marine Biogeochemie*.

References

- Bange, H.W., and M.O. Andreae, Nitrous oxide in the deep waters of the world's oceans, *Global Biogeochemical Cycles*, *13*, 1127 – 1135, 1999.
- Bange, H.W., S. Rapsomanikis, and M.O. Andreae, Nitrous oxide cycling in the Arabian Sea, *Journal of Geophysical Research*, *106*, 1053 – 1065, 2001.
- Boyd, P, Ironing out algal issues in the Southern Ocean, *Science*, *304*, 396 – 397, 2004.
- Boyd, P.W., The role of iron in the biogeochemistry of the Southern Ocean and equatorial Pacific: A comparison of in situ iron enrichments, *Deep-Sea Research II*, *49*, 1803 – 1821, 2002.
- Butler, J.H., J.M. Lobert, S.A. Yvon, and L.S. Geller, The distribution and cycling of halogenated trace gases, *Berichte zur Polarforschung*, *168*, 27 – 40, 1995.
- David, H.A., Further applications of range to analysis of variance, *Biometrika*, *38*, 393 – 409, 1951.
- Fuhrman, J.A., and D.G. Capone, Possible biogeochemical consequences of ocean fertilization, *Limnology and Oceanography*, *36* (8), 1951 – 1959, 1991.
- Hoffmann, L., I. Peeken, K. Lochte, P. Assmy, and M. Veldhuis, Different reactions of Southern Ocean phytoplankton size classes to iron fertilisation, *Limnology and Oceanography*, *submitted*, 2005.
- Jain, A.K., B.P. Briegleb, K. Minschwaner, and D.J. Wuebbles, Radiative forcing and global warming potentials of 39 greenhouse gases, *Journal of Geophysical Research*, *105*, 20,773 – 20,790, 2000.
- Jin, X., and N. Gruber, Offsetting the radiative benefit of ocean iron fertilization by enhancing N₂O emissions, *Geophysical Research Letters*, *30*, 2249, doi:10.1029/2003GL018458, 2003.
- Law, C.S., and R.D. Ling, Nitrous oxide flux and response to increased iron availability in the Antarctic Circumpolar Current, *Deep-Sea Research II*, *48*, 2509 – 2527, 2001.
- Martin, J.H., R.M. Gordon, and S.E. Fitzwater, The case for iron, *Limnology and Oceanography*, *36*, 1793 – 1802, 1991.
- Nevison, C., J.H. Butler, and J.W. Elkins, Global distribution of N₂O and Δ N₂O-AOU yield in the subsurface ocean, *Global Biogeochemical Cycles*, *17*, 1119, doi:10.1029/2003GB002068, 2003.
- Peeken, I., L. Hoffmann, P. Assmy, U. Bathmann, B. Cisewski, H. Leach, K. Lochte, O. Sachs, E. Sauter, and V. Strass, Export of fresh algal material during the Southern Ocean iron fertilisation experiment, EIFEX, *Abstract presented at the ASLO Summer Meeting 2005, Santiago de Compostela, Spain, 19-24 June 2005*.

- Prinn, R.G., R.F. Weiss, P.J. Fraser, P.G. Simmonds, D.M. Cunnold, F.N. Alyea, S.O. O'Doherty, P. Salameh, B.R. Miller, J. Huang, R.H.J. Wang, D.E. Hartley, C. Harth, L.P. Steele, G. Sturrock, P.M. Midgley, and A. McCulloch, A history of chemically and radiatively important gases in air deduced from ALE/GAGE/AGAGE, *Journal of Geophysical Research*, 105, 17,751 – 17,792, 2000.
- Rees, A.P., N.J.P. Owens, and R.C. Upstill-Goddard, Nitrous oxide in the Bellingshausen Sea and Drake Passage, *Journal of Geophysical Research*, 102, 3383 – 3391, 1997.
- Röttgers, R., F. Colijn, and M. Dibbern, Algal physiology and biooptics, *Berichte zur Polar- und Meeresforschung*, 500, 82 – 88, 2005.
- Schmidt, I., T. Gries, and T. Willuweit, Nitrification – Fundamentals of the metabolism and problems at the use of ammonia oxidizers (in German with English abstract), *Acta Hydrochimica et Hydrobiologica*, 27, 121 – 135, 1999.
- Smetacek, V., and cruise participants, The expedition ANT XXI/3 of R/V Polarstern, *Berichte zur Polar- und Meeresforschung*, 500, 1 – 134, 2005.
- Strass, V., B. Cisewski, S. Gonzalez, H. Leach, K.-D. Loquay, H. Prandke, H. Rohr, and M. Thomas, The physical setting of the European Iron Fertilisation Experiment 'EIFEX' in the Southern Ocean, *Berichte zur Polar- und Meeresforschung*, 500, 15 – 46, 2005.
- Walter, S., Nitrous oxide in the Atlantic Ocean and the Baltic Sea, Ph.D. thesis, University of Kiel, Kiel, 2005.
- Weiss, R.F., and B.A. Price, Nitrous oxide solubility in water and seawater, *Marine Chemistry*, 8, 347 – 359, 1980.
- Weiss, R.F., F.A. Van Woy, and P.K. Salameh, Surface water and atmospheric carbon dioxide and nitrous oxide observations by shipboard automated gas chromatography: Results from expeditions between 1977 and 1990, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA, 1992.

Chapter 5

Nitrous oxide water column distribution during the transition from anoxic to oxic conditions in the Baltic Sea

**Sylvia Walter, Ulrich Breitenbach, Hermann W. Bange,
Günther Nausch, Douglas W.R. Wallace**

Abstract

In January 2003, a major inflow of cold and oxygen-rich North Sea Water in the Baltic Sea terminated an ongoing stagnation period in parts of the central Baltic Sea. In order to investigate the role of North Sea Water inflow to the Baltic Sea with regard to the production of nitrous oxide (N_2O), we measured dissolved and atmospheric nitrous oxide at 26 stations in the southern and central Baltic Sea in October 2003.

At the time of our cruise, water renewal had proceeded to the eastern Gotland Basin, whereas the western Gotland Basin was still unaffected by the inflow. The deep water renewal was detectable in the distributions of temperature, salinity, and oxygen concentrations as well as in the distribution of the N_2O concentrations: Shallow stations in the Kiel Bight and Pomeranian Bight were well-ventilated with uniform N_2O concentrations near equilibrium throughout the water column. In contrast, stations in the deep basins, such as the Bornholm and the Gotland Deep, showed a clear stratification with deep water affected by North Sea Water. Inflowing North Sea Water led to changed environmental conditions, especially enhanced O_2 or declining H_2S concentrations, thus, affecting the conditions for the production of N_2O . Pattern of N_2O profiles and correlations with parameters like oxygen and nitrate differed between the basins. The dominant production pathway seems to be nitrification rather than denitrification.

No indications for advection of N_2O by North Sea Water were found. A rough budget revealed a significant surplus of in situ produced N_2O after the inflow. However, due to the permanent halocline, it can be assumed that the formed N_2O does not reach the atmosphere. Hydrographic aspects therefore are decisive factors determining the final release of produced N_2O to the atmosphere.

1. Introduction

1.1 Nitrous oxide

Nitrous oxide (N_2O) is an important atmospheric trace gas which influences, directly and indirectly, the Earth's climate: In the troposphere, it acts as a greenhouse gas with a relatively long atmospheric lifetime of 114 years [Prather *et al.*, 2001]. In the stratosphere it is the major source for nitric oxide radicals, which are involved in one of the main ozone reaction cycles [WMO, 2003].

N_2O is mainly formed during microbial processes such as nitrification and denitrification. Nitrification is an aerobic two-step process in which ammonium is oxidized to nitrate. In this process, in which typically two groups of bacteria are involved, N_2O is assumed to be a by-product, the exact metabolism however is still under discussion [Ostrom *et al.*, 2000]. In suboxic habitats, nitrate can be reduced by denitrification to molecular nitrogen, with N_2O as an intermediate [Cohen and Gordon, 1978]. N_2O may also be produced by coupled nitrification and denitrification at oxic/suboxic boundaries, due to the transfer of intermediates such as nitrate and nitrite [Yoshinari *et al.*, 1997]. Other possibilities are the production of N_2O during nitrifier-denitrification or aerobic denitrification [Wrage *et al.*, 2001]. Both processes enable nitrifiers to oxidize NH_4^+ to NO_2^- , followed by the reduction of NO_2^- to N_2O or N_2 [Robertson and Kuenen, 1984; Robertson *et al.*, 1988; Richardson, 2000]. In anoxic habitats N_2O is used, instead of oxygen, as an electron acceptor [Elkins *et al.*, 1978; Cohen and Gordon, 1978].

The yield of N_2O during these processes strongly depends on the concentration of dissolved oxygen and nitrate [Brettar and Rheinheimer, 1991; Goreau *et al.*, 1980; Vollack and Zumft, 2001; Wetzel, 1983], with maximal N_2O accumulation at the interface between oxic and suboxic layers and depletion in anoxic layers [Codispoti *et al.*, 2005]. Positive correlations between N_2O and oxygen or nitrate are commonly interpreted as an indication of N_2O production by nitrification [Yoshinari, 1976; Yoshida *et al.*, 1989; Cohen and Gordon, 1978]. In contrast, production by denitrification is inferred by missing correlations [Elkins *et al.*, 1978; Cohen and Gordon, 1978]. However, up to now the dominant production pathway for N_2O on the global scale remains unclear and is discussed controversially [Codispoti *et al.*, 2001; Popp *et al.*, 2002; Yamagishi *et al.*, 2005]. Oceans emit more than 25 % of natural produced N_2O and contribute significantly to the global N_2O budget [Prather *et al.*, 2001; Seitzinger *et al.* 2000]. Particularly coastal re-

gions, including estuarine and upwelling regions play a major role for the formation and release of N_2O to the atmosphere [Bange *et al.*, 1996; Naqvi *et al.*, 2000; Seitzinger *et al.*, 2000]. In the Baltic Sea, first investigations were made by Rönner [1983] who found the Baltic Sea to be a source of atmospheric N_2O . In contrast to open ocean areas coastal regions are expected to be more influenced by conversion processes in sediments or by riverine inputs. In the Bodden waters and Danish fjords of the Baltic Sea enhanced N_2O concentrations were correlated with seasonal riverine input [Jørgensen and Sørensen, 1985; Dahlke *et al.*, 2000]. Additionally, denitrification processes in sediments were shown to contribute to the release of N_2O in Danish fjords [Jørgensen and Sørensen, 1985].

1.2 Study area

Samples of dissolved N_2O were measured at 26 stations in the western, southern and central Baltic Sea. The cruise took place on board the German research vessel Gauss (expedition no. 11/03/04) from 13th October to 25th October 2003 as part of the Cooperative Monitoring in the Baltic Sea Environment (COMBINE) program of the Baltic Marine Environment Protection Commission (Helsinki Commission, HELCOM, see <http://www.helcom.fi>). The locations of sampled stations are shown in Fig. 1.

The Baltic Sea is a mediterranean sea of the Atlantic Ocean and part of the European continental shelf. It consists of a series of basins (Arkona, Bornholm, and Gotland Basin; see Fig. 1), with restricted horizontal and vertical water exchange due to shallow sills and a clear salinity stratification of water masses.




Fig. 1: Map of the western, southern and central Baltic Sea with locations of the stations. The stations were grouped as follows: well-mixed stations are number 10, 12, 22, 30, 41, 46, 121, 130, 133, 360, OB Boje and OB 4; the Arkona Basin is represented by station 109 and 113; the Bornholm Basin is represented by station 140, 200, 213 and 222; in the eastern Gotland Basin station 250, 259, 260, 271 and 286 were grouped; and the western Gotland Basin is represented by station 240, 245 and 284. The arrow indicates the main flow direction of North Sea Water.

In January 2003 a major inflow of cold, highly saline and oxygen-rich North Sea Water was observed. It was the most important inflow event since 1993 and terminated the ongoing stagnation period in southern parts of the central Baltic Sea [Feistel *et al.*, 2003; Nausch *et al.*, 2003]. This inflow event was preceded by a minor inflow of warmer and less oxygenated water in August 2002. Due to the inflow of North Sea Water oxygen conditions changed from anoxic to oxic in most parts of the Baltic Sea. From the inflow in January 2003 until our cruise in October 2003 water renewal was already detectable at the Farö Deep (# 286), however the western Gotland Basin was still unventilated [Feistel *et al.*, 2003; Nausch *et al.*, 2003].

Due to the fact that N_2O production highly depends on environmental conditions such as e.g. oxygen concentration [e.g., Naqvi *et al.*, 2000] any natural or anthropogenic-induced shifts of coastal ecosystems will modulate the formation and subsequent release of N_2O to the atmosphere. In this context the inflow of North Sea Water into the Baltic Sea offered a good opportunity to investigate naturally changing environmental conditions with regard to the production of N_2O .

1.3 Definition of water masses

We refer to four different water masses, characterized by temperature, salinity and oxygen concentrations (Fig. 3). The definition of water masses follows the description of the 'Institut für Ostseeforschung' (IOW) cruise reports [Nausch, 2003a; Nagel, 2003; Feistel, 2003; Nausch, 2003b; Nausch, 2003c; Wasmund, 2003] and the hydrographic-chemical report of the Baltic Sea in 2003 [Nausch et al., 2004].




Fig. 2: Characterization of different water masses in the Baltic Sea, for example at station 271 in the Eastern Gotland Basin (triangles: temperature [°C], circles: salinity, squares: oxygen [$\mu\text{mol } 10^1 \text{ L}^{-1}$])

The Surface Water layer (sw) was characterized by uniform temperature and salinity, in combination with high oxygen concentrations. Below this layer, rapidly decreasing temperatures indicated Winter Water (ww), which is formed annually during convection in winter. Salinity and oxygen concentrations were still uniform. The 'old' Bottom Water (bw) was visible by increasing temperature and simultaneously increasing salinity. In this water mass, located below the Winter Water, oxygen concentrations decreased rapidly, to anoxic conditions at some stations. A permanent halocline between Winter Water and Bottom Water strongly restricts the vertical exchange and is the reason for the development of stagnant deep waters with oxygen depletion up to anoxia accompanied by accumulation of hydrogen sulphide (H_2S). Bottom Water, affected by the North Sea Water inflow in January 2003 (abw) was characterized by decreasing temperature and enhanced oxygen concentrations compared to previous Bottom Water (bw) values. Due to its higher density the affected Bottom Water lifts up the 'old' Bottom Water.

2. Methods

Water samples for N₂O analysis were collected in triplicate from various depths, taken with a 24-Niskin-bottle rosette, equipped with a CTD-probe. The analytical method applied was a modification of the method described by [Bange *et al.*, 2001]. Bubble free samples were taken immediately following oxygen sampling from the rosette in 24 mL glass vials, sealed directly with butyl rubber stoppers and crimped with aluminium caps. To prevent microbial activity, samples were poisoned with 500 µL of a 2 mM mercury chloride solution. 10 mL of the sample were then replaced with a helium headspace for each vial, and the samples were equilibrated for at least two hours at room temperature (temperature was recorded continuously). A 9 mL subsample from the headspace was used to flush a 2 mL sample loop after passing through a moisture trap (filled with Si-capent[®], Merck Germany). Gaschromatographic separation was performed at 190 °C on a packed molecular sieve column (6 ft x 1/8" SS, 5 A, mesh 80/100, Alltech GmbH, Germany). The N₂O was detected with an electron capture detector. A mixture of argon with 5 percent by volume methane was used as carrier gas with a flow of 21 mL min⁻¹. For the two-point calibration procedure we used standard gas mixtures with 311.8 ± 0.2 ppb and 346.5 ± 0.2 ppb N₂O in synthetic air (Deuste Steininger GmbH, Mühlhausen Germany). The standard mixtures have been calibrated against the NOAA (National Oceanic and Atmospheric Administration, Boulder, Co.) standard scale in the laboratories of the Air Chemistry Division of the Max Planck Institute for Chemistry, Mainz, Germany.

2.1 Calculations

N₂O water concentrations (C_{N₂O}) were calculated as follows:

$$C_{\text{N}_2\text{O}} \left[\text{nmol L}^{-1} \right] = \left(\beta \times P V_{\text{wp}} + \frac{x P}{R T} V_{\text{hs}} \right) / V_{\text{wp}} \quad (1)$$

where β stands for the Bunsen solubility in nmol L⁻¹ atm⁻¹ [Weiss and Price, 1980], x is the dry gas mole fraction of N₂O in the headspace in ppb, P is the atmospheric pressure in atm, V_{wp} and V_{hs} stand for the volumes of the water and headspace phases, respectively. R is the gas constant (8.2054 10⁻² L atm mol⁻¹ K⁻¹) and T is the temperature during equilibration. The salinity was measured by the CTD-Sensor during water sample collec-

tion; the temperature was measured while subsampling the headspace of the sample vial (i.e. the equilibration temperature). The overall relative mean analytical error was estimated to be $\pm 1.8\%$.

The excess N_2O ($\Delta\text{N}_2\text{O}$) was calculated as the difference between the calculated N_2O equilibrium concentration and the measured concentration of N_2O as follows

$$\Delta\text{N}_2\text{O} [\text{nmol L}^{-1}] = \text{N}_2\text{O} (\text{observed}) - \text{N}_2\text{O} (\text{equilibrium}). \quad (2)$$

Since the water masses in the Baltic Sea are comparably young (e.g. 11 years for the oldest bottom water at the Landsort Deep) [Meier, 2005] it is reasonable to calculate the equilibrium value with the actual atmospheric N_2O mole fraction. During the cruise we measured a mean of 318 ppb (± 3 ppb, $n = 84$) in the atmosphere, which is in good agreement with the monthly mean of 318.5 ± 0.2 ppb in October 2003 measured at Mace Head, Ireland. This value was taken from the Advanced Global Atmospheric Gases Experiment (AGAGE) data set (updated version from May 2005, available at <ftp://cdiac.esd.ornl.edu> (subdirectory pub/ale_gage_Agage/Agage/gc-md/monthly) at the Carbon Dioxide Information Analysis Center in Oak Ridge, Tennessee).

The apparent oxygen utilization (AOU) was calculated as follows:

$$\text{AOU} [\mu\text{mol L}^{-1}] = \text{O}_2 (\text{equilibrium}) - \text{O}_2 (\text{observed}). \quad (3)$$

The equilibrium values of dissolved oxygen (O_2) were calculated with the equation given by [Weiss, 1970]. The concentration of H_2S is expressed as the negative oxygen equivalent ($1 \mu\text{mol L}^{-1} \text{H}_2\text{S} = 2.62 \mu\text{mol L}^{-1} \text{O}_2$). Dissolved nutrients and CTD data were provided by the participating working groups.

The N_2O inventory of single basins $m_{\text{N}_2\text{O}}$ was calculated as follows:

$$m_{\text{N}_2\text{O}} [\text{tons}] = \bar{c}_{\text{N}_2\text{O}} * n_{\text{N}_2\text{O}} * V * 10^{-3} \quad (4)$$

where $\bar{c}_{\text{N}_2\text{O}}$ is the mean measured N_2O concentration in the single basins from the upper halocline to the bottom [nmol L^{-1}], $n_{\text{N}_2\text{O}}$ is the mole weight of N_2O (44 g mol^{-1}) and V is the water volume of the single basins [km^3].

The water volumes are based on data published in chapter 4.4.1 [HELCOM, 1996], available at: www.vtt.fi/inf/baltic/balticinfo/index.html.

The N₂O content of basins was calculated with data of the following stations: Bornholm Basin: station 140, 200, 213, 222, eastern Gotland Basin: station 250, 259, 260, 272, western Gotland Basin: station 240, 245, 284. Station 286 is located in the northern part of the Gotland Basin and thus has not been taken into account.

Nitrification rates [N] were estimated for the Bornholm Basin and the eastern Gotland Basin.

$$N \left[\text{nmol L}^{-1} \text{d}^{-1} \right] = \frac{\Delta m_{\text{N}_2\text{O}}}{d_{\text{basin}} * V_{\text{basin}} * n * 10^{-9}} * r_{\text{N}_2\text{O}} \quad (5)$$

where $\Delta m_{\text{N}_2\text{O}}$ is the difference of calculated N₂O content of the basins before and after the inflow event in tons, d_{basin} is the number of days from the first observation of the intrusion of North Sea Water until our measurements (assumed by data of the cruise reports [Nausch, 2003a; Nagel, 2003; Feistel, 2003; Nausch, 2003b; Wasmund, 2003; Nausch, 2003c]).

V_{basin} is the calculated volume of the basins [km³] (based on data published in chapter 4.4.1 [HELCOM, 1996], available at: www.vtt.fi/inf/baltic/balticinfo/index.html), n is the mole weight of N₂O (44 g mol⁻¹), and $r_{\text{N}_2\text{O}}$ is the assumed N₂O release of 0.3 % in continental shelves during nitrification [Seitzinger and Kroeze, 1998].

3. Results

In order to account for the hydrographic characteristics of the Baltic Sea and the direction of the inflow of North Sea Water, we present the results according to the following classifications: I) well-mixed basins such as the Kiel, Lübeck and Pomeranian Bights and II) clearly stratified basins such as the Arkona, the Bornholm, the western and the eastern Gotland Basin (see Fig. 2). For each basin selected profiles are shown.

3.1 Well-mixed basins

At shallow stations, with depths < 30 m (station 10, 12, 22, 30, 41, 46, 121, 130, 133, 360, OB Boje, OB 4, Fig. 1), water masses were well mixed, and profiles showed nearly uniform vertical distributions of all parameters (Fig. 3a). Concentrations of N₂O were

near equilibrium; however the Pomeranian Bight (station 130, 133, OB Boje, OB 4) showed enhanced saturation values ($104.6 \pm 7.9 \%$) in comparison with the Kiel Bight (station 360) and the Lübeck and Mecklenburg Bight (station 10, 12, 22, 41, 46; $79.3 \pm 10.7 \%$). No correlations were found between ΔN_2O and other parameters like O_2 and NO_3^- (Fig. 3b-c).




Fig. 3: well mixed basins; a) left plot with profiles of N_2O , calculated N_2O equilibrium concentration, NO_3^- , NO_2^- at station 41 in the Mecklenburg Bight and right plot with profiles of temperature, salinity and oxygen at station 41 in the Mecklenburg Bight; b) ΔN_2O plotted against oxygen at all stations < 30 m; c) ΔN_2O plotted against NO_3^- at all stations < 30 m

3.2 Stratified basins

Basins with water depths > 30 m (Fig. 4-7) were clearly stratified into layers of well mixed Surface Water (sw), Winter (ww) and Bottom Water (bw) as described above. At several stations Bottom Water was affected by North Sea Water (abw), up to the Farö Deep in the northern part of the central Baltic Sea (Fig. 1, station 286) [Feistel *et al.*, 2003]. However, below 110 m the deep water of the Farö Deep was still anoxic, though with decreasing H_2S concentrations from 125 m to the bottom (Fig. 6a, lower profiles). Stations in the western Gotland Basin such as the Landsort Deep (station 284, Fig. 7a) or

the Karlsö Deep (station 245, not shown) were still unaffected by the inflow event, thus below 80 m H_2S concentrations were uniform.

3.2.1 Arkona Basin

In the Arkona Basin (stations 109 and 113 (Fig. 4a)), N_2O concentrations were constant and near equilibrium ($10.9 \pm 0.7 \text{ nmol L}^{-1}$) throughout the water column. In the Winter Water below the thermocline at 15 m O_2 concentrations decreased, associated with increasing NO_2^- , NO_3^- and NH_4^+ . $\Delta\text{N}_2\text{O}$ was slightly negatively correlated with O_2 (Fig. 4b), and positively correlated with NO_3^- (Fig. 4c). At the bottom below 40 m inflowing North Sea (arrow in Fig. 4a) water formed a 5 to 10 m thick oxygen enriched layer, however with no clear influence on the N_2O concentration.




Fig. 4: Arkona Basin; a) station 113 (Arkona Deep): left plot with profiles of N_2O , N_2O equilibrium concentration, NO_3^- , NO_2^- , right plot with profiles of temperature, salinity and oxygen, the arrow indicate the influence of North Sea Water; abbreviations see Fig. 2.; b) $\Delta\text{N}_2\text{O}$ plotted against oxygen (at all stations in the Arkona Basin, $y = -0.011x + 3.132$, $R^2 = 0.67$); c) $\Delta\text{N}_2\text{O}$ plotted against NO_3^- (at all stations in the Arkona Basin, $y = 0.546x - 0.807$, $R^2 = 0.66$)

3.2.2 Bornholm Basin

In the Bornholm Basin (Fig. 5, stations 140, 200, 213 and 222), N_2O profiles in the central basin (stations 200 (not shown) and 213 (Fig. 5a)) can be clearly distinguished from stations where water flows into and out of the basin. At station 140 (inflow, not shown) concentrations and distribution of N_2O and ΔN_2O were comparable to the Arkona Basin. At station 222 (outflow, not shown) N_2O concentrations in the surface layer were uniform near equilibrium at approximately 10 nmol L^{-1} , below the surface layer concentrations were uniform around 15.4 nmol L^{-1} . In the central Bornholm Basin, at station 200 (not shown) and 213 (Fig. 5a) N_2O concentrations increased rapidly within the layer affected by North Sea Water (abw, below 60 m), with N_2O values up to 31.3 nmol L^{-1} (station 200). These were the highest values measured during the entire cruise. In water masses above, N_2O was near equilibrium, with slightly enhanced ΔN_2O values in the 'old' Bottom Water (bw, 40 – 60 m). In the Bornholm Basin ΔN_2O was clearly negatively correlated with oxygen and positively with NO_3^- (Fig. 5b-c), however, both correlations were nonlinear and were fitted best by polynomials.





Fig. 5: central Bornholm Basin; a) station 213 (Bornholm Deep): left plot with profiles of N_2O , N_2O equilibrium concentration, NO_3^- , NO_2^- , right plot with profiles of temperature, salinity and oxygen, abbreviations see Fig. 2; b) ΔN_2O plotted against oxygen (at all stations in the Bornholm Basin, $y = 0.0003 x^2 - 0.1531 x + 19.517$, $R^2 = 0.88$); c) ΔN_2O plotted against NO_3^- (at all stations in the Bornholm Basin, $y = 0.0585 x^2 + 0.1438 x - 0.6155$, $R^2 = 0.90$)


3.2.3 Eastern Gotland Basin

The situation became more complex in the eastern Gotland basin (stations 259, 250, 260, 271 and 286). Profiles were not as homogeneous as in the Arkona or Bornholm Basin. Again, N_2O concentrations were near equilibrium in the surface layer (sw, 0 - 20/30 m) and the Winter Water (ww, 20/30 – 60 m). At station 271 (Fig. 6a, upper profiles) the Bottom Water (bw) was completely oxygenated, with N_2O values at approximately 20 nmol L^{-1} and positive ΔN_2O . At station 286 (Fig. 6a, lower profiles) the Bottom Water (bw) was affected by the North Sea Water too, but was still anoxic. Inflow of North Sea Water was detectable by decreasing H_2S concentrations down to the bottom. Throughout the Bottom Water N_2O concentrations remained near zero. At station 250 (not shown), 271 (Fig. 6a, upper profiles) and 286 (Fig. 6a, lower profiles) a sharp local minimum of N_2O concentrations was observed at depths between 90 and 110 m (see arrows in Fig. 6a), combined with a local minimum in NO_3^- values. Except for the anoxic water masses, ΔN_2O was linearly correlated with O_2 and NO_3^- (Fig. 6b-c).

a) station 271 (Gotland Deep)



station 286 (Farö Deep)







Fig. 6: Eastern Gotland Basin; a) station 271 (Gotland Deep, upper plots) and 286 (Farö Deep, lower plots): left plots with profiles of N_2O , N_2O equilibrium concentration, NO_3^- , NO_2^- ; right plots with profiles of temperature, salinity and oxygen, abbreviations see Fig. 2; b) $\Delta\text{N}_2\text{O}$ plotted against oxygen (at all stations in the Eastern Gotland Basin, $y = -0.019x + 5.625$, $R^2 = 0.67$ (except for $\text{O}_2 < 3 \mu\text{mol L}^{-1}$)); c) $\Delta\text{N}_2\text{O}$ plotted against NO_3^- (at all stations in the Eastern Gotland Basin, $y = 0.639x - 0.459$, $R^2 = 0.62$ (except for $\text{O}_2 < 3 \mu\text{mol L}^{-1}$))

3.2.4 Western Gotland Basin

The western Gotland Basin with stations 284 (Fig. 7a), 245 and 240 revealed the “old” conditions, showing characteristics as yet unaffected by the latest intrusion of oxidic North Sea Water. N_2O in the surface layer (sw, 0 - 20/40 m) and Winter Water (ww, 20/40 - 60 m) was near equilibrium. Below 50 m, oxygen concentrations decreased rapidly and N_2O concentrations dropped sharply within the oxidic/anoxic interface and remained near zero in the anoxic deep waters. $\Delta\text{N}_2\text{O}$ values were negative and were not correlated with NO_3^- (Fig. 7c). $\Delta\text{N}_2\text{O}$ was logarithmically correlated with oxygen (Fig. 7b).






Fig. 7: Western Gotland Basin; a) station 284 (Landsort Deep): left plot with profiles of N_2O , N_2O equilibrium concentration, NO_3^- , NO_2^- , right plot with profiles of temperature, salinity and oxygen, abbreviations see Fig. 2; b) $\Delta\text{N}_2\text{O}$ plotted against oxygen (at all stations in the Western Gotland Basin; $y = 2.2467 \ln(x) - 13.322$, $R^2 = 0.86$, (with exception of $\text{O}_2 < 0 \mu\text{mol L}^{-1}$)) c) $\Delta\text{N}_2\text{O}$ plotted against NO_3^- (at all stations in the Western Gotland Basin)

3.3 Estimated contribution of the North Sea Water inflow to the production of N_2O

The North Sea Water inflow consisted of a water volume of 200 km^3 [Feistel and Nausch, 2003]. With an assumed N_2O concentration of $10 \pm 2 \text{ nmol L}^{-1}$ [Law and Owens, 1990], the North Sea Water transported approximately $88 \pm 18 \text{ tons N}_2\text{O}$ into the Baltic Sea.

Before the North Sea Water inflow, the deep waters below the halocline were anoxic, not only in the western but also in the eastern Gotland Basin and the Bornholm Basin [Schmidt, 2002]. Thus, N_2O concentrations near zero similar to measured profiles in the western Gotland Basin in October 2003 (Fig. 7a) can be assumed. This is supported by the drop in concentration at station 286 (Fig. 7a, lower profile), which is assumed to be related to the previously anoxic bottom water. The mean N_2O concentration in the western Gotland Basin was $0.97 \pm 0.34 \text{ nmol L}^{-1}$, on the basis of these values the calculated N_2O content of the Bornholm Basin, the eastern and the western Gotland Basin was approximately $92 \pm 33 \text{ tons}$ before the inflow.

After the inflow event the Bornholm Basin and the eastern Gotland Basin are clearly influenced by the North Sea Water, whereas the western Gotland Basin was still unaffected [Nausch, 2003a; Nagel, 2003; Feistel, 2003; Nausch, 2003b; Wasmund, 2003; Nausch, 2003c]. The N_2O content of the basins, calculated with the mean of measured N_2O concentrations below the halocline in these basins, was about $1222 \pm 266 \text{ tons}$ (Table 1).

Table 1: Estimated N₂O content of single basins in the Baltic Sea below the halocline, before and after the inflow of North Sea Water in January 2003

	mean N ₂ O conc. below the halocline [nmol L ⁻¹]	Water volume [km ³]	N ₂ O content before the inflow event [tons]	N ₂ O content after the inflow event [tons]
Bornholm Basin	> 50 m 16.59 ± 5.61	306	13 ± 5	223 ± 76
eastern Gotland Basin	> 70 m 18.46 ± 3.43	1195	51 ± 18	971 ± 180
western Gotland Basin	> 70 m 0.97 ± 0.34	657	28 ± 10	28 ± 10
Σ		2158	92 ± 33	1222 ± 266

4. Discussion

Over the past two decades the previously frequent inflows of North Sea Water became rather rare [Feistel and Nausch, 2003], and oxygen levels in deep waters decreased. Thus, oxygen conditions in the Baltic Sea deep water cover a continuum from permanently oxic (i.e. Arkona Basin) to almost permanently anoxic conditions (i.e. western Gotland Basin), with changes at non-regular intervals between anoxic and oxic (i.e. Bornholm Basin, eastern Gotland Basin) [Feistel, 2003; Nausch, 2003a; Nausch, 2003b; Nausch, 2003c; Nagel, 2003; Wasmund, 2003].

The inflow event in January 2003 rapidly changed the environmental conditions of the deep basins. With respect to the oxygen dependent production of N₂O, our measured N₂O concentrations reflect the continuum of unaffected and changing oxygen conditions quite well. In oxic and well mixed waters, vertical N₂O profiles were homogenous, with concentrations near equilibrium (Fig. 3a). Anoxic deep water layers, unaffected by North Sea Water (i.e. in the western Gotland Basin), had N₂O concentrations near zero (Fig. 7a). Therefore, in both cases no correlations between N₂O and either oxygen or nitrate were found (Fig. 3b-c, Fig. 7b-c). In contrast, stratified and recently ventilated water bodies in the Bornholm and eastern Gotland Basin revealed N₂O distributions that were clearly correlated with oxygen and nitrate (Fig. 5b-c, Fig. 6b-c).

These vertical N₂O distributions are in general agreement with the few previously published N₂O profiles from the central Baltic Sea [Rönner, 1983; Rönner and Sörensson, 1985; Brettar and Rheinheimer, 1992]. However, the past environmental settings of the deep central Baltic Sea basins were different: N₂O profiles from the central Baltic Sea reported by [Rönner, 1983] were measured when oxic conditions prevailed during Au-

gust-September 1977 after a strong inflow event in 1976/1977 [Schinke and Matthäus, 1998]. Their N₂O profiles are comparable to our profiles, measured in the completely oxygenated Bornholm Basin during October 2003 (Fig. 5a). Anoxic conditions were re-established in July 1979 and May-June 1980. The shape of the N₂O profiles from the then anoxic Gotland Deep, measured by Rönner and Sörensson [1985] is comparable to the N₂O profiles measured in the western Gotland Basin (e.g., the Landsort Deep, Fig. 7a). This is the same for profiles measured by Brettar and Rheinheimer [1991] in August 1986 and July 1987 during the 1983-1993 stagnation periods [Schinke and Matthäus, 1998].

In the following sections we discuss the processes that may cause the observed distributions of N₂O in the different basins.

4.1 Non-biological aspects

In surface layers and well-mixed water bodies of shallow stations, observed N₂O concentrations were near the equilibrium due to exchange with the atmosphere. In the Winter Water N₂O concentrations were also near equilibrium, however with higher absolute values than in the surface layer (see Fig. 5a-7a). Mainly hydrographic aspects were here responsible for the observed N₂O distribution. This water mass is formed during winter convection, when N₂O concentrations were in equilibrium with the atmosphere and this signal is conserved during stratification of the upper layer in summer. The lower temperature and hence higher N₂O solubility during formation of the Winter Water are the reason for the enhanced N₂O concentrations in this layer.

A non-biological factor affecting N₂O in the deep water of the Baltic Sea might be advection with inflowing North Sea Water. Intrusion of N₂O by North Sea Water should be detectable at stratified stations, where the inflow of North Sea Water was clearly identified. In the Arkona Basin (station 109 and 113) this inflow was detectable at the bottom by lower temperature and higher oxygen concentrations; however, N₂O concentrations did not increase and remained close to equilibrium (Fig. 4a-b). These results point to only low advection of N₂O by North Sea Water, and are supported by measurements of [Law and Owens, 1990]. They found N₂O concentrations close to equilibrium up to approximately 10 nmol L⁻¹ in the North Sea. Thus, the enhanced N₂O values detected in layers affected by North Sea Water, for example in the Bornholm Basin (station 200 and 213), must originate from biological *in situ* production since the inflow, rather than advection.

4.2 Biological aspects

Previous studies demonstrated the existence of N₂O producing bacteria and investigated the biological pathways, namely nitrification and denitrification in the Baltic Sea [Bauer, 2003; Brettar and Höfle, 1993; Brettar et al., 2001]. Both processes are commonly inferred by correlations between N₂O and oxygen or nitrate [Yoshinari, 1976; Yoshida et al., 1989; Cohen and Gordon, 1978; Butler et al., 1989].

4.2.1 Anoxic waters

In general, in anoxic and H₂S containing bottom waters N₂O concentrations were constantly near zero, and therefore no correlation with either O₂ or NO₃⁻ was found. The N₂O production by nitrification and denitrification might probably be inhibited by the presence of H₂S [Joye and Hollibaugh, 1995; Knowles, 1982, Sørensen et al., 1980], and while changing to anoxic conditions, N₂O can be consumed during denitrification as an electron acceptor instead of oxygen [Elkins et al., 1978; Cohen and Gordon, 1978]. However, in contrast to other authors [Rönnner et al., 1983; Brettar and Rheinheimer, 1992] we found low and uniformly distributed concentrations of N₂O (up to 1.7 nmol L⁻¹) in the anoxic water masses, which may have been residuals of a previous production process during oxic conditions.

4.2.2 Suboxic waters

In suboxic waters and at the boundary to anoxic water masses N₂O is expected to be mainly produced by denitrification processes [Codispoti et al., 2001], usually indicated by decreasing NO₃⁻ concentrations and a secondary NO₂⁻ peak [Wrage et al., 2001; Kristiansen and Schaanning, 2002]. These indicators for denitrification were found only at the Farö Deep (station 286, 90m). However, no accumulation of N₂O was observed, rather a local minimum of N₂O was found (Fig. 6a, indicated by arrows). Hannig et al. [2005] recently investigated denitrification associated microorganisms in the Gotland Basin (station 271 and 286) in October 2003. They did not find denitrification activities in suboxic water masses, but a high denitrifying potential restricted to a narrow depth range at the oxic-anoxic interface and the sulfidic zone. However, in these depths an accumulation of N₂O was not found either.

The local minimum of N₂O was observed not only at the Farö Deep, but also at the Gotland Deep (Fig. 6a, indicated by arrows) and station 250 (profile not shown). A residual signal of the small inflow event in August 2002 could be observed at these depths be-

tween 90 and 110 m [Feistel *et al.*, 2003]. We assume that this minimum of N_2O is a previous signal of former anoxic bottom water, pushed up by the small inflow event in August 2002. The restriction of denitrification activity to a narrow depth range at anoxic-oxic boundaries was not only reported by Hannig *et al.* [2005] but also by Brettar *et al.* [2001]. Therefore, the lack of denitrification signals leads to the question, which processes might cause the measured N_2O concentrations.

4.2.3 Correlation between N_2O and O_2

In general, in oxic waters N_2O is positively correlated with nitrate, negatively with oxygen, indicating a production by nitrification. However, below a distinct threshold of oxygen concentration a clear inversion of relationship was found. Figure 8 shows the correlation between ΔN_2O and O_2 in the Baltic Sea. At O_2 concentrations $> 50 \mu\text{mol L}^{-1}$ ΔN_2O is clearly negatively correlated with O_2 , indicating production by nitrification (see Fig. 8, green data points). At O_2 concentrations $< 20 \mu\text{mol L}^{-1}$ ΔN_2O and O_2 were significantly positively correlated (see Fig. 8, red data points), data between $20 \mu\text{mol L}^{-1}$ and $50 \mu\text{mol L}^{-1}$ were extremely scattered (see Fig. 8, black data points).




Fig. 8. Correlation between ΔN_2O and O_2 in the Baltic Sea. Correlations were calculated for oxic waters with O_2 concentrations $> 50 \mu\text{mol L}^{-1}$ (green coloured, $y = -0.019x + 5.41$, $R^2 = -0.70$) and $< 20 \mu\text{mol L}^{-1}$ (red coloured, $y = 1.038x - 11.36$, $R^2 = 0.81$). These concentrations were empirically tested and gave the best fittings for both correlations.

These findings suggest a change in N_2O converting processes. Particularly in environments with rapidly changing conditions it is advantageous for microorganisms to be able to switch between different metabolic pathways. The change between aerobic and an-

aerobic metabolism and thus the yield of N₂O during these processes is probably controlled particularly by the O₂ concentration, although little is known about the detailed mechanisms [Baumann *et al.*, 1996; John, 1977; Sørensen, 1987]. Our results suggest a production of N₂O during nitrification until an oxygen threshold of around 20 - 50 µmol L⁻¹, whereas the exact concentration is not to be determined due to the scattered data. Below this threshold N₂O seemed to be degraded; probably used as an electron acceptor instead of oxygen and thereby reduced to N₂ [Elkins *et al.*, 1978; Cohen and Gordon, 1978]. In the literature, threshold values of 2 µmol L⁻¹ for nitrification are reported [Carlucci and McNally, 1969; Gundersen *et al.*, 1966]. For several nitrifiers the ability to switch between 'classical' nitrification, nitrifier-denitrification and aerobic denitrification was shown [Wrage *et al.*, 2001, Whittaker *et al.*, 2000, Zart *et al.*, 2000, Zehr and Ward, 2002]. The oxygen sensitivity is species-specific and also enzyme-specific; therefore the scatter of data might reflect the variety of involved species and enzymes [Jiang and Bakken, 2000; Goreau *et al.*, 1980; Wetzel, 1983; Robertson *et al.*, 1988; Richardson, 2000]. Bauer [2003] investigated NH₄⁺ oxidizing bacteria in the eastern Gotland Basin, and found similar bacterial communities at different depths; their nitrification activities however depended on O₂ concentrations.

Therefore, the ability of nitrifiers to perform denitrifying processes and the lack of 'classical' denitrifying signals, a switch of N₂O producing processes by nitrifiers can be assumed. These findings are in agreement with the assumptions of Rønner [1983], who also assumed, that nitrification is the main N₂O production pathway in the Baltic Sea.

Alternatively, it is also possible to interpret the data from the hydrographical or temporal point of view. Figure 9 shows the same data set as shown in Fig. 8. This time the data set is grouped not according to the oxygen concentrations but to the affiliation to different basins. Station 286 was excluded due to its transitional character. At this station anoxic conditions in the deep waters were found similar to other stations in the western Gotland Basin, but H₂S concentrations were decreasing towards the bottom. This indicates beginning ventilation, however still too weak to lead to oxic conditions.




Fig. 9. Correlation between $\Delta\text{N}_2\text{O}$ and O_2 in the Baltic Sea. Correlations were calculated for the Bornholm Basin (station 140, 200, 213, 222, green coloured, $y = -6.83 \ln(x) + 37.88$, $R^2 = 0.86$), the eastern Gotland Basin (station 259, 250, 260, 271, blue coloured, $y = -0.02x + 5.88$, $R^2 = 0.70$) and the western Gotland Basin (station 284, 240, 245, red coloured, $y = 2.25 \ln(x) - 13.32$, $R^2 = 0.86$). Anoxic data and station 286 were excluded.

In the stratified basins such as the Bornholm Basin, and the eastern and western Gotland Basin correlations of $\Delta\text{N}_2\text{O}$ and O_2 were regionally different and not always linear (Fig. 5b-c, 6b-c, 7a, 9). Particularly in the Bornholm Basin, N_2O and oxygen as well as N_2O and nitrate showed significant non-linear relationships (Fig. 5b-c, 9). The Bornholm Basin, which was anoxic before the inflow [Schmidt, 2002], was ventilated by North Sea Water in January 2003, months before the northern part of the eastern Gotland Basin was affected by the inflow [Nausch, 2003a, Nausch et al., 2004]. In October 2003 the oxygen conditions were already switching back to suboxic conditions [Nausch, 2003c; Wassmund, 2003], visible by decreasing oxygen concentrations compared to the beginning of the year. Accordingly the duration of elevated oxygen concentration in the respective basins may contribute to the observed accumulation of N_2O . In the eastern Gotland Basin (Fig. 6b-c, 9) the anoxic conditions changed a few months after the Bornholm Basin: the Gotland Deep was ventilated by North Sea Water in May 2003 [Nausch, 2003b]. Thus, there was less time for N_2O accumulation. For various communities of NH_4^+ oxidizing bacteria different lag times after switching from anoxic to oxic incubations were shown and the production of N_2O might not have started directly after the ventilation by North Sea Water [Bodelier et al., 1996]. In the western Gotland Basin (Fig. 7b-c, 9) no ventilation by North Sea Water had occurred by October 2003, therefore degradation of N_2O at

the oxic-anoxic interface was found. We suspect that the correlation between $\Delta\text{N}_2\text{O}$ and O_2 in the Bornholm Basin and the eastern Gotland Basin will become similar to those of the western Gotland Basin with time, when the conditions change to anoxic.

Summarizing, we assume a conversion of N_2O mainly by nitrifiers, depending on oxygen concentration and with significant spatial and temporal characteristics.

4.3 Estimated contribution of the North Sea Water inflow to the production of N_2O

The estimated N_2O content in the stratified basins showed distinctly higher values after the inflow of the North Sea Water than before. The N_2O concentration in the North Sea Water was assumed to be near equilibrium, so there was no significant advection of N_2O from the North Sea. Thus, the observed elevated N_2O concentrations in the Baltic Sea basins result from a stimulation of N_2O production by the inflow, most likely by advection of oxygen (see Table 1).

Although more than 1000 tons of N_2O were produced, it is questionable whether the North Sea Water inflow makes the Baltic Sea a source of atmospheric N_2O . Due to the strong salinity stratification, it can be assumed that the formed N_2O stays below the permanent halocline, and therefore it will not reach the atmosphere. Commonly N_2O budgets were modelled as a function of nitrification and denitrification. *Seitzinger and Kroeze [1998]* modelled the distribution of N_2O production, amongst others based on the input of nitrogen compounds into estuaries by rivers. However, estimations of global N_2O emissions do not or only to a small extent take into account the hydrographic aspects. The stratification of the water column probably lead to a reduced release of calculated amounts, and accordingly to an overestimation of N_2O emissions.

The assumption of remaining N_2O below the halocline leads to the question, whether and to what extent the nitrogen cycle might be influenced.

Based on the calculated N_2O content of the basins and the assumption of nitrification as the main production pathway N_2O production rates and nitrification rates were estimated (Table 2). These nitrification rates are in good agreement with previously published rates for the Baltic Sea [*Enoksson, 1986; Bauer 2003*]. *Bauer [2003]* calculated for the eastern Gotland Basin mean nitrification rates of $21.6 \pm 11.1 \text{ nmol L}^{-1}$ at 60 m depth, respectively $44.3 \pm 33.1 \text{ nmol L}^{-1}$ at 100 m depth.

Table 2: Estimated nitrification rates in the Bornholm Basin and the eastern Gotland Basin, based on the assumption of 0.3 % N₂O release during nitrification [Seitzinger and Kroeze, 1998]

	$\Delta m_{\text{N}_2\text{O}}$ [tons]	d_{basin} [day]	Water volume [km ³]	N ₂ O production rate [nmol L ⁻¹ d ⁻¹]	nitrification rate [nmol L ⁻¹ d ⁻¹]
Bornholm Basin	220 ± 81	265	306	0.059 ± 0.023	19.62 ± 7.57
eastern Gotland Basin	920 ± 198	167	1195	0.105 ± 0.021	34.92 ± 6.87

These rates are comparably low [e.g. *Bianchi et al., 1999*]; therefore the influence on the nitrogen cycle in the Baltic Sea might be small, too. N₂O might play a minor role as a reserve- or buffer-molecule during the change to anoxic conditions, preserving nitrifying processes in exchange for oxygen for a short while.

5. Conclusions

In January 2003 a major inflow of cold, highly saline and oxygen-rich North Sea Water was observed, terminating the ongoing stagnation period in parts of the central Baltic Sea.

- ⊕ In agreement with previous studies, we found N₂O production mainly in oxic water masses below the Winter Water layer.
- ⊕ We found no indication for advection of N₂O by North Sea Water; however, the environmental conditions for N₂O production were clearly changed due to the North Sea Water inflow.
- ⊕ The inflow leads to a stimulation of N₂O production below the permanent halocline, but due to the halocline, the Baltic Sea is not a significant source of N₂O to the atmosphere.
- ⊕ There was no indication for an accumulation of N₂O during denitrification. In oxic and suboxic water masses nitrification seems to be the main production pathway. The occurrence of nitrifier-denitrification and aerobic denitrification is possible, but needs further investigations.

Acknowledgements

We thank the officers and crew of R/V Gauss for their excellent support. We especially thank R. Hoffmann [MPI for Chemistry, Mainz] for the calibration of the standard gas mixtures, and the colleagues from the IOW for providing the CTD data and the nutrients. The investigations were financially supported by the Deutsche Forschungsgemeinschaft through grant WA1434/1. The German part of the HELCOM COMBINE program is conducted by the Leibniz Institut für Ostseeforschung Warnemünde on behalf of the Bundesamt für Seeschifffahrt und Hydrographie, Hamburg and is funded by the Bundesministerium für Verkehr, Bau und Stadtentwicklung, Berlin.

References

- Bange, H.W., S. Rapsomanikis, and M.O. Andreae, Nitrous oxide in coastal waters, *Global Biogeochemical Cycles*, 10 (1), 197-207, 1996.
- Bange, H.W., S. Rapsomanikis, and M.O. Andreae, Nitrous oxide cycling in the Arabian Sea, *Journal of Geophysical Research-Oceans*, 106 (C1), 1053-1065, 2001.
- Bauer, S., Structure and function of nitrifying bacterial communities in the Eastern Gotland Basin (Central Baltic Sea), *Rostock, Univ., Diss., 2003, H 2003 B 4373*, 2003.
- Baumann, B., M. Snozzi, A.J.B. Zehnder, and J.R. van der Meer, Dynamics of denitrification activity of *Paracoccus denitrificans* in continuous culture during aerobic-anaerobic changes, *Journal of Bacteriology*, 178 (15), 4367-4374, 1996.
- Bianchi, M., C. Fosset, and P. Conan, Nitrification rates in the NW Mediterranean Sea, *Aquatic Microbial Ecology*, 17(3), 267-278, 1999.
- Bodelier, P.L.E., J.A. Libochant, C.W.P. Blom, H.J. Laanbroek, Dynamics of nitrification and denitrification in root-oxygenated sediments and adaptation of ammonia-oxidizing bacteria to low-oxygen or anoxic habitats, *Applied and Environmental Microbiology*, 62, 4100-4107, 1996.
- Brettar, I., and M.G. Höfle, Nitrous oxide producing heterotrophic bacteria from the water column of the central Baltic: abundance and molecular identification, *Marine Ecology Progress Series*, 94, 253-265, 1993.
- Brettar, I., E.R.B. Moore, and M.G. Höfle, Phylogeny and abundance of novel denitrifying bacteria isolated from the water column of the central Baltic Sea, *Microbial Ecology*, 42 (3), 295-305, 2001.

- Brettar, I., and G. Rheinheimer, Denitrification in the central Baltic: evidence for hydrogen sulfide oxidation as motor of denitrification at the oxic-anoxic interface, *Marine Ecology Progress Series* 77 (2-3), 157-169, 1991.
- Brettar, I., and G. Rheinheimer, Influence of carbon availability on denitrification in the Central Baltic Sea, *Limnology and Oceanography*, 37 (6), 1146-1163, 1992.
- Butler, J.H., J.W. Elkins, T.M. Thompson, and K.B. Egan, Tropospheric and dissolved N₂O of the West Pacific and East Indian Oceans during the El-Nino Southern Oscillation event of 1987, *Journal of Geophysical Research-Atmospheres*, 94 (D12), 14865-14877, 1989.
- Carlucci, A.F. and P.M. McNally, Nitrification by marine bacteria in low concentrations of substrate and oxygen, *Limnology and Oceanography*, 14, 736-739, 1969.
- Codispoti, L.A., J.A. Brandes, J.P. Christensen, A.H. Devol, S.W.A. Naqvi, H.W. Paerl, and T. Yoshinari, The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the anthropocene?, *Scientia Marina*, 65, 85-105, 2001.
- Codispoti, L.A., T. Yoshinari, and A.H. Devol, Suboxic respiration in the oceanic water column, in *Respiration in aquatic ecosystems*, edited by P.A. Del Giorgio, and P.J.I.B. Williams, pp. 225-247, Oxford University Press, Oxford, 2005.
- Cohen, Y., and L.I. Gordon, Nitrous oxide in the oxygen minimum of the eastern tropical North Pacific: Evidence for its consumption during denitrification and possible mechanisms for its production, *Deep-Sea Research*, 25 (6), 509-524, 1978.
- Dahlke, S., S. Wolff, L.-A. Meyer-Reil, H.W. Bange, R. Ramesh, S. Rapsomanikis and M.O. Andreae, Bodden waters (southern Baltic Sea) as a source of methane and nitrous oxide, in *Proceedings in Marine Sciences, Volume 2: Muddy Coast Dynamics and Resource Management*, edited by B.W. Flemming, M.T. Delafontaine, and G. Liebezeit, pp. 137-148, Elsevier Science, Amsterdam, 2000
- Elkins, J.W., S.C. Wofsy, M.B. McElroy, C.E. Kolb, and W.A. Kaplan, Aquatic sources and sinks for nitrous oxide, *Nature*, 275 (5681), 602-606, 1978.
- Enoksson, V., Nitrification rates in the Baltic Sea: Comparison of three isotope techniques, *Applied and Environmental Microbiology*, 51(2), 244-250, 1986.
- Feistel, R., *IOW Cruise report 11/03/02, March 2003*, 2003.
<http://www.io-warnemuende.de/projects/monitoring/documents/cr110302.pdf>
- Feistel, R., G. Nausch, W. Matthäus, and E. Hagen, Temporal and spatial evolution of the Baltic deep water renewal in spring 2003, *Oceanologia*, 45 (4), 623-642, 2003.
- Feistel, R. and G. Nausch, Water exchange between the Baltic Sea and the North Sea and conditions in the deep basins, *HELCOM indicator fact sheets / Baltic Marine Environment Protection Commission – Helsinki Commission*, 2003.
<http://www.helcom.fi/environment/indicators2003/inflow.html>

- Goreau, T.J., W.A. Kaplan, S.C. Wofsy, M.B. McElroy, F.W. Valois, and S.W. Watson, Production of nitrite and nitrous oxide by nitrifying bacteria at reduced concentrations of oxygen, *Applied and Environmental Microbiology* 40 (3), 526-532, 1980.
- Gundersen, K., A.F. Carlucci, and K. Boström, Growth of some chemoautotrophic bacteria at different oxygen tensions, *Experientia*, 22, 229-230, 1966.
- Hannig, M., G. Braker, G. Lavik, M. Kuypers, J. W. Dippner, K. Jürgens, Structure and activity of denitrifying bacteria in the water column of the Gotland Basin (Baltic Sea), *Abstract presented at the SPOT-ON conference 2005, Warnemünde, June 26 – July 1, 2005*.
- HELCOM, Third periodic assessment of the state of the marine environment of the Baltic Sea 1989-1993, 64b, 1996.
<http://www.baltic.vtt.fi/balticinfo/index.html>
- Jiang, Q.Q., and L.R. Bakken, Nitrous oxide production and methane oxidation by different ammonia-oxidizing bacteria, *Applied and Environmental Microbiology*, 65, 2679-2684, 2000.
- Jørgensen, B B, and J. Sørensen, *Seasonal cycles of O₂, NO₃⁻ and SO₄²⁻ reduction in estuarine sediments: The significance of a NO₃⁻ reduction maximum in spring*, Marine Ecology Progress Series, 24, 65-74, 1985.
- John, P., Aerobic and anaerobic bacterial respiration monitored by electrodes, *Journal of General Microbiology*, 98, 231-238, 1977.
- Joye, S.B., and J.T. Hollibaugh, Influence of sulfide inhibition of nitrification on nitrogen regeneration in sediments, *Science*, 270, 623-625, 1995.
- Knowles, R. (1982), Denitrification, *Microbiological Reviews*, 46, 43-70.
- Kristiansen, S., and M.T. Schaanning, Denitrification in the water column of an intermitently anoxic fjord, *Hydrobiologia*, 469, 77-86, 2002.
- Law, C.S., and N.J.P. Owens, Denitrification and nitrous oxide in the North Sea, *Netherlands Journal of Sea Research*, 25 (1-2), 65-74, 1990.
- Meier, M.H.E., Modeling the age of Baltic seawater masses: Quantification and steady state sensitivity experiments, *Journal of Geophysical*, 110 (C02006), 2005.
- Nagel, K., IOW Cruise report 11/03/01, February 2003, 2003.
<http://www.io-warnemuende.de/projects/monitoring/documents/cr110301.pdf>
- Naqvi, S.W.A., D.A. Jayakumar, P.V. Narvekar, H. Naik, V. Sarma, W. D'Souza, S. Joseph, and M.D. George, Increased marine production of N₂O due to intensifying anoxia on the Indian continental shelf, *Nature*, 408 (6810), 346-349, 2000.
- Nausch G., IOW Cruise Report 40/03/22, January 2003, 2003a.
<http://www.io-warnemuende.de/projects/monitoring/documents/cr400322.pdf>

- Nausch G., *IOW Cruise Report 44/03/03, May 2003*, 2003b.
<http://www.io-warnemuende.de/projects/monitoring/documents/cr440303.pdf>
- Nausch, G., *Cruise report 11/03/04, October 2003*, 2003c.
<http://www.io-warnemuende.de/projects/monitoring/documents/cr110304.pdf>
- Nausch, G., R. Feistel, H.-U. Lass, K. Nagel, and H. Siegel, Hydrographisch-chemische Zustandseinschätzung der Ostsee 2003, *Meereswissenschaftliche Berichte*, 59 (1), 1-80, 2004.
- Nausch, G., W. Matthäus, and R. Feistel, Hydrographic and hydrochemical conditions in the Gotland Deep area between 1992 and 2003, *Oceanologia*, 45 (4), 557-569, 2003.
- Ostrom, N.E., M.E. Russ, B. Popp, T.M. Rust, and D.M. Karl, Mechanisms of nitrous oxide production in the subtropical North Pacific based on determinations of the isotopic abundances of nitrous oxide and di-nitrogen, *Chemosphere: Global Change Science*, 2 (3-4), 281-290, 2000.
- Popp, B.N., M.B. Westley, S. Toyoda, T. Miwa, J.E. Dore, N. Yoshida, T.M. Rust, F.J. Sansone, M.E. Russ, N.E. Ostrom, and P.H. Ostrom, Nitrogen and oxygen isotopic constraints on the origins and sea-to-air flux of N₂O in the oligotrophic subtropical North Pacific gyre, *Global Biogeochemical Cycles*, 16 (4), doi: 10.1029/2001GB001806, 2002.
- Prather, M., D. Ehhalt, F. Dentener, R. Derwent, E. Dlugokencky, E. Holland, I. Isaksen, J. Katima, V. Kirchhoff, P. Matson, P. Midgley, and M. Wang, Atmospheric chemistry and greenhouse gases, in *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by J.T. Houghton, Y. Ding, D.J. Griggs, M. Noguer, P.J. Van der Linden, X. Dai, K. Maskell, and C.A. Johnson, pp. 239-287, Cambridge University Press, Cambridge, UK, 2001.
- Richardson, D.J., Bacterial respiration: A flexible process for a changing environment, *Microbiology*, 146, 551-571, 2000.
- Robertson, L.A., and J.G. Kuenen, Aerobic denitrification - Old wine in new bottles, *Antonie Van Leeuwenhoek Journal of Microbiology*, 50 (5-6), 525-544, 1984.
- Robertson, L.A., E.W.J. Vanniell, R.A.M. Torremans, and J.G. Kuenen, Simultaneous nitrification and denitrification in aerobic chemostat cultures of *Thiosphaera pantotropha*, *Applied and Environmental Microbiology*, 54 (11), 2812-2818, 1988.
- Rönner, U., Distribution, production and consumption of nitrous oxide in the Baltic Sea, *Geochimica et Cosmochimica Acta*, 47, 2179-2188, 1983.
- Rönner, U., F. Sörensson, and O. Holmhansen, Nitrogen assimilation by phytoplankton in the Scotia Sea, *Polar Biology*, 2 (3), 137-147, 1983.
- Rönner, U., and F. Sörensson, Denitrification rates in the low-oxygen waters of the stratified Baltic Proper, *Applied and Environmental Microbiology*, 50, 801-806, 1985.

- Schinke, H., and W. Matthäus, On the causes of major Baltic inflows - an analysis of long time series, *Continental Shelf Research*, 18, 67-97, 1998.
- Schmidt, M., *IOW Cruise Report 11/02/03, October 2002*, 2002.
<http://www.io-warnemuende.de/projects/monitoring/documents/cr110203.pdf>
- Seitzinger, S.P., and C. Kroeze, Global distribution of nitrous oxide production and N inputs in freshwater and coastal marine ecosystems, *Global Biogeochemical Cycles*, 12 (1), 93-113, 1998.
- Seitzinger, S.P., C. Kroeze, and R.V. Styles, Global distribution of N₂O emissions from aquatic systems: Natural emissions and anthropogenic effects, *Chemosphere: Global Science Change*, 2, 267-279, 2000.
- Sørensen, J., Nitrate reduction in marine sediment: Pathways and interactions with iron and sulfur cycling, *Geomicrobiology Journal*, 5 (3-4), 401-422, 1987.
- Sørensen, J., J.M. Tiedje, R.B. Firestone, Inhibition by sulfide of nitric and nitrous oxide reduction by denitrifying *Pseudomonas fluorescens*, *Applied and Environmental Microbiology*, 39(1), 105-108, 1980.
- Vollack, K.U., and W.G. Zumft, Nitric oxide signaling and transcriptional control of denitrification genes in *Pseudomonas stutzeri*, *Journal of Bacteriology*, 183 (8), 2516-2526, 2001.
- Wasmund, N., *IOW Cruise Report 44/03/07, July-August 2003*, 2003
<http://www.io-warnemuende.de/projects/monitoring/documents/cr440307.pdf>
- Weiss, R.F., The solubility of nitrogen, oxygen and argon in water and seawater, *Deep-Sea Research*, 17, 721-735, 1970.
- Weiss, R.F., and B.A. Price, Nitrous oxide solubility in water and seawater, *Marine Chemistry*, 8, 347-359, 1980.
- Wetzel, R.G., *Limnology*, Saunders College Publishing, Philadelphia, Pa., 1983.
- Whittaker, M., D. Bergnamm, D. Arciero, and A.B. Hooper, Electron transfer during the oxidation of ammonia by the chemolithotrophic bacterium *Nitrosomonas europaea*, *Biochimica et Biophysica Acta*, 1459, 346-355, 2000.
- WMO, *Scientific assessment of ozone depletion: 2002*, pp. 498, WMO (World Meteorological Organization), Geneva, 2003.
- Wrage, N., G.L. Velthof, M.L. van Beusichem, and O. Oenema, Role of nitrifier denitrification in the production of nitrous oxide, *Soil Biology and Biochemistry*, 33 (12-13), 1723-1732, 2001.
- Yamagishi, H., N. Yoshida, S. Toyoda, B.N. Popp, M.B. Westley, and S. Watanabe, Contributions of denitrification and mixing on the distribution of nitrous oxide in the North Pacific, *Geophysical Research Letters*, 32 (4), 2005.

Yoshida, N., H. Morimoto, M. Hirano, I. Koike, S. Matsuo, E. Wada, T. Saino, and A. Hattori, Nitrification rates and ^{15}N abundances of N_2O and NO_3^- in the western North Pacific, *Nature*, 342, 895-897, 1989.

Yoshinari, T., Nitrous oxide in the sea, *Marine Chemistry*, 4, 189-202, 1976.

Yoshinari, T., M.A. Altabet, S.W.A. Naqvi, L. Codispoti, A. Jayakumar, M. Kuhland, and A. Devol, Nitrogen and oxygen isotopic composition of N_2O from suboxic waters of the eastern tropical North Pacific and the Arabian Sea - Measurement by continuous-flow isotope-ratio monitoring, *Marine Chemistry*, 56 (3-4), 253-264, 1997.

Zart, D., I. Schmidt, and E. Bock, Significance of gaseous NO for ammonia oxidation by *Nitrosomonas eutropha*, *Antonie van Leeuwenhoek*, 77, 49-55, 2000.

Zehr, J.P., and B.B. Ward, Nitrogen cycling in the ocean: New perspectives on processes and paradigms, *Applied and environmental microbiology*, 68 (3), 1015-1024, 2002.

Chapter 6

Analysis of bacterial communities in the subtropical North Atlantic by 16S rRNA genes

Sylvia Walter, Jörg Süling, Marcus Tank, Hermann W. Bange, Johannes F. Imhoff

Abstract

The vertical composition of bacterial communities in the oceans is largely unknown, though bacteria play an important role in biogeochemical cycles, beside others in the production and consumption of greenhouse gases such as methane or nitrous oxide. Studies showed possible correlations between single picoplankton species and chemical compounds, i.e. between species related to the genus *Roseobacter* and the turnover of DMSP to DMS. Therefore, the hypothesis was that it might also be possible to find species which are associated with the distribution of nitrous oxide, an atmospheric trace gas with considerable relevance for the Earth's climate. We investigated the vertical distribution of bacteria in the western and eastern subtropical North Atlantic (Meteor 60-5 cruise), identified by 16S rRNA gene sequences, with the intention to find correlations between the community structure of bacteria and the distribution of nitrous oxide. In the subtropical North Atlantic N₂O is assumed to be mainly produced during nitrification. However, the functional genes encoding the enzymes for the conversion of N₂O during nitrification are unknown.

The yield of N₂O depends on environmental conditions and also on the species involved in conversion processes. *Proteobacteria* were found to be the predominant group in both basins. In the western basin the contribution of *Proteobacteria* increased significantly with depth, and also the composition changed with depth: α -*Proteobacteria* were predominant at shallower depth and γ -*Proteobacteria* increased in proportion in deeper water masses. Particularly *Acinetobacter* and *Alteromonas* seem to become more important with depth. In contrast, in the eastern basin *Proteobacteria* were also the predominant group of identified bacteria, but with minor proportions. Here no depth depending community structures were found. In both basins β - and δ -*Proteobacteria* hardly occurred. Particularly *Proteobacteria* are known to perform N₂O producing processes, however, we found no obviously predominant bacterial group related to the distribution of N₂O.

1. Introduction

Pelagic bacteria are a major component of biomass in oceanic oligotrophic systems [Binder *et al.*, 1996; Cho and Azam, 1990; Carlson *et al.*, 1996]. It is generally accepted that major bacterioplankton groups show a vertical stratification [Lee and Fuhrman, 1991; Field *et al.*, 1997], and community structures shift with changing environmental factors [Bollmann and Laanbroek, 2002]. Some studies showed that correlations between single picoplankton species and chemical compounds (e.g. species of *Roseobacter* and Dimethylsulfoniopropionate, DMSP) can be found [Zubkov *et al.*, 2001; González *et al.*, 2000]. Thus, there might exist a similar pattern for nitrous oxide (N₂O). N₂O is an atmospheric trace gas with a high climatic relevance. It is an intermediary and a by-product in several processes of the nitrogen cycle. The contribution of single processes of the nitrogen cycle to the global N₂O budget is discussed controversially [Codispoti *et al.*, 2001; Popp *et al.*, 2002; Yamagishi *et al.*, 2005], and the knowledge of biological processes and controlling factors of the production and conversion of N₂O in situ are still unsatisfactory.

Up to now nitrification and denitrification are commonly accepted to be the main N₂O producing processes in the nitrogen cycle [Bonin *et al.*, 2002]. In general, nitrification is an aerobic process in which organic or inorganic nitrogen compounds are oxidized to nitrate (NO₃⁻) via hydroxylamine (NH₂OH) and nitrite (NO₂⁻). In this process N₂O is considered as a by-product. It is assumed that N₂O is either produced by an incomplete conversion of NH₂OH or by oxidation of NO₂⁻ [Ostrom *et al.*, 2000; Popp *et al.*, 2002]. Denitrification is the reduction of NO₃⁻ to N₂, with N₂O as a regular intermediate product [Naqvi *et al.*, 2000]. The yield of N₂O strongly depends on abiotic factors like concentrations of oxygen and nitrate but is also species specific [Brettar and Rheinheimer, 1991; Goreau *et al.*, 1980; Vollack and Zumft, 2001; Wetzel, 1983; Abou Seada and Ottow, 1985].

1.1 Bacteria involved in the production of N₂O

1.1.1 Nitrifying bacteria

Nitrification can be conducted by both autotrophic and heterotrophic organisms. Typically two groups of microorganisms are involved in autotrophic nitrification: NH₄⁺-oxidizing bacteria or primary nitrifiers and NO₂⁻-oxidizing bacteria or secondary nitrifiers [Bock *et al.*, 1986]. Both groups are affiliated with *Proteobacteria*. NH₄⁺-oxidizing bacteria, obligatory lithotrophic, belong to the β- and γ-*Proteobacteria*, divided into the genera *Nitrosomonas*, *Nitrosococcus*, and *Nitrospira* (including the genera *Nitrospira*, *Nitrosovibrio* and *Nitrosolobolus*) [Bothe *et al.*, 2000; Krümmel and Harms, 1982]. NO₂⁻-oxidizing bacteria belong to the α-*Proteobacteria* [Schramm *et al.*, 1998; Woese *et al.*, 1984a; Woese *et al.*, 1984b; Woese *et al.*, 1985], divided into the genera *Nitrobacter*, *Nitrospina*, *Nitrococcus* and *Nitrospira*. Among these, *Nitrobacter* and *Nitrospira* strains are capable of heterotrophic or mixotrophic growth, the rest being strict chemolithotrophs [Bartosch *et al.*, 1999]. The *Nitrospina* group was shown to be affiliated with the δ-subdivision of the *Proteobacteria* but did not share a specific relationship to each other or to other members of the δ-subdivision [Abeliovich, 2001]. Formerly autotrophic nitrifiers were classified as nitrifying bacteria in the family *Nitrobacteraceae* [Buchanan, 1917], but phylogenetically they are not closely related to each other [Woese *et al.*, 1984a; Woese *et al.*, 1984b; Woese *et al.*, 1985]. Both groups are widely distributed in marine environments [Venter *et al.*, 2004; Gallagher *et al.*, 2004; Ward and O'Mullan, 2002; Morris *et al.*, 2002].

Besides autotrophic nitrifiers also heterotrophic nitrifiers are known, which are phylogenetically more widespread and include a variety of species [Mével and Prieur, 2000]. They are unspecialized heterotrophic microorganisms for which their nitrogen oxidation property is not regarded as taxonomically significant [Focht and Verstraete, 1977].

1.1.2 Denitrifying bacteria

Denitrification reactions are widely distributed in various subclasses of the *Proteobacteria*, but also organisms of other eubacterial and archaeal genera are able to denitrify [Chen *et al.*, 2002; Shapleigh, 2001; Cabello *et al.*, 2004]. 16S rRNA and functional gene phylogenies are not congruent for denitrifiers, implying substantial horizontal gene transfer of the functional genes over time [Zehr and Ward, 2002; Ward, 2005]. However, this topic is discussed controversially [Petri and Imhoff, 2000].

Nitrifiers possess the ability to perform denitrifying processes as well, so called nitrifier-denitrification and aerobic denitrification [Wrage *et al.*, 2001]. Under aerobic conditions, both processes can be performed by either autotrophic NH_4^+ oxidizing bacteria or heterotrophic nitrifiers [Wrage *et al.*, 2001; Whittaker *et al.*, 2000; Zart *et al.*, 2000; Zehr and Ward, 2002]. Although the processes have identical educts and products as 'classical' denitrification, they differ by the structure of used enzymes and their genetics [Moreno-Vivián *et al.*, 1999; Richardson and Watmough, 1999; Robertson and Kuenen, 1988]. Many organisms are able to use different enzymes for similar reactions or to switch between these pathways depending on the environmental conditions [Moir *et al.*, 1995; Richardson, 2000; Wrage *et al.*, 2001].

Our studies focused on the vertical structure of bacterial communities in the subtropical North Atlantic, investigated by 16S rRNA genes. Aim was to obtain more information about the vertical distribution of bacteria, with the intention to find correlations between the structure of bacterial communities and the distribution of the greenhouse gas N_2O .

2. Methods

2.1 Sampling location

Samples were taken in the subtropical North Atlantic during March / April 2004 onboard the research vessel 'Meteor'. The cruise (M 60-5) started in Fort de France, Martinique (French Antilles) in the western part of the Atlantic and ended in Lisbon (Portugal).




Fig. 1: Cruise track of M 60-5 in the subtropical North Atlantic; numbers mark sampled stations for molecular analysis and N_2O profiles, dots mark sampled stations only for N_2O profiles

2.2 N_2O measurements

Water samples for N_2O analysis were collected in triplicate at 37 stations from various depths, taken with a 24-Niskin-bottle rosette, equipped with a CTD-probe. N_2O was measured directly on board with a gaschromatograph, equipped with an ECD (Electron Capture Detector). The analytical method applied is described by [Walter et al. \[2005\]](#).

2.3 Sampling for molecular analysis

70 water samples were taken with a 24-Niskin-bottle rosette at 8 stations (see Fig. 1 for station numbers), at depths from the surface to the bottom. Depths were selected according to the shape of profiles of nitrous oxide and oxygen. A water volume of 1 L was filtered through 0.2 μm polycarbonate filter, which were immediately stored at -80°C .

2.4 DNA-extraction

Until DNA-extraction filters were stored at -80°C , not exceeding five months after sampling. For extractions of total DNA one quarter of the filter was used. The extraction protocol followed standard methods [Ward *et al.*, 1993] using a QIAamp DNA Mini Kit (QIAGEN, Hilden, Germany). Extracted nucleic acids were stored in DNA-free water at -20°C .

2.5 PCR

16S rRNA genes were amplified using PCR by two different protocols for DGGE [Muyzer *et al.*, 1993] and cloning, with primer pairs specific for *Eubacteria* (see Tab. 1 and Tab. 2). The PCR for DGGE was performed as a touchdown-PCR; a GC-clamp was attached to the used primer.

For these reactions 1 μL of each primer (10 pmol L^{-1} , see table 1) and 1 μL DNA-extract were used, and PCR was performed using Ready-To-GoTM PCR Beads (Amersham Pharmacia Biotech). The amplification results were checked by agarose gels (1% agarose, 1 x TBE). The PCR products were purified with a High Pure PCR purification Kit (Roche, Mannheim, Germany), and used directly for following DGGE or cloning experiments.

Table 1: PCR protocol for DGGE, * temperature was reduced for 1 $^{\circ}\text{C}$ at each cycle

cycles	annealing	elongation		denaturation	
1				94 $^{\circ}\text{C}$	120 s
15	*65 – 50 $^{\circ}\text{C}$	40 s	72 $^{\circ}\text{C}$	40 s	94 $^{\circ}\text{C}$ 30 s
40	50 $^{\circ}\text{C}$	40 s	72 $^{\circ}\text{C}$	40 s	94 $^{\circ}\text{C}$ 30 s
1	42 $^{\circ}\text{C}$	60 s	72 $^{\circ}\text{C}$	300 s	
99	4 $^{\circ}\text{C}$				
primer: 5'342 GC ([CGC CCG CCG CGC GCG GCG GGC GGG GCG GGG GCA CGG GGG GC] CTA CGG GAG GCA GCA G)					
3'534 (ATT ACC GCG GCT GCT GG)					

Table 2: PCR protocol for cloning

cycles	annealing		elongation		denaturation	
1					94 °C	120 s
30	50 °C	40 s	72 °C	40 s	94 °C	30 s
1	42 °C	60 s	72 °C	300 s		
99	4 °C					

primer: 5'27F (GAG TTT GAT CCT GGC TCA G)
3'1387 (CGG GCG GTG TGT ACA AGG)

2.6 DG-DGGE

Bacterial diversity was qualitatively investigated by a double gradient modification DGGE (DG-DGGE) [Petri and Imhoff, 2001] at 8 stations (see Fig.1, stations marked with numbers). DG-DGGE was performed using a CBS 2001 Scientific System (Del Mar, Calif.). The gels consisted of a denaturant gradient (50-80%), superimposed with an acrylamide gradient (6 – 8 %). For electrophoresis of 25 μ L PCR-product a 0.5 TAE buffer was used at a constant voltage of 80 V and 60 °C for 14 h. Afterwards gels were stained with an ethidium bromide solution (0.5 mg L⁻¹) for 30 min, washed with MilliQ water for 30 min, and documented by a Kodak 1D Image analysis system .

2.7 Cloning and sequencing

For cloning and sequencing experiments two representative stations with four, respectively three depths were selected: station 175 in the western basin of the subtropical North Atlantic (31.747 °N, -50.752 °E) and station 194 in the eastern basin (35.525 °N, -23.982 °E). The depths were selected by measured N₂O profiles. For both stations the depth of the N₂O maximum was chosen, additionally depths above and below. Cloning experiments were performed with a TOPO TA Cloning[®] Kit for Sequencing (Invitrogen, Carlsbad, CA, USA), according to the manufacturer's manual. Clones were cultivated (for 24 h at 37 °C) on a LB medium containing 50 μ g mL⁻¹ canamycin, and afterwards 96 colonies of each cloning experiment were picked for sequencing. Sequencing was performed commercially (AGOWA, Berlin).

2.8 Sequence analysis and phylogeny

Sequences were manually refined and edited with the program Seqman (DNASTar, Laser-gene, Madison, USA), and compared to the Ribosomal Database Project, release 0.9, (<http://rdp.cme.msu.edu/classifier/classifier.jsp>), on the basis of the Bergey's Manual of Systematic Bacteriology, with a confidence threshold of 80 %. Phylogenetic analyses were made with the ARB software [Strunk *et al.*, 1999].

3. Results

3.1 Physical and chemical parameters

The physical and chemical parameters (at the representative stations 175 in the western basin and station 194 in the eastern basin) at depths sampled for sequence analyses are shown in Table 1.

Table 3: Physical and chemical conditions

	station 175				station 194		
	400 m	800 m	1200 m	4000 m	400 m	1000 m	1500 m
temperature (CTD)	16.63	9.28	5.88	2.31	13.71	9.07	5.76
salinity (CTD)	36.33	35.29	35.20	34.95	35.85	35.59	35.25
oxygen [$\mu\text{mol L}^{-1}$]	199	160	225	263	213	188	239
oxygen sat. [%]	70.3	55.7	72.5	77.1	82.0	65.4	76.7
nitrate [$\mu\text{mol L}^{-1}$]	7.36	22.2	19.8	20.3	9.8	18.9	18.2
nitrite [$\mu\text{mol L}^{-1}$]	0.01	0.0	0.0	0.0	0.0	0.0	0.0
phosphate [$\mu\text{mol L}^{-1}$]	0.41	1.4	1.3	1.4	0.7	1.2	1.2
silicate [$\mu\text{mol L}^{-1}$]	2.52	12.1	12.9	35.9	3.7	11.3	12.3
pot. density [kg m^{-3}]	26.6	27.3	27.7	27.9	26.9	27.6	27.8

N_2O distributions and profiles in the subtropical North Atlantic showed strong variations with water depth (Fig. 2). The profiles generally had one distinct maximum. In the surface layer concentrations were uniform, increasing below the thermocline up to a maximum and decreasing down to approximately 2000 m. Below 2000 m N_2O concentrations were nearly constant with depth in both basins.




Fig. 2: Profiles of a) N₂O and b) ΔN₂O at station 175 and 194; depth selected for sequencing are marked

Surface concentrations were $8.7 \pm 0.7 \text{ nmol L}^{-1}$, maximum values were found at depths between 600 to 1000 m; values ranged from 14.0 in the eastern basin (station 195) to 21.3 nmol L^{-1} in the western basin (station 156). Below 2000 m, concentrations were nearly constant at $13.1 \pm 0.9 \text{ nmol L}^{-1}$. Profiles in the western subtropical North Atlantic showed distinct maxima, while in the eastern basin no clear maximum was expressed. From the western to the eastern basin maximum concentrations decreased slightly from $17.7 \pm 1.4 \text{ nmol L}^{-1}$ to $15.1 \pm 0.7 \text{ nmol L}^{-1}$. East of the Midatlantic Ridge maxima were not clearly expressed and were broader. Additionally, maximum ΔN₂O values were lower in the eastern ($5.5 \pm 0.6 \text{ nmol L}^{-1}$) than in the western basin ($7.9 \pm 1.3 \text{ nmol L}^{-1}$).

3.2 DGGE

Figure 3 shows the gels of two DG-DGGE from station 175 in the western and station 194 in the eastern Basin of the subtropical North Atlantic (see Fig. 1). On both gels clearly different band patterns in different depths were found, although most bands were common to all samples. The band patterns of both stations differ in the composition of bands and the strength of those (at depths selected for sequencing some bands differing between the depths are exemplarily marked by red and blue circles). Therefore, a qualitatively different bacterial community in different depths of the water column can be assumed.




Fig. 3: DG-DGGE of eubacterial amplified 16S rDNA fragments covering a depth profile of the water column of station 175 (a) and station 194 (b); the left column shows an internal marker. The depth selected for cloning and sequencing experiments were underlined, differences between them were exemplarily marked with red and blue circles, note the position of marker and bands at each station.

3.3 Sequence analysis and phylogeny




Fig. 4: Total number of clones of bacterial groups, at station 175 and 194 at different depths, identified by the Ribosomal Database Project using a confidence threshold of 80 % <http://rdp.cme.msu.edu/index.jsp>. Lines between the bars separate *Proteobacteria* from other identified groups of bacteria, n gives the total number of sequenced clones

In total 640 clones were sequenced (mean length of sequences was 703 bases), and first analysis of sequences using the Ribosomal Database Project (Fig. 4) supported the results of the DGGE-gels that the bacterial communities differed with depths.

All clones of the clone libraries were strained into six major lineages of the domain bacteria: *Proteobacteria*, *Planctomycetes*, *Nitrospira*, *Actinobacteria*, *Cyanobacteria*, and *Bacteroidetes*. A large part of clones were most closely related to uncultured, environmental sequences. At station 175, the contribution of clones related to uncultured sequences decreased continuously with depth, whereas at station 194 those clones were mainly found at 1000 m.

Sequences affiliated to the *Proteobacteria* were the predominant group in both basins, followed by those related to uncultured bacteria. The composition and contribution of *Proteobacteria* was different depending on location and depth. In both basins sequences of the β - and δ -subdivision of *Proteobacteria* were hardly represented, whereas sequences related to α - and γ -*Proteobacteria* contribute to a major part to the total number of clones.

In the western basin the contribution of *Proteobacteria*-sequences increased significantly with depth (48.9 – 85.2 %), with changing composition: sequences related to α -

Proteobacteria were predominant at shallower depth, whereas sequences related to γ -*Proteobacteria* increased in proportion in deeper water masses. In contrast, at station 194 in the eastern basin sequences of α - and γ -*Proteobacteria* were also the predominant group, but showed no clear depth depending distribution. The number of clones affiliated to α -*Proteobacteria* was higher in the eastern basin than in the western, conversely to the number of clones affiliated to γ -*Proteobacteria*.

This is also obvious in the correlation of clone numbers of α - and γ -*Proteobacteria* with the potential density (Fig. 5). No correlations were found in the eastern basin at station 194. In contrast, in the western basin at station 175 the number of clones related to the α -*Proteobacteria* were found negatively correlated with the potential density, whereas those of γ -*Proteobacteria* were positively correlated.




Fig. 5: Correlation between the clone number of α - and γ -*Proteobacteria* (given as a percentage of the number of clones in total) and the potential density [kg m⁻³] at station 175 and station 194

Beside the increasing contribution of *Proteobacteria*-sequences and the differences between the contributions of sequences related to the α - and γ -*Proteobacteria* with depth, the composition of these subdivisions also changed depending on depth (see Fig. 6). This is obvious especially within the γ -subdivision.

At station 194, a depth depending contribution of the γ -subdivision of the *Proteobacteria* was not found, but the number of genera increased more than threefold (3 up to 10) from 400 m to 1500 m. In 1500 m particularly the clone numbers of members of the family *Alteromonadaceae* (*Alteromonas*, *Pseudoalteromonas*) increased.

At station 175 the total number of clones of the genera *Alteromonas* and *Acinetobacter* increased clearly with depth, at 4000 m a contribution of 68 % to the number of sequences affiliated to the *Proteobacteria* was found.




Fig. 6: Number of clones of the *Proteobacteria* groups, at different depths of station 175 and 194, n gives the number of identified clones by the Ribosomal data base using a confidence threshold of 80 % <http://rdp.cme.msu.edu/index.jsp>, lines between the bars separate the subdivisions of *Proteobacteria* from each other, the order of genera given in the legend correspond to the order of genera in the bars (top down)

A more detailed picture of phylogenetic relationships is given by analysis with the program ARB [Strunk *et al.*, 1999]. In general, the classification of sequences by ARB and RDP coincide. Most of the groups were found at both stations, with some exceptions. Seven sequences were affiliated with the group *Acidobacteria* (Fig. 10, highlighted in orange). These sequences were only found at station 194 in the eastern basin. Sequences affiliated to the groups *Piscirickettsiaceae*, *Spirochaetes*, *Verrucomicrobiae*, *Planctomycetes*, *Nitrospira*, and to one cluster related to uncultured sequences (Fig. 8, 10), were only found at station 175 in the western basin of the subtropical North Atlantic. In the α - and δ -subdivision of *Proteobacteria* (Fig. 7, 9) we found also cluster associated with only one basin.

A major number of sequences are affiliated to uncultured, environmental sequences, particularly in the α - and the δ -subdivision of the *Proteobacteria*. The closest relatives of most sequences related to the α -*Proteobacteria* (Fig. 7, group of uncultured bacteria with 151 of our sequences) are *Pelagibacter ubique* (1972; AF510191 and 1974; AF510192), a very small, heterotrophic bacterium, which is present throughout the oceans [Giovannoni *et al.*, 2005; Morris *et al.*, 2002]. A more detailed picture about the phylogenetic classification is given by trees exported from ARB (Fig. 7-10)




Fig. 7: Phylogenetic tree of 16S rDNA sequences affiliated to α -*Proteobacteria* exported from ARB (Maximum Parsimony); signatures of sequences obtained during this study start with the letters “sw” and “pl”; the number of sequences in each group is given in parentheses; groups of sequences highlighted in blue were found at station 175; uncoloured groups contain sequences of both stations




Fig. 8: Phylogenetic tree of 16S rDNA sequences affiliated to γ - and β -Proteobacteria exported from ARB (Maximum Parsimony); signatures of sequences obtained during this study start with the letters “sw” and “pl”; the number of sequences in each group is given in parentheses; groups of sequences highlighted in blue were found at station 175; uncoloured groups contain sequences of both stations




Fig. 9: Phylogenetic tree of 16S rDNA sequences affiliated to δ -Proteobacteria exported from ARB (Maximum Parsimony); signatures of sequences obtained during this study start with the letters “sw” and “pl”; groups of sequences highlighted in blue were found at station 175, groups of sequences highlighted in orange were found at station 194; uncoloured groups contain sequences of both stations




Fig. 10: Phylogenetic tree of 16S rDNA sequences affiliated to non-Proteobacteria exported from ARB (Maximum Parsimony); signatures of sequences obtained during this study start with the letters “sw” and “pl”; the number of sequences in each group is given in parentheses; groups of sequences highlighted in blue were found at station 175; groups of sequences highlighted in orange were found at station 194; uncoloured groups contain sequences of both stations

4. Discussion

16S rRNA genes have been used in several studies to analyze bacterial communities in the oceans [Bano and Hollibaugh, 2002; Braker et al., 2001; Corre et al., 2001; Fuhrman and Davis, 1997; Gallagher et al., 2004; Kato et al., 1996; López-García et al., 2003; Mullins et al., 1995; Schmidt et al., 1991; Ward and O'Mullan, 2002]. However, most work was done on specific groups of bacteria or restricted to single depths, mainly to the surface layer or the upper water column and the sediments. Only few studies provide data of the oceanic water column from the surface to the deep sea, using “universal” primers to investigate the bacteria community in total [Gallagher et al., 2004; Tholosan et al., 1999; Lee and Fuhrman, 1991; Maruyama et al., 1997]. The prokaryotes inhabiting the water column at depth greater than 1000 m are still largely unknown [Gallagher et al., 2004].

The structure and vertical distribution of bacterial communities presented here generally coincide with previous findings. Sequences were affiliated only with a few major lineages of the domain *Bacteria*, dominated by members of the group *Proteobacteria* [Bano and Hollibaugh, 2002; López-García et al., 2001; Morris et al., 2002]. The dominance of subdivisions of the *Proteobacteria* depended on depths: sequences related to the α -subdivision were more present in the surface layer [Bano and Hollibaugh, 2002], conversely to members of the γ -subdivision, which were predominant in the deep sea [López-García et al., 2001]. The vertical stratification leads to the interpretation that these groups are specialized to exploit vertical patterns in physical, chemical, and biological factors. The specialization to life at certain depths was shown in the ocean particularly for bacteria living at the surface and in the deep sea. Barotolerant and barophilic deep-sea bacterial strains belong to the genera of the γ -subdivision of *Proteobacteria*, namely *Shewanella*, *Photobacterium*, *Colwellia*, *Moritella*, and one new group [Kato et al., 1996; DeLong et al., 1997]. We found only small numbers of sequences belonging to these genera; the majority of sequences related to the γ -*Proteobacteria* in the deep sea belonged to the genera *Alteromonas* and *Acinetobacter*, with the closest affiliate to *Alteromonas macleodii* and *Acinetobacter junii*. In contrast, Maruyama et al. [1997] investigated the bacterial community at different depth in the Japan Trench, and found *Acinetobacter* as a main species in surface communities. However, DeLong et al. [1997] assume that the ability for growth under high pressure conditions might be fairly widely

distributed among the γ -*Proteobacteria*, and *Imada et al.* [2003] described barophilic strains of *Acinetobacter* in sediment samples collected from Sagami Bay and Enshu-Nada. Beside this, the concentration and composition of available nutrients differ clearly between surface and deep waters, whereas easily decomposable nutrients are depleted in deep waters [*Handa and Tominaga, 1969; Handa et al., 1992*]. The genera *Acinetobacter* and *Alteromonas* are known to be able to use an extremely wide variety of organic compounds [*Towner, 2001; Mikahailov et al., 2002*]. Therefore, the large number of sequences affiliated with *Acinetobacter* and *Alteromonas* might be not uncommon.

Differences between the stations might likely be explained by hydrographical aspects (Fig. 11) Phylogenetic analyses show that the majority of bacterial small subunit rRNA (SSU rRNA) gene lineages retrieved from near the coast of Oregon in the Pacific Ocean are related to those recovered from open ocean regions of both the Pacific and Atlantic Oceans, the Mediterranean Sea, and previously investigated coastal regions [*Rappé et al., 2000*]. With some exceptions, at both stations the same major lineages of Bacteria were found (Fig. 4, 6, 7-10), but with clearly different contributions of different groups to the bacterial community. Particularly at depths of 1200 m in the eastern and 1500 m in the western basin the chemical and physical conditions are not such different to explain the variety of bacterial community solely (Table 3, Fig. 11). Therefore, the distribution and contribution of water masses seem to have a strong influence on the bacterial community in the subtropical North Atlantic.




Fig. 11: T-S-diagram of the M 60-5 cruise, colour-divided into western and eastern basin, depths selected for sequencing of bacterial 16S rDNA clones are marked

Aim of this study was to investigate the composition of bacterial communities in the subtropical North Atlantic, with the intention to find correlations between the structure of bacterial communities and the distribution of the greenhouse gas N₂O. In the North Atlantic N₂O is assumed to be produced mainly by nitrification [see [Nitrous oxide in the North Atlantic Ocean, chapter 2 of this thesis](#)], therefore we expected sequences related to well-known aerobic and autotrophic nitrifying communities, such as *Nitrosomonas*, *Nitrosococcus*, *Nitrosospira*, and *Nitrobacter*, *Nitrospina*, *Nitrococcus* and *Nitrospira*. Surprisingly, only a few of our sequences were affiliated to these genera, particularly to the genera *Nitrospina* and *Nitrospira*. This might be a methodical artefact, i.e. caused by PCR bias or due to different rRNA gene numbers [v. [Wintzingerode et al., 1997](#)]. [Krawiec and Riley \[1990\]](#) showed a trend between fewer copies of rRNA gene regions (*rrn* operons) and slower growth of organisms, which is known for nitrifiers [[Ward, 2005](#)]. Both would lead to an underestimation of the contribution of autotrophic nitrifiers, respectively an overestimation of bacteria, whose sequences amplify with more efficiency to the primers or reveal more copies of the rRNA gene. However, autotrophic nitrifiers are very slow growing organisms with generation times on the order of a day, and they occur in relatively low abundances in natural environments [[Ward, 2005](#)]. So, despite methodical artefacts, there is also the possibility that the small number of sequences affiliated with nitrifiers might approximately mirror the contribution of autotrophic nitrifiers to the bacterial community.

Beside autotrophic nitrification, heterotrophic nitrification is also an opportunity for N₂O production. Commonly the relevance of heterotrophic nitrifiers within the nitrogen cycle was considered to be low, due to their lower nitrification rates in comparison to autotrophic nitrifiers [[Knowles, 1985](#); [Jetten et al., 1997](#)]. However, in recent years increased attention has been paid to heterotrophic nitrifiers because many of them are found to denitrify their nitrification products simultaneously [[Robertson et al., 1988](#); [Nishio et al., 1998](#)]. This might lead to a significant underestimation of their nitrification ability and the release of N₂O by heterotrophic nitrifiers. [Papen et al. \[1989\]](#) found heterotrophic nitrifiers producing much more N₂O per cell than autotrophic nitrifiers. Thus, since heterotrophic nitrifiers are more widespread and include a variety of species [[Mével and Prieur, 2000](#)], they may contribute significantly to the global N₂O budget [[Jetten et al., 1997](#)]. Heterotrophic nitrifiers which are able to produce N₂O are reported i.e. for strains of the γ -proteobacterial genus *Stenotrophomonas* [[Finkmann et al., 2000](#)]. Due to the

relation of a major part of sequences to uncultured clones, we have little or no information on the physiology of these groups. Additionally, suggestions of shared metabolic characteristics between neighbouring groups are sometimes dangerous due to the immense diversity and capability of bacteria for horizontal gene transfer and rapid mutation [Ward, 2005].

Further investigations are necessary, to obtain a more detailed picture about the activities of bacterial communities. Fluorescence-in-situ-hybridization experiments might be helpful to prove the activity of involved genes, or antibody-based proof of enzymes and measurements of their activity. Beside the focus on members of the domain *Bacteria*, also *Archaea* should be taken into account. *Archaea* are also widely distributed in the oceans [DeLong, 1992; Fuhrman and Davis, 1997], and also able to perform different reductive pathways of the N-cycle, including both assimilatory processes, such as nitrate assimilation and N₂ fixation, and dissimilatory reactions, such as nitrate respiration and denitrification [Cabello *et al.*, 2004]. Archaeal denitrifiers generally stop denitrification activities at the production of N₂O [Ward, 2005], therefore *Archaea* might contribute, maybe even under oxic conditions, to the observed distribution of N₂O.

5. Summary

- ④ The structure of the bacterial communities in the North Atlantic Ocean clearly depends on depths and sampling sites.
- ④ *Proteobacteria* were predominant in the western as well as in the eastern basin, but with clearly different contribution of subdivisions and genera.
- ④ Due to limited data an obvious relationship between N₂O distribution and single bacteria groups could not be proven.
- ④ Heterotrophic nitrifiers or other N₂O producer might be important due to the low number of sequences affiliated with known autotrophic nitrifiers.

Acknowledgements

We thank the officers and crew of R/V Meteor for their excellent support. Thanks to R. Hoffmann (Max Planck Institute for Chemistry, Mainz, Germany) for calibration of our standard gas mixtures. Special thanks to Tim Staufenger for experimental help, and to Vera Thiel and Martina Blümel for their essential mental support. The investigations were financially supported by the Deutsche Forschungsgemeinschaft through grant WA 1434/6-1 and BA 1990/3-1.

References

- Abeliovich, A., The nitrite oxidizing bacteria, in *The Prokaryotes: An evolving electronic resource for the microbiological community*, Springer Verlag, New York, 2001.
<http://link.springer-ny.com/link/service/books/10125/>
- Abou Seada, M.N.I., and J.C.G. Ottow, Effect of increasing oxygen concentration on total denitrification and nitrous oxide release from soil by different bacteria, *Biology and Fertility of Soils*, 1(1), 31-38, 1985.
- Bano, N., and J.T. Hollibaugh, Phylogenetic composition of bacterioplankton assemblages from the Arctic Ocean, *Applied and Environmental Microbiology*, 68 (2), 505-518, 2002.
- Bartosch, S., Wolgast, I., E. Spieck, and E. Bock, Identification of nitrite-oxidizing bacteria with monoclonal antibodies recognizing the nitrite oxidoreductase, *Applied and Environmental Microbiology*, 65, 4126-4133, 1999.
- Binder, B.J., S.W. Chisholm, R.J. Olson, S.L. Frankel, and A.Z. Worden, Dynamics of picophytoplankton, ultraphytoplankton and bacteria in the central equatorial Pacific, *Deep-Sea Research Part II - Topical Studies in Oceanography*, 43(4-6), 907-931, 1996.
- Bock, E., H.-P. Koops, and H. Harms, Cell biology of nitrifying bacteria, in: *Prosser, J.I. (Ed.), Nitrification, Special Publications of the Society for General Microbiology*, 20, 17-38, 1986.
- Bollmann, A., and H.J. Laanbroek, Influence of oxygen partial pressure and salinity on the community composition of ammonia-oxidizing bacteria in the Schelde estuary, *Aquatic Microbial Ecology*, 28(3), 239-247, 2002.

- Bonin, P., C. Tamburini, and V. Michotey, Determination of the bacterial processes which are sources of nitrous oxide production in marine samples, *Water Research*, 36(3), 722-732, 2002.
- Bothe, H., G. Jost, M. Schloter, B.B. Ward, and K.P. Witzel, Molecular analysis of ammonia oxidation and denitrification in natural environments, *FEMS Microbiology Reviews*, 24(5), 673-690, 2000.
- Braker, G., H.L. Ayala-del-Rio, A.H. Devol, A. Fesefeldt, and J.M. Tiedje, Community structure of denitrifiers, *Bacteria*, and *Archaea* along redox gradients in pacific north-west marine sediments by terminal restriction fragment length polymorphism analysis of amplified nitrite reductase (*nirS*) and 16S rRNA genes, *Applied and Environmental Microbiology*, 67(4), 1893-1901, 2001.
- Brettar, I., and G. Rheinheimer, Denitrification in the central Baltic: evidence for hydrogen sulfide oxidation as motor of denitrification at the oxic-anoxic interface, *Marine Ecology Progress Series* 77(2-3), 157-169, 1991.
- Buchanan, R.E., Studies on the nomenclature and classification of the *Bacteria* III. The families of the *Eubacteriales*, *Journal of Bacteriology*, 2, 347-350, 1917.
- Cabello, P., M.D. Roldan, and C. Moreno-Vivián, Nitrate reduction and the nitrogen cycle in *Archaea*, *Society for General Microbiology*, 150, 3527-3546, 2004.
- Carlson, C.A., H.W. Ducklow, and T.D. Sleeter, Stocks and dynamics of bacterioplankton in the northwestern Sargasso Sea, *Deep-Sea Research II*, 43, 491-515, 1996.
- Chen, M.Y., S.S. Tsay, K.Y. Chen, Y.C. Shi, Y.T. Lin, and G.H. Lin, *Pseudoxanthomonas taiwanensis* sp nov., a novel thermophilic, N₂O producing species isolated from hot springs, *International Journal of Systematic and Evolutionary Microbiology*, 52, 2155-2161, 2002.
- Cho, B.C., and F. Azam, Biogeochemical significance of bacterial biomass in the oceans euphotic zone, *Marine Ecology Progress Series*, 63, 253-259, 1990.
- Codispoti, L.A., J.A. Brandes, J.P. Christensen, A.H. Devol, S.W.A. Naqvi, H.W. Paerl, and T. Yoshinari, The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the anthropocene?, *Scientia Marina*, 65, 85-105, 2001.
- Corre, E., A.L. Reysenbach, and D. Prieur, ϵ -proteobacterial diversity from a deep-sea hydrothermal vent on the Mid-Atlantic Ridge, *FEMS Microbiology Letters*, 205(2), 329-335, 2001.
- DeLong, E.F., Archaea in coastal marine bacterioplankton, Proceedings of the National Academy of Science of the United States of America, 89, 5685-5689, 1992.
- DeLong, E.F., D.G. Franks, and A.A. Yayanos, Evolutionary relationship of cultivated psychrophilic and barophilic deep-sea bacteria, *Applied and Environmental Microbiology*, 63(5), 2105-2108, 1997.

- Field, K.G., D. Gordon, T. Wright, M. Rappé, E. Urbach, K. Vergin, and S.J. Giovannoni, Diversity and depth-specific distribution of SAR11 cluster rRNA genes from marine planktonic bacteria, *Applied and Environmental Microbiology*, 63(1), 63-70, 1997.
- Finkmann, W., K. Altendorf, E. Stackebrandt, and A. Lipski, Characterization of N₂O-producing *Xanthomonas*-like isolates from biofilters as *Stenotrophomonas nitritireducens* sp nov., *Luteimonas mephitis* gen. nov., sp nov and *Pseudoxanthomonas broegbernensis* gen. nov., sp nov, *International Journal of Systematic and Evolutionary Microbiology*, 50, 273-282, 2000.
- Focht, D.D., and W. Verstraete, Biochemical ecology of nitrification and denitrification, in *Advances in microbial ecology*, edited by M. Alexander, pp. 135–214, Plenum Press, New York, 1977.
- Fuhrman, J.A., and A.A. Davis, Widespread *Archaea* and novel bacteria from the deep sea as shown by 16S rRNA gene sequences, *Marine Ecology Progress Series*, 150(1-3), 275-285, 1997.
- Gallagher, J.M., M.W. Carton, D.F. Eardly, and J.W. Patching, Spatio-temporal variability and diversity of water column prokaryotic communities in the eastern North Atlantic, *FEMS Microbiology Ecology*, 47(2), 249-262, 2004.
- Giovannoni, S.J., H.J. Tripp, S. Givan, M. Podar, K.L. Vergin, D. Baptista, L. Bibbs, J. Eads, T.H. Richardson, M. Noordewier, M.S. Rappé, J.M. Short, J.C. Carrington, and E.J. Mathur, Genome streamlining in a cosmopolitan oceanic bacterium, *Science*, 309, 1242-1245, 2005.
- González, J.M., R. Simó, R. Massana, J.S. Covert, E.O. Casamayor, C. Pedrós-Alió, and M.A. Moran, Bacterial community structure associated with a dimethylsulfoniopropionate-producing North Atlantic algal bloom, *Applied and Environmental Microbiology*, 66(10), 4237-4246, 2000.
- Goreau, T.J., W.A. Kaplan, S.C. Wofsy, M.B. McElroy, F.W. Valois, and S.W. Watson, Production of nitrite and nitrous oxide by nitrifying bacteria at reduced concentrations of oxygen, *Applied and Environmental Microbiology* 40(3), 526-532, 1980.
- Handa, N., T. Nakatsuka, M. Fukuchi, H. Hattori, and T. Hoshiai, Vertical fluxes and ecological significance of organic materials during the phytoplankton bloom during austral summer in Breid Bay, Antarctica, *Marine Biology*, 112(3), 469-478, 1992.
- Handa, N., and H. Tominaga, A detailed analysis of carbohydrates in marine particulate matter, *Marine Biology*, 2, 228-235, 1969.
- Imada, C., T. Kobayashi, N. Hamada-Sato, and E. Watanabe, Isolation and characterization of barophilic bacteria obtained from sediment samples collected from Sagami Bay and Enshu-Nada, *Nippon Suisan Gakkaishi*, 69(3), 347-351, 2003.

- Jetten, M.S.M., S. Logemann, G. Muyzer, L.A. Robertson, S. deVries, M.C.M. van Loosdrecht, and J.G. Kuenen, Novel principles in the microbial conversion of nitrogen compounds, *Antonie van Leeuwenhoek International Journal of General and Molecular Microbiology*, 71(1-2), 75-93, 1997.
- Kato, C., A. Inoue, and K. Horikoshi, Isolating and characterizing deep-sea microorganisms, *Trends in Biotechnology*, 14, 6-12, 1996.
- Knowles, R., Microbial transformations as sources and sinks for nitrogen oxides, in J. A. Brierley (ed.), *Plantary ecology*, Van Nostrand Reinhold Co., Inc., New York, 411-426, 1985.
- Krawiec, S., and M. Riley, Organization of the bacterial chromosome, *Microbiological Reviews*, 54 (4), 502-539, 1990.
- Krümme, A., and H. Harms, Effect of organic matter on growth and cell yield of ammonia-oxidizing bacteria, *Archives of Microbiology*, 133, 50-54, 1982.
- Lee, S., and J.A. Fuhrman, Spatial and temporal variation of natural bacterioplankton assemblages studied by total genomic DNA cross-hybridization, *Limnology and Oceanography*, 36, 1277-1287, 1991.
- López-García, P., S. Duperron, P. Philippot, J. Foriel, J. Susini, and D. Moreira, Bacterial diversity in hydrothermal sediment and ϵ -proteobacterial dominance in experimental microcolonizers at the Mid-Atlantic Ridge, *Environmental Microbiology*, 5(10), 961-976, 2003.
- López-García, P., A. López-López, D. Moreira, and F. Rodríguez-Valera, Diversity of free-living prokaryotes from a deep-sea site at the Antarctic Polar Front, *FEMS Microbiology Ecology*, 36(2-3), 193-202, 2001.
- Maruyama, A., R. Taniguchi, H. Tanaka, and T. Higashihara, Low-temperature adaptation of deep-sea bacteria isolated from the Japan Trench, *Marine Biology*, 128(4), 705-711, 1997.
- Mével, G., and D. Prieur, Heterotrophic nitrification by a thermophilic *Bacillus* species as influenced by different culture conditions, *Canadian Journal of Microbiology*, 46, 465-473, 2000.
- Mikhailov, V.V., L.A. Romanenko, and E.P. Ivanova, The genus *Alteromonas* and related *Proteobacteria*, in *The Prokaryotes: An Evolving Electronic Resource for the Microbiological Community*, Springer Verlag, New York, 2002.
<http://link.springer-ny.com/link/service/books/10125/>
- Moir, J.W.B., D.J. Richardson, and S.J. Ferguson, The expression of redox proteins of denitrification in *Thiosphaera pantotropha* grown with oxygen, nitrate, and nitrous oxide as electron acceptors, *Archives of Microbiology*, 164(1), 43-49, 1995.

- Moreno-Vivián, C., P. Cabello, M. Martínez-Luque, R. Blasco, and F. Castillo, Prokaryotic nitrate reduction: Molecular properties and functional distinction among bacterial nitrate reductases, *Journal of Bacteriology*, 181(21), 6573-6584, 1999.
- Morris, R.M., M.S. Rappé, S.A. Connon, K.L. Vergin, W.A. Siebold, C.A. Carlson, and S.J. Giovannoni, SAR11 clade dominates ocean surface bacterioplankton communities, *Nature*, 420(19), 806-810, 2002.
- Mullins, T.D., T.B. Britschgi, R.L. Krest, and S.J. Giovannoni, Genetic comparisons reveal the same unknown bacterial lineages in Atlantic and Pacific bacterioplankton communities, *Limnology and Oceanography*, 40(1), 148-158, 1995.
- Muyzer, G., E.C. De Waal, and A.G. Uitterlinden, Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA, *Applied and Environmental Microbiology*, 59, 695-700, 1993.
- Naqvi, S.W.A., D.A. Jayakumar, P.V. Narvekar, H. Naik, V. Sarma, W. D'Souza, S. Joseph, and M.D. George, Increased marine production of N₂O due to intensifying anoxia on the Indian continental shelf, *Nature*, 408(6810), 346-349, 2000.
- Nishio, T., T. Yoshikura, H. Mishima, Z. Inouye, and H. Itoh, Conditions for nitrification and denitrification by an immobilized heterotrophic nitrifying bacterium *Alcaligenes faecalis* OKK17, *Journal of Fermentation and Bioengineering*, 86(4), 351-356, 1998.
- Ostrom, N.E., M.E. Russ, B. Popp, T.M. Rust, and D.M. Karl, Mechanisms of N₂O production in the subtropical North Pacific based on determinations of the isotopic abundances of N₂O and N₂, *Chemosphere: Global Change Science*, 2(3-4), 281-290, 2000.
- Papen, H., R. Vonberg, I. Hinkel, B. Thoene, and H. Rennenberg, Heterotrophic nitrification by *Alcaligenes faecalis* – NO₂⁻, NO₃⁻, N₂O, and NO production in exponentially growing cultures, *Applied and Environmental Microbiology*, 55(8), 2068-2072, 1989.
- Petri, R., and J.F. Imhoff, The relationship of nitrate reducing bacteria on the basis of *narH* gene sequences and comparison of *narH* and 16S rDNA based phylogeny, *Systematic and Applied Microbiology*, 23, 47-57, 2000.
- Petri, R., and J.F. Imhoff, Genetic analysis of sea-ice bacterial communities of the western Baltic Sea using an improved double gradient method, *Polar Biology*, 24(4), 252-257, 2001.
- Popp, B.N., M.B. Westley, S. Toyoda, T. Miwa, J.E. Dore, N. Yoshida, T.M. Rust, F.J. Sansone, M.E. Russ, N.E. Ostrom, and P.H. Ostrom, Nitrogen and oxygen isotopic constraints on the origins and sea-to-air flux of N₂O in the oligotrophic subtropical North Pacific gyre, *Global Biogeochemical Cycles*, 16(4), doi: 10.1029/2001GB001806, 2002.

- Rappé, M. S., K. Vergin, and S.J. Giovannoni, Phylogenetic comparisons of a coastal bacterioplankton community with its counterparts in open ocean and freshwater systems, *FEMS Microbiology Ecology*, 33(3), 219-232, 2000.
- Richardson, D.J., and N.J. Watmough, Inorganic nitrogen metabolism in bacteria, *Current Opinion in Chemical Biology*, 3, 207-219, 1999.
- Richardson, D.J., Bacterial respiration: a flexible process for a changing environment, *Microbiology*, 146, 551-571, 2000.
- Robertson, L.A., and J.G. Kuenen, Heterotrophic nitrification in *Thiosphaera pantotropha* - oxygen uptake and enzyme studies, *Journal of General Microbiology*, 134, 857-863, 1988.
- Robertson, L.A., E.W.J. Vanniell, R.A.M. Torremans, and J.G. Kuenen, Simultaneous nitrification and denitrification in aerobic chemostat cultures of *Thiosphaera pantotropha*, *Applied and Environmental Microbiology*, 54 (11), 2812-2818, 1988.
- Schmidt, T., E.F. DeLong, and N.R. Pace, Analysis of a marine picoplankton community by 16S rRNA gene cloning and sequencing, *Journal of Bacteriology*, 173(14), 4371-4378, 1991.
- Schramm, A., D. De Beer, M. Wagner, and R. Amann, Identification and activities in situ of *Nitrosospira* and *Nitrospira* spp. as dominant populations in a nitrifying fluidized bed reactor, *Applied and Environmental Microbiology*, 64, 3480-3485, 1998.
- Shapleigh, J.P., The denitrifying prokaryotes, in *The Prokaryotes: An Evolving Electronic Resource for the Microbiological Community*, 3rd edition, release 3.1, Springer Verlag, New York, 2001.
<http://141.150.157.117:8080/prokPUB/index.htm>
- Strunk, O., O. Gross, B. Reichel, M. May, S. Hermann, N. Stuckman, B. Nonhoff, M. Lemke, A. Ginhart, A. Vilbig, T. Ludwig, K.-H. Schleifer, and W. Ludwig, ARB: a software environment for sequence data, in <http://www.mikro.biologie.tu-muenchen.de>, Department of Microbiology, Technische Universität, München, 1999.
- Tholosan, O., J. Garcin, and A.M. Bianchi, Effects of hydrostatic pressure on microbial activity through a 2000 m deep water column in the NW Mediterranean Sea, *Marine Ecology Progress Series*, 183, 49-57, 1999.
- Towner, K., The Genus *Acinetobacter*, in *The Prokaryotes: An Evolving Electronic Resource for the Microbiological Community*, Springer Verlag, New York, 2001.
<http://link.springer-ny.com/link/service/books/10125/>
- v. Wintzingerode, F., U.B. Göbel, and E. Stackebrandt, Determination of microbial diversity in environmental samples: pitfalls of PCR-based rRNA analysis, *FEMS Microbiology Reviews*, 21, 213-229, 1997.

- Venter, C.J., K. Remington, J.F. Heidelberg, A.L. Halpern, D. Rusch, J.A. Eisen, D. Wu, I. Paulsen, K.E. Nelson, W. Nelson, D.E. Fourts, S. Levy, A.H. Knap, M.W. Lomas, K. Nealson, O. White, J. Peterson, J. Hoffman, R. Parsons, H. Baden-Tillson, C. Pfannkoch, Y-H. Rogers, and H.O. Smith, Environmental genome shotgun sequencing of the Sargasso Sea, *Science*, 304, 66-74, 2004.
- Vollack, K.U., and W.G. Zumft, Nitric oxide signaling and transcriptional control of denitrification genes in *Pseudomonas stutzeri*, *Journal of Bacteriology*, 183(8), 2516-2526, 2001.
- Walter, S., I. Peeken, K. Lochte, and H.W. Bange, Nitrous oxide measurements during EIFEX, the European Iron Fertilisation Experiment in the subpolar South Atlantic Ocean, *Geophysical Research Letters* (in press), 2005.
- Ward, B.B., Molecular approaches to marine microbial ecology and the marine nitrogen cycle, *Annual Review of Earth and Planetary Sciences*, 33, 301-333, 2005.
- Ward, B.B., A.R. Cockcroft, and K.A. Kilpatrick, Antibody and DNA probes for detection of nitrite reductase in seawater, *Journal of General Microbiology*, 139, 2285-2293, 1993.
- Ward, B.B., and G.D. O'Mullan, Worldwide distribution of *Nitrosococcus oceani*, a marine ammonia-oxidizing γ -*Proteobacterium*, detected by PCR and sequencing of 16S rRNA and *amoA* genes, *Applied and Environmental Microbiology*, 68(8), 4153-4157, 2002.
- Wetzel, R.G., Limnology, *Saunders College Publishing, Philadelphia, Pa.*, 1983.
- Whittaker, M., D. Bergmann, D. Arciero, and A.B. Hooper, Electron transfer during the oxidation of ammonia by the chemolithotrophic bacterium *Nitrosomonas europaea*, *Biochimica et Biophysica Acta*, 1459, 346-355, 2000.
- Woese, C.R., E. Stackebrandt, W.G. Weisburg, B.J. Paster, M.T. Madigan, V.J. Fowler, C.M. Hahn, P. Blanz, R. Gupta, Nealson, and G.E. Fox, The phylogeny of purple bacteria: the α -subdivision, *Systematic and Applied Microbiology*, 5, 315-326, 1984a.
- Woese, C.R., J.B. Weisburg, B.J. Paster, C.M. Hahn, R.S. Tanner, N.R. Krieg, H.-P. Koops, H. Harms, and E. Stackebrandt, The phylogeny of purple bacteria: the β -subdivision, *Systematic and Applied Microbiology*, 5, 327-336, 1984b.
- Woese, C.R., W.G. Weisburg, C.M. Hahn, B.J. Paster, L.B. Zablen, B.J. Lewis, T.J. Macke, W. Ludwig, and E. Stackbrandt, The phylogeny of purple bacteria: the γ -subdivision, *Systematic and Applied Microbiology*, 6, 25-33, 1985.
- Wrage, N., G.L. Velthof, M.L. van Beusichem, and O. Oenema, Role of nitrifier denitrification in the production of nitrous oxide, *Soil Biology and Biochemistry*, 33(12-13), 1723-1732, 2001.

- Yamagishi, H., N. Yoshida, S. Toyoda, B.N. Popp, M.B. Westley, and S. Watanabe, Contributions of denitrification and mixing on the distribution of nitrous oxide in the North Pacific, *Geophysical Research Letters*, 32(4), 2005.
- Zart, D., I. Schmidt, and E. Bock, Significance of gaseous NO for ammonia oxidation by *Nitrosomonas eutropha*, *Antonie van Leeuwenhoek International Journal of General and Molecular Microbiology*, 77, 49-55, 2000.
- Zehr, J.P., and B.B. Ward, Nitrogen cycling in the ocean: New perspectives on processes and paradigms, *Applied and environmental microbiology*, 68(3), 1015-1024, 2002.
- Zubkov, M.V., B.M. Fuchs, S.D. Archer, R.P. Kiene, R. Amann, and P.H. Burkill, Linking the composition of bacterioplankton to rapid turnover of dissolved dimethylsulphoniopropionate in an algal bloom in the North Sea, *Environmental Microbiology*, 3(5), 304-311, 2001.

Chapter 7

Influence of HgCl₂-poisoning and temperature on N₂O-concentrations during storage

Sylvia Walter, Hermann W. Bange

Abstract

Sometimes it is necessary to store samples until measurements. Commonly samples for N₂O determination are poisoned with saturated aqueous HgCl₂ solution to prevent biological activity until measurements. To investigate the effect of storage on N₂O concentrations at different conditions, we started an experiment over more than ten months. Samples were poisoned in three different ways, and stored at two different temperatures. With one exception, no significant change in N₂O concentrations was found after 10 months either in different poisoned treatments or in treatments stored at different temperatures. Only not poisoned samples, stored at room temperature, showed a distinct decrease in N₂O concentrations after 33 days to a concentration of approximately zero. Thus, it is possible to store samples for N₂O measurements for at least ten months after sampling.

1. Introduction

Due to temporal and spatial limitations during research cruises, it is not always possible to measure the obtained samples onboard. In this case samples need to be stored until the measurements can be conducted. However, in order to use stored samples it is necessary to know, how long samples can be stored without significant changes of parameters of interest e.g. dissolved N_2O . Commonly, to prevent biological activity until measurements, aqueous samples are poisoned with a saturated aqueous HgCl_2 solution, which acts as a strong cell and protoplasm toxin [Glud *et al.*, 1995; Hemond and Duran, 1989; Jensen *et al.*, 1984; Kaplan *et al.*, 1978; McElroy *et al.*, 1978; Naqvi *et al.*, 1994; Oudot *et al.*, 2002]. However, the use of saturated solutions seems to be not necessary because non-saturated aqueous HgCl_2 solutions are already highly toxic for aquatic organisms. For example the EC_5 (effective concentration) for organisms such as *Pseudomonas putida* is 0.01 mg L^{-1} (16h) [Safety data sheets of HgCl_2 , <http://uk.chemdat.info/mda/uk/>, product number 104419]. Additionally low concentrated aqueous HgCl_2 solution may facilitate the handling of the solution during ship expeditions.

To investigate the effect of storage time on dissolved N_2O we collected surface water samples from the Kiel Fjord (Baltic Sea), and measured N_2O concentrations after different time periods during more than ten months. We applied different treatments of the samples regarding the degree of poisoning and storage temperatures.

2. Methods

We took surface water samples from the pier in front of the “Leibniz-Institut für Meereswissenschaften” (Kiel) at the 4th of September 2003. The analytical method applied is a modification of the method described by *Bange et al.* [2001]. 162 bubble free samples were taken, sealed directly with butyl rubber stoppers, and crimped with aluminium caps. Directly after sampling one third of the samples was treated with 300 μL per vial of a low concentrated mercury chloride (Merck, for analysis) solution of 40 mg L^{-1} , one third with 300 μL saturated mercury chloride solution (6 g L^{-1}) and the last third of the samples were not poisoned. At the first day ($t = 0$) we measured six samples of each treatment. Then the remaining samples were divided into storage at room temperature (in the lab, temperatures were between 17 – 25 $^{\circ}\text{C}$) and storage at 4 $^{\circ}\text{C}$ in the fridge. All samples were stored in a paper board; therefore they were not exposed to light. For each treatment (poison, temperature) we measured six samples.

N_2O water concentrations (C_w) were calculated as follows:

$$C_w \left[\text{nmol L}^{-1} \right] = \left(\beta \times P V_{wp} + \frac{x P}{R T} V_{hs} \right) / V_{wp}$$

where β stands for the Bunsen solubility in $\text{nmol L}^{-1} \text{atm}^{-1}$ [*Weiss and Price, 1980*], x is the dry gas mole fraction of N_2O in the headspace in ppb, P is the atmospheric pressure in atm (set to 1 atm), V_{wp} and V_{hs} stand for the volumes of the water (14 mL) and headspace phases (10 mL), respectively. R is the gas constant ($8.2054 \cdot 10^{-2} \text{ L atm mol}^{-1} \text{K}^{-1}$) and T is the temperature during equilibration.

For calibration we used standard gas mixtures with 311.8 ± 0.2 ppb and 346.5 ± 0.2 ppb N_2O in synthetic air (DEUSTE Steininger GmbH, Mühlhausen, Germany). The standard mixtures have been calibrated against the NOAA (National Oceanic and Atmospheric Administration, Boulder, Co.) standard scale in the laboratories of the Air Chemistry Division of the Max Planck Institute for Chemistry, Mainz, Germany). The overall mean analytical precision was $\pm 0.66 \%$.

The effectiveness of low concentrated HgCl_2 was additionally tested optically. 300 μL were added to a surface sea water probe with living flagellates, recognizable by agile movements.

3. Results and discussion

N₂O concentrations of different treatments are shown in Figure 1.




Fig. 1: N₂O concentration after different treatments, lines give an estimated error of ± 1 nmol L⁻¹, results of N₂O measurements of not poisoned samples stored at room temperature are red circled

N₂O concentrations measured directly after sampling ($t = 0$) showed a mean concentration of 9.97 ± 0.14 nmol L⁻¹, independent from poisoning. These concentrations correspond to the expected equilibrium concentration of 9.63 nmol L⁻¹ in surface waters. The equilibrium concentration was calculated with an atmospheric N₂O concentration of 318 ppb, and the ambient salinity and temperature. 10.68 ± 1.07

With one exception, after 10 months we found no significant change in N₂O concentrations either in different poisoned treatments or in treatments stored at different temperatures. The mean N₂O concentration of the treated samples was 10.68 ± 1.07 nmol L⁻¹ (samples measured after 20 days and not poisoned samples stored at room temperature were excluded). However, there might be a weak trend to increasing N₂O concentrations in poisoned samples with time; more distinct in cooled samples than in samples stored at room temperature (see Fig. 1).

Only not poisoned samples stored at room temperature showed a distinct decrease in N₂O concentrations after 33 days to approximately zero (see Fig. 1, red circles). The decrease in measured N₂O concentrations after 20 days was probably an artefact due to problems with the carrier gas flow. However, to demonstrate that the decrease of N₂O concentrations of not poisoned samples stored at room temperature does not occur before 20 days of storage these data were also shown.

[*Elkins, 1980*] published results of storage experiments with freshwater from the Boston's Muddy river, where 50 mL samples were poisoned with 200 µL saturated HgCl₂ and stored at 5 to 10 °C. He found no significant storage effect on poisoned samples after 45 days. This is in general agreement with our results. However, based on the fact that *Elkins [1980]* used a freshwater sample with a comparable high N₂O concentration of 591 nmol L⁻¹, these experimental conditions were not representative for oceanic N₂O concentrations.

Yoshinari [1976] tested the effect of HgCl₂ on water samples from the open Atlantic Ocean. 650 mL samples of different depths were either poisoned with 0.5 mL of 3% HgCl₂ solution or not poisoned. With the exception of water from the oxygen minimum layer he found no differences between both treatments. In the oxygen minimum layer N₂O concentrations in the not poisoned samples were two times higher than in those poisoned with HgCl₂. The effect of HgCl₂ was also tested by *Kieskamp et al. [1988]*. They found an increase of N₂O concentrations in sediment pore water after addition of HgCl₂; assuming a reduction of nitrite to N₂O by Fe²⁺. However, mechanisms of these reactions are not completely understood.

The distinct decrease of N₂O concentrations in not poisoned samples stored at room temperature indicates a conversion of N₂O by biological activities. Without poisoning a degradation of O₂ by ongoing biological activities can be assumed, which lead to a change to anoxic conditions in the vials. Under anoxic conditions N₂O can be used as an electron acceptor instead of O₂ [*Elkins et al., 1978; Cohen and Gordon, 1978*]. Therefore, the concentrations clearly decrease and remain near zero.

4. Conclusions

- ④ There was no difference between poisoned samples with low concentrated or saturated HgCl_2 solution. Thus, it is not necessary to work with saturated HgCl_2 solution.
- ④ There is no difference between poisoned samples stored at room temperature or at 4 °C, on condition that samples are poisoned. Thus, samples must not be transported with cooling.
- ④ If samples are measured closely within a few days after sampling, it is not necessary to poison them.

References

- Bange, H.W., S. Rapsomanikis, and M.O. Andreae, Nitrous oxide cycling in the Arabian Sea, *Journal of Geophysical Research-Oceans*, 106 (C1), 1053-1065, 2001.
- Cohen, Y., and L.I. Gordon, Nitrous oxide in the oxygen minimum of the eastern tropical North Pacific: Evidence for its consumption during denitrification and possible mechanisms for its production, *Deep-Sea Research*, 25 (6), 509-524, 1978.
- Elkins, J.W., S.C. Wofsy, M.B. McElroy, C.E. Kolb, and W.A. Kaplan, Aquatic sources and sinks for nitrous oxide, *Nature*, 275 (5681), 602-606, 1978.
- Elkins, J.W., Determination of dissolved nitrous oxide in aquatic systems by gas chromatography using electron capture detection and multi phase equilibration, *Analytical Chemistry*, 52, 263-267, 1980
- Glud, R. N., K. Jensen, and N.P. Revsbech, Diffusivity in Surficial Sediments and Benthic Mats Determined By Use of a Combined $\text{N}_2\text{O} - \text{O}_2$ Microsensor, *Geochimica Et Cosmochimica Acta*, 59(2), 231-237, 1995.
- Hemond, H.F. and A.P. Duran, Fluxes of N_2O at the sediment-water and water-atmosphere boundaries of a nitrogen-rich river, *Water Resources Research*, 25(5), 839-846, 1989.
- Jensen, H. B., K.S. Jørgensen, and J. Sørensen, Diurnal variation of nitrogen cycling in coastal, marine sediments II. Nitrous oxide emission, *Marine Biology*, 83, 177-183, 1984.
- Kaplan, W.A., J.W. Elkins, C.E. Kolb, M.B. McElroy, S.C. Wofsy, and A.P. Duran, Nitrous oxide in fresh water systems: An estimate for the yield of atmospheric N_2O associated with the disposal of human waste, *Pure and Applied Geophysics*, 116, 423-438, 1978.
- Kieskamp, W.M., K. Eijkman, and W. Helder, The effect of mercury(II)chloride on the determination of nitrous oxide profiles in marine sediments, *Netherlands Journal of Sea Research*, 22(3); 199-203; 1988.

McElroy, M. B., J.W. Elkins, S.C. Wofsy, C.E. Kolb, A.P. Duran, and W.A. Kaplan, Production and release of N₂O from the Potomac Estuary, *Limnology and Oceanography*, 23, 1168-1182, 1978.

Naqvi, S. W. A., D.A. Jayakumar, M. Nair, M. Dileep Kumar, and M.D. George, Nitrous oxide in the western Bay of Bengal, *Marine Chemistry*, 47, 269-278, 1994.

Oudot, C., P. Jean-Baptiste, E. Fourre, C. Mormiche, M. Guevel, J.-F. TERNON, and P. Le Corre, Transatlantic equatorial distribution of nitrous oxide and methane, *Deep-Sea Research Part I*, 49, 1175-1193, 2002.

Safety data sheets of HgCl₂, <http://uk.chemdat.info/mda/uk/>, product number 104419

Weiss, R.F., and B.A. Price, Nitrous oxide solubility in water and seawater, *Marine Chemistry*, 8, 347-359, 1980.

Yoshinari, T., Nitrous oxide in the sea, *Marine Chemistry*, 4, 189-202, 1976.

Acknowledgements

Zuallererst möchte ich mich ganz herzlich bei Herrn Dr. Bange bedanken. Seine ständige Diskussionsbereitschaft, seine Eigenschaft als wandelndes Literaturlexikon und sein unerschütterlicher Optimismus haben entscheidend zum Erfolg dieser Arbeit beigetragen.

Vielen Dank auch an Herrn Prof. Wallace und Frau Prof. Lochte, die diese Arbeit nicht nur ermöglichten, sondern beide auch eine stetige Quelle hilfreicher Kommentare und guter Ratschläge waren. Auch Herrn Prof. Imhoff vielen Dank für die Möglichkeit, in seiner Abteilung meine Kenntnisse zu vertiefen. In diesem Zusammenhang gilt mein Dank ebenfalls der DFG, welche meine Forscherambitionen finanziell unterstützt hat.

Ute Weidinger als Seele der Abteilung möchte ich für die vielen netten Stunden danken, für ihre ständige Hilfsbereitschaft und ihr Vertrauen, irgendwann in Rente gehen zu dürfen. Auch Peter Streu und Hergen Johanssen, Birgit Quack sowie Tobias Steinhoff vielen Dank für die nette Arbeitsatmosphäre.

In der vorliegenden Arbeit könntewerdewollte ich mich bei Martina Blümel bedanken, die mir immer zur Seite stand, mein ausschweifendes Gejammer ertragen hat, und deren Ohr zu jeder Tages- und Nachtzeit und in jedem Zustand offen war. Auch Isabel Schunck möchte ich für ihre Geduld, Ohrverleihung und Hilfsbereitschaft danken. Ich kann nur sagen: Mit Euch durch die Gegend zu paddeln ist (in Abwesenheit vom Westwind und seinen fiesigen Anverwandten) Entspannung pur!

Bedanken möchte ich mich auch bei Heiko Trautmann und Michael Schunck, welche mich wöchentlich badmintonisch mit Wonne gescheucht haben, und so die hemmungslose Aufnahme von jeder Menge kalorienhaltiger Nervennahrung ermöglichten. Dies gilt auch für Marcus Tank, der außerdem durch seine ARBsolute Ausdauer im Einladen von Sequenzen eine große Hilfe war.

Jörg Süling, Vera Thiel und Tim Staufenberger vielen Dank für die unermüdliche Beantwortung all meiner Fragen, und die anregenden Diskussionen über mögliche und unmögliche Antworten. Dank auch im Namen aller Denitrifikationskaninchen!

Der unermüdlichen Messwut von Imke Görtz und Ulrich Breitenbach ist es zu verdanken, dass meine Probensammelleidenschaft zu zahlreichen schönen Daten geführt hat. Außerdem hatten wir auch noch viel Spaß dabei!

Bei Sarah Gebhardt und Karsten Friis bedanke ich mich für eine klasse Büro-Gemeinschaft, viele lustige Stunden und musische Klassikqualen.

Herrn Dr. Nausch und seinen Kollegen vom IOW möchte ich ganz herzlich für die Möglichkeit danken, in der Ostsee Proben zu sammeln, und auch für die problemlose Bereitstellung von Daten.

Mein ganz besonderer Dank aber gilt Roland D.T. Mumm, der immer an meiner Seite war und mich gut genug kennt, um stets die richtigen Worte zu finden. Seine liebevolle Betreuung und ständige Boykottierung meiner hausfraulichen Ambitionen war der Schlüssel zum Ende.

Dieser besondere Dank gilt auch meiner Oma, meiner Schwester und meiner Mutter, die mich immer unterstützt, mir den Rücken freigehalten und an mich geglaubt haben. Ohne sie wäre ich heute nicht hier. Danke!

Abschließend möchte ich mich noch bei all denen bedanken, die ich hier namentlich nicht mehr aufgeführt habe. In der Zeit hier am Institut habe ich viele nette, aufgeschlossene und hilfsbereite Menschen kennen gelernt, die alle ihren Teil dazu beigetragen haben, dass ich mich gerne an diese Zeit zurückerinnern werde.