

**Potential of Phytonutrients to act as
Mimetics for Caloric Restriction –
Identification of the Transcriptional and
Functional Overlap in *silico***

Dissertation
zur Erlangung des Doktorgrades
der Agrar- und Ernährungswissenschaftlichen Fakultät
der Christian-Albrechts-Universität zu Kiel

vorgelegt von
M.Sc. Stefanie Rieper
Kiel, 2016

**Aus dem Institut für Humanernährung und Lebensmittelkunde
der Christian-Albrechts-Universität Kiel**

**Potential of Phytonutrients to act as Mimetics for Caloric Restriction –
Identification of the Transcriptional and Functional Overlap in *silico***

Dissertation
zur Erlangung des Doktorgrades
der Agrar- und Ernährungswissenschaftlichen Fakultät
der Christian-Albrechts-Universität zu Kiel

vorgelegt von

M.Sc. Stefanie Rieper
aus Neuenkirchen

Kiel, 2016

Dekan: Prof. Dr. Eberhard Hartung
1. Berichterstatter: Prof. Dr. Gerald Rimbach
2. Berichterstatter: Prof. Dr. Frank Döring
Tag der mündlichen Prüfung: 13.07.2016

Gedruckt mit Genehmigung der Agrar- und Ernährungswissenschaftlichen Fakultät der
Christian-Albrechts-Universität zu Kiel

Summary

Caloric restriction (CR) is known for its beneficial effects on life- and healthspan (lifespan without the occurrence of age-related diseases) in several model organisms. Studies in rhesus monkeys and humans are still ongoing, but the first results are promising. Nonetheless, it is assumed to be very difficult for humans to reach long-term CR because food is always available and because CR has many side effects, such as hunger and sensitivity to cold. Therefore, it is of interest to identify CR mimetics which are able to enhance health- and lifespan without affecting calorie intake. Some promising pharmaceutical substances have been identified (e.g., rapamycin and metformin) but most of them also show side effects. Therefore it will be of interest to identify food derived compounds such as plant bioactives (phytonutrients) as potential CR mimetics.

The aim of this study is to develop a new approach that will allow identifying potential CR mimetics *in silico*. Therefore, the transcriptional and functional overlap between CR and putative plant derived CR mimetics is analysed by using data of previously published investigations from the databases ArrayExpress and Gene Expression Omnibus (GEO).

In the present thesis nine CR experiments are identified, four experiments investigating the impact of different phytonutrients supplementing a normal diet (e.g., resveratrol, phloridzin and quercetin) and one experiment investigating the influence of phytonutrients administered intraperitoneally (quercetin, genistein). The results of these experiments were summarised with the R package RankProd to identify genes that were consistently differently regulated by CR, resveratrol, phloridzin, quercetin and genistein. These gene lists were compared to identify the transcriptional and functional overlap between CR and the putative mimetics.

Our results show that resveratrol had the largest overlap in gene expression regulation, so it is assumed that resveratrol can mimic many functions of CR. However, our results were unable to show that resveratrol increases the expression of sirtuin 1 (Sirt1; typically mentioned as a mediator of CR effects) or influences insulin-like growth factor-1 (Igf1) signalling. However, our results indicate that quercetin administered into a diet could be a CR mimetic, although only a smaller overlap in gene expression compared to resveratrol could be identified. Quercetin increases the expression of Sirt1 and regulates the expression of several genes, which indicate that Igf1 signalling could be reduced, probably leading to increased activity of Foxo1 and enhanced defence against oxidative stress. Quercetin increases the lifespan of *C. elegans* and *Saccharomyces cerevisiae*, and it modulates PI3K/Akt activation, Foxo1 phosphorylation, mTOR signalling and oxidative stress defence, as has already been seen in previous investigations.

Therefore, it is concluded that quercetin could be a CR mimetic, as it and CR address similar mechanisms.

In conclusion, our results show that in *silico* analyses can be used to identify potential CR mimetics. This is in sense of ethical principles of animal welfare and avoids additional animal studies because already existing array data are used. Unfortunately, however, our results are limited to a few substances because no further data are available on ArrayExpress or GEO. Thus the results of our in *silico* analysis are limited to the availability and quality of existing data.

Zusammenfassung

Kalorienrestriktion (CR, engl.: caloric restriction) führt bei verschiedenen Modellorganismen zu einer gesteigerten Lebenserwartung und einer verlängerten Gesundheitsspanne (Lebensspanne ohne altersbedingte Erkrankungen). Erste Studienergebnisse aus Rhesusaffen und Menschen sind bereits vielversprechend. Allerdings ist davon auszugehen, dass eine langfristige Kalorienrestriktion für Menschen nur sehr schwer durchzuhalten ist, weil Lebensmittel so gut wie immer verfügbar sind und CR häufig mit Nebenwirkungen wie Hunger und frieren einhergeht. Deshalb ist es von Interesse Mimetika für CR zu identifizieren, die ebenfalls einen positiven Einfluss auf die Lebenserwartung und die Gesundheitsspanne haben, ohne dabei die Kalorienzufuhr zu beeinflussen. Bisher konnten potentiell wirksame pharmakologische Substanzen wie Rapamycin und Metformin identifiziert werden, deren Einnahme jedoch mit zum Teil schweren Nebenwirkungen verbunden ist. Deswegen ist die Suche nach potentiell wirksamen Substanzen aus Lebensmitteln wie etwa sekundären Pflanzenstoffen besonders interessant.

Ziel dieser Studie ist es, eine neue Methode zu entwickeln, mit der potentielle CR Mimetika in *silico* (d.h. computergestützt) identifiziert werden können. Hierfür wird die transkriptionelle und funktionelle Übereinstimmung zwischen CR und verschiedenen Substanzen analysiert. Dazu wird auf bereits publizierte Ergebnisse (Expressionsprofile) früherer Studien zurückgegriffen. Datenbanken wie ‚arrayexpress‘ und ‚Gene expression omnibus (GEO)‘ wurden genutzt um entsprechende Microarray-Experimente zu identifizieren, die den Einfluss von CR bzw. sekundären Pflanzenstoffen auf die Genexpression untersucht haben.

Von den als relevant für die Fragestellung eingestuften Studien untersuchten neun den Einfluss von CR, vier den Einfluss von sekundären Pflanzenstoffen, die mit der Diät zugeführt wurden (Resveratrol, Phloridzin und Quercetin), und eine den Einfluss von sekundären Pflanzenstoffen, die intraperitoneal verabreicht wurden (Quercetin und Genistein). Die Ergebnisse dieser Experimente wurden mit Hilfe des R Pakets ‚RankProd‘ zusammengefasst, um Gene zu identifizieren, deren Expression konsistent über alle Behandlungen (CR und sekundäre Pflanzenstoffe) reguliert wurde. Die so erhaltenen Genlisten wurden miteinander verglichen, um die transkriptionelle und funktionelle Übereinstimmung herauszufinden.

Unsere Ergebnisse zeigten, dass Resveratrol die größte Übereinstimmung bezüglich der Genexpression zeigte, und damit als potentielles Mimetikum am ehesten in Frage käme. Allerdings konnte die in der Literatur beschriebene Induktion von Sirt1 (ein bekannter Vermittler einiger CR Wirkungen) oder die Modulation des Igf1 Signalweges

durch Resveratrol hier nicht gezeigt werden. Interessanterweise wurde für Quercetin gezeigt, dass trotz geringer numerischer Übereinstimmung in der Genexpression eine übereinstimmende Modulation wichtiger metabolischer Schlüssel signalwege analog zu CR bestand. Quercetin steigerte die Expression von Sirt1 und modulierte mehrere Gene innerhalb des Igf1 Signalweges, was zu einer gesteigerten Foxo1 Aktivität und damit zu einer verstärkten Abwehr oxidativer Schäden führen kann. Es ist bereits bekannt, dass Quercetin einen positiven Effekt auf die Lebenserwartung von *C. elegans* und *Saccharomyces cerevisiae* und einen regulierenden Einfluss auf die PI3K/Akt Aktivierung, die Foxo1 Phosphorylierung, den mTOR Signalweg und die Abwehr von oxidativem Stress hat. Demnach wird Quercetin hier als ein potentielles CR Mimetikum herausgestellt da ähnliche molekulare Signalwege wie durch CR moduliert werden.

Zusammenfassend kann gesagt werden, dass der hier vorgestellte *in silico* Ansatz grundsätzlich geeignet ist, um potentielle CR Mimetika zumindest hypothetisch zu identifizieren. Dies ist im Sinne ethischer Grundregeln des Tierschutzes und vermeidet zusätzliche tierexperimentelle Studien, indem bereits vorhandene Array Daten verwendet werden. Allerdings ist die Aussagekraft der *in silico* Analyse beschränkt auf die in den Datenbanken (‘arrayexpress’ und ‘GEO’) frei verfügbaren Datensätze und abhängig von der Qualität der Daten.

Table of Contents

List of Tables	IV
List of Figures	VI
Abbreviations	VII
1. Introduction	1
1.1. Mechanisms of CR effects.....	2
1.2. Mimetics	7
1.3. Identification of CR mimetics	10
1.3.1. Methods for identification of relevant biological functions addressed by the differently expressed genes (enrichment tools).....	11
1.3.2. Advantage of <i>in silico</i> analysis for identifying CR mimetics	14
1.4. Aim of the present analysis	14
2. Methods	15
2.1. Public data collection	15
2.2. Pre-processing of raw data and standardisation of array information	15
2.3. Identification of significantly regulated genes observable over all data sets.....	17
2.4. Functional Analysis	18
2.4.1. Overrepresented pathways.....	18
2.4.2. Functions of the most strongly regulated CR genes identified by fold change or q-value.....	19
2.4.3. Functions of the genes which are investigated for CR and all dietary phytonutrients	20
2.5. Investigation of genes with special regard for longevity	20
2.6. Analysis of transcription factor binding sites	20
3. Results	21
3.1. Public Data Collection.....	21
3.2. Comparison of analysis methods used to identify significantly regulated genes/Finding the appropriate p-value test.....	24
3.3. Functional analysis	27
3.3.1. Identification of over-represented pathways for the regulated CR genes with CPDB.....	28

3.3.2. Determining the overlap between CR and the dietary plant bioactives regarding the over-represented pathways.....	33
3.3.3. Analysis of the top genes up- and downregulated by CR identified with fold change (> 1.5)	42
3.3.4. Overlap between the top 50 regulated genes identified by q-value.....	49
3.3.5. Calculation of Pearson correlation for gene expression between CR and phytonutrients in over-represented pathways for a better comparison between phytonutrients and their mimicking effect	53
3.4. Analysis of longevity genes in mouse	59
3.5. Transcription factor binding sites (TFBs)	60
4. Discussion	63
5. Conclusion	68
References	70
Appendix.....	100
A: Number of regulated genes by the different interventions dependent on the used test.....	100
B: Overlapping genes between C and the plant bioactives identified by the different analysis methods	101
C: All significantly regulated genes by caloric restriction which are identified in at least one approach sorted by alphabet	108
D: Pathways mentioned by CR in overrepresentation analysis and the number of overlapping genes with plant bioactives.....	127
E: Pathways which are excluded from CR analysis	145
F: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and resveratrol.....	146
G: Overlapping genes between CR and resveratrol assigned to the overrepresented pathways mentioned in separate and overlapped analysis	147
H: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and phloridzin.....	148
I: Overlapping genes between CR and phloridzin assigned to the overrepresented pathways mentioned in separate and overlapped analysis	149

J: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and dietary quercetin	151
K: Overlapped genes between CR and dietary quercetin assigned to the overrepresented pathways mentioned in separate and overlapped analysis	151
L: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and quercetin administered intraperitoneally	151
M: Overlapping genes between CR and quercetin administered intraperitoneally assigned to the overrepresented pathways mentioned in separate and overlapped analysis	154
N: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and genistein administered intraperitoneally.....	155
O: Overlapping genes between CR and genistein administered intraperitoneally assigned to the overrepresented pathways mentioned in separate and overlapped analysis	157
P: All significantly regulated genes by caloric restriction with a fold change > 1.5 and the regulation of these genes by the different phytonutrients	158
Q: Overrepresented pathways identified by the regulated CR genes with fold change >1.5	163
R: Overlap between the upregulated CR genes (q-value < 0.05) based on the entrez IDs which are investigated by all dietary interventions (15552 IDs)	165
S: Overlap between the downregulated CR genes (q-value < 0.05) based on the entrez IDs which are investigated by all dietary interventions (15552 IDs).....	178
T: Pearson correlation values between CR and the plant bioactives (including synergistic analysis) for all pathways which include at least one CR gene with a fold change > 1.5	196
U: Transcription factor binding sites (TFBs) with a z-score > 5 or a Fischer-score > 5 addressed by the upregulated CR genes and the overlap with the phytonutrients .	206
V: Transcription factor binding sites (TFBs) with a z-score > 5 or a Fischer-score > 5 addressed by the downregulated CR genes and the overlap with the phytonutrients	209
W: All mentioned functions / biological processes revealed by UniProt database for the transcription factors (TF), which address the top TFBs of caloric restriction	210

List of Tables

Table 1: Overview of the approaches used in the functional analysis	19
Table 2: Overview of the characteristics of the analyzed caloric restriction studies	22
Table 3: Overview of the characteristics of the plant bioactive studies.....	23
Table 4: Number of genes that are regulated by caloric restriction or dietary plant bioactives (different analysis methods).....	25
Table 5: Total and percental ¹ overlap in gene expression between caloric restriction and plant bioactives.....	26
Table 6: Total and percental ¹ overlap in gene expression regulation between caloric restriction and plant bioactives based on the results of the rank product analysis	27
Table 7: Total and percental overlap between the pathways affected by caloric restriction and plant bioactives in separate analyses	34
Table 8: Overlapping pathways between CR and plant bioactives identified in separate analysis	35
Table 9: Summarized results of pathway analysis of the overlapping genes of CR and resveratrol.....	38
Table 10: Genes that are regulated by caloric restriction and resveratrol in selected pathways.....	38
Table 11: Summarized results of pathway analysis of the overlapping genes of CR and phloridzin.....	40
Table 12: Genes that are regulated by caloric restriction and phloridzin in selected pathways.....	41
Table 13: Summarized results of pathway analysis of the overlapping genes of CR and dietary quercetin	41
Table 14: Genes that are regulated by caloric restriction and dietary quercetin in selected pathways	41
Table 15: Genes that are regulated by caloric restriction and quercetin administered intraperitoneally in selected age-related pathways.....	42
Table 16: Summarized results of pathway analysis of the overlapping genes of CR and quercetin administered intraperitoneally	43
Table 17: Summarized results of pathway analysis of the overlapping genes of CR and genistein administered intraperitoneally	44

Table 18: Summarized results of pathway analysis of the top regulated genes of caloric restriction	45
Table 19: Pathways and biological functions of the overlapping genes between the top 50 upregulated genes of caloric restriction and the plant bioactives.....	51
Table 20: Pathways and biological functions of the overlapping genes between the top 50 downregulated genes of caloric restriction and the plant bioactives	52
Table 21: Number of regulated genes based on the q-value or the q-value in combination with a fold change (FC) > 1.5	53
Table 22: A.)Pearson correlation between CR and the plant bioactives (including synergistic analysis) based on all 15552 IDs; B.) Number of pathways with a high Pearson correlation between caloric restriction (CR) and the plant bioactives (total: 358 CR pathways)	56
Table 23: Pathways with a high Pearson correlation (≥ 0.9) in gene expression between caloric restriction (CR) and the plant bioactives	57
Table 24: Influence of caloric restriction and plant bioactives on longevity genes	59
Table 25: CR longevity genes which are influenced by at least one of the plant bioactive.....	59
Table 26: Total and percental overlap in transcription factor binding sites (TFBs) putatively affected by CR and the plant bioactives	60
Table 27: Top transcription factor binding sites (TFBs) putatively affected by upregulated genes of caloric restriction identified with the z- or F-score and the overlap with the plant bioactives	61
Table 28: Top transcription factor binding sites (TFBs) putatively affected by downregulated genes of caloric restriction identified with the z- or F-score and the overlap with the plant bioactives	61
Table 29: Overview on the transcription factors and their biological function	62

List of Figures

Figure 1: Metabolic changes and underlying molecular mechanisms upon caloric restriction	3
Figure 2: Data collection, preprocessing, standardization and analysis methods	16
Figure 3: Venn diagram of the shared gene pool between CR and the dietary plant bioactives	21
Figure 4: Venn diagram of the upregulated CR genes identified from rank product analysis with the different gene pools	24
Figure 5: Venn diagram for the downregulated CR genes identified from rank product analysis with the different gene pools	24
Figure 6: Overlap and opposite regulation of up- and downregulated CR genes in presence of plant bioactives.....	26
Figure 7: Overview of the potential mechanism mediated by Nnmt.....	39
Figure 8: Heat map showing the fold change of selected genes regulated by caloric restriction and plant bioactives.....	46
Figure 9: Interaction of the overlapped genes between caloric restriction and quercetin (administered per diet).....	50
Figure 10: Heat map showing the fold changes of all upregulated CR genes with a fold change > 1.5 in presence of the different plant bioactives (individual and synergistic analysis).....	54
Figure 11: Heat map showing the fold changes of all downregulated CR genes with a fold change > 1.5 in presence of the different plant bioactives (individual and synergistic analysis)	55
Figure 12: Overlap between CR and plant bioactives including synergistic effects.....	56
Figure 13: Heat map of genes of the pathway ‘Regulation of Insulin-like Growth Factor (IGF) Transport and Uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)’ (Reactome)	58
Figure 14: Heat map of genes of the pathway ‘Nicotinate and nicotinamide metabolism - Mus musculus (mouse)’ (KEGG)	58

Abbreviations

AGEs	advanced glycosylation end-products
Akt	Protein kinase B
AMPK	adenosine monophosphate-activated protein kinase
CPDB	online tool ConsensusPathDB
CR	caloric restriction
Cry1	cryptochrome 1
DAVID	Database for Annotation, Visualization and Integrated Discovery
FDR	False discovery rate
Foxo	forkhead member of the class O
G6pc	glucose 6-phosphatase
GH	growth hormone
Ghr	growth hormone receptor
Gi	genistein administered intraperitoneal
Glut4	glucose transporter type 4
GO	Gene ontology
GSEA	gene set enrichment analysis
HIF-1 α	hypoxia inducible factor 1-alpha
Igf1	insulin-like growth factor-1
Igflals	insulin-like growth factor binding protein, acid labile subunit
Igfbp1	Insulin-like growth factor binding protein 1
Irs1	insulin receptor substrate 1
MEA	modular enrichment analysis
MNAM	N1-methylnicotinamide
mTOR	mechanistic target of rapamycin
NA	nicotinic acid
NAD	nicotinamide adenine dinucleotide
NAM	nicotinamide
NF- κ B	nuclear factor κ B
Nnmt	nicotinamide N-methyltransferase
Nrf1	nuclear respiratory factor 1
Nrf2	Nuclear factor erythroid 2-related factor 2
P	phloridzin
Pepck	phosphoenolpyruvate carboxykinase
Per1/2	period circadian clock 1/2
PGC-1 α	peroxisome proliferator-activated receptor gamma coactivator 1-alpha
PI3K	phosphoinositide 3-kinase

PPAR	peroxisome proliferator activated receptor
Qd	quercetin administered per diet
Qi	quercetin administered intraperitoneal
R	resveratrol
ROS	reactive oxygen species
S6k	ribosomal protein 6 kinase
SAM	S-adenosylmethionine
SEA	singular enrichment analysis
Sirt	sirtuin
Srebp1c	sterol regulatory element binding protein 1c
TFAM	transcription factor A, mitochondrial
TF	transcription factor
TFBs	transcription factor binding sites
TGF beta	transforming growth factor beta
TNF- α	tumor necrosis factor alpha

1. Introduction

Life expectancy has been increasing for several decades. However, large differences nonetheless exist between individuals and between regions and countries (1). This leads to the question of which factors are responsible for the determination of lifespan. In addition to genetic disposition, lifestyle and nutrition seem to be important. In antiquity, Hippocrates had already recognised a relationship between high calorie intake and early death (2; 3). Since then, many more hints have become available that food amount and quality are associated with health, ageing and lifespan (3). The first studies to investigate the relationship between food or calorie restriction (CR) and ageing or lifespan occurred in the early 20th century. OSBORNE et al. (1917) showed that food restriction for four female rats increased lifespan, retarded reproduction and reduced growth rate (4; 5). Moreover, MCCAY et al. (1935) observed an increased lifespan in rats following a reduced calorie intake based on a diet containing 20% indigestible cellulose (2; 6). Currently, many more investigations are available indicating that CR increases the maximum and/or median lifespan in many model organisms, such as yeast, flies, worms, fish, mice and rats (2). CR refers to reducing calorie intake by 10–50% for animals fed ad libitum. These diets are often supplemented with micronutrients to prevent deficits and malnutrition (2).

However, it is important to not only increase lifespan but also enhance the occurrence of healthy ageing. Age-related diseases increase costs for health care systems and decrease the quality of life drastically (7; 8; 9). Interventions to increase both lifespan and healthy ageing have been the subject of research for several decades. CR seems to be effective in preventing physiological decline and age-related diseases such as cancer, cardiovascular diseases, neurodegenerative diseases (e.g., Alzheimer's disease) and insulin resistance (which is linked to diabetes mellitus) as well, leading to increased healthspan (lifespan without age-related diseases) (10; 11; 12; 13).

If CR is able to increase lifespan and improve health in model organisms, then it is important to investigate whether CR is beneficial in humans, too. Experiments with humans and rhesus monkeys are still ongoing. The available results for rhesus monkeys to date are inconsistent regarding lifespan but already indicate that CR improves health, cardiovascular factors and insulin sensitivity (2; 12; 14). The differences in lifespan expansion could be explained by differences in dietary composition and genetic background (14). The CALERIE (Comprehensive Assessment of the Long-term Effects of Reducing Intake of Energy) study in humans indicates that CR also shows health benefits but that bone mineral density is reduced (14). Nonetheless, further investigations are necessary to determine conclusively whether CR improves health- and increases lifespan in humans too. However, even if CR is effective, it will be very diffi-

cult for humans to achieve a lifelong restriction in calorie intake because CR shows some side effects (such as hunger and sensitivity to cold) and because food is available all the time for most people in Western countries. Therefore, identifying other interventions that can improve health and increase lifespan could be of interest. To determine these factors, it may be useful to examine the underlying mechanisms of beneficial CR effects.

1.1. Mechanisms of CR effects

To identify mechanisms that could be responsible for the lifespan-expanding effect of CR, it is important to know what happens during ageing and which risk factors promote the development of age-related diseases. Ageing is known to be associated with increased metabolic and oxidative stress, higher concentrations of reactive oxygen species (ROS), reduced antioxidant defence, enhanced lipid and DNA damage, reduced capability in DNA damage-repair mechanisms, mitochondrial dysfunction, chronic inflammation, nicotinamide adenine dinucleotide (NAD⁺) depletion, and higher rates of apoptosis (13; 15; 16). Interestingly, previous studies indicate that CR regulates many of these mechanisms, such as apoptosis, ROS production and defence systems (17). However, how CR influences these factors and which factors are necessary for improved health and increased lifespan are not clear. SINCLAIR (2005) reviewed the many hypotheses that exist regarding the potential mechanisms of CR (17). For example, endocrine changes based on reduced fat mass, reduced ROS availability or increased cell survival could be responsible for CR's longevity effect. However, some studies have refuted these hypotheses and most of the mentioned mechanisms alone have not been able to explain the multitudinous effects of CR, which have made it impossible until now to completely identify the underlying CR mechanisms (17). However, the most accepted theory to date is the hormesis hypothesis, which states that CR functions as a mild stressor that activates several defence mechanisms. This activation protects against the effects of factors that are involved in the ageing process or in the development of age-related diseases (17). Nonetheless, further studies are necessary to finally identify the CR mechanisms.

However, even if the CR effect is not completely understood, many pathways and key players have been identified that are most likely involved, including sirtuins, the mechanistic targets of rapamycin (mTOR) signalling, adenosine monophosphate-activated protein kinase (AMPK) signalling, forkhead members of the class O (Foxos), peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1α), growth hormone (GH) / insulin-like growth factor-1 (Igf1) signalling, Ras-protein kinase A (PKA), glycolysis, autophagy and possibly fibroblast growth factor 21 (Fgf21) (2; 10; 12;

14; 15; 18; 19). Moreover, CR is known to influence the regulation of oxidative stress, suppress inflammation, have proangiogenic effects and increase DNA repair mechanisms and protein synthetic capacity (20; 21). Furthermore, CR may increase the serum concentration of glucocorticoids (e.g., cortisol), which results in the activation of anti-stress and anti-inflammatory pathways (22). In the following, it will be provided an overview of the potential mechanisms that have been identified. Figure 1 summarises these results.




Figure 1: Metabolic changes and underlying molecular mechanisms upon caloric restriction

GH and Igf1 signalling; Foxo1

CR lowers the levels of growth hormone (GH), Igf1, plasma insulin and plasma glucose, which leads to reduced activity in GH/Igf1 signalling, improved glucose homeostasis and increased insulin sensitivity (20; 12). GH activity is associated with ageing, and reduced plasma levels of GH and Igf1 or reduced Igf1 signalling leads to an increased lifespan (21; 23). This is supported by BABAR et al. (1999), who showed that an inhibition of the phosphoinositide 3-kinase (PI3K) pathway is associated with longevity in *C. elegans* (24; 25). PI3K/Protein kinase B (Akt) signalling is a target of GH/Igf1 signalling, which leads, for example, to the activation of mTOR/ribosomal protein S6 kinase (S6k) signalling (12). Moreover, activation of PI3K/Akt signalling leads to

the inactivation of Foxo1 by phosphorylation (26). Foxos, especially Foxo1, are involved in oxidative stress responses, apoptosis, cell cycles, detoxification and immune homeostasis (27; 26). Foxo1 increases the expression of antioxidant genes such as superoxide dismutase (Sod), promotes the scavenging of ROS, and prevents damage to DNA and cells. Moreover, Foxos interacts with Sirt1, another important factor for the effects of CR (26). In particular, the increased oxidative defence mechanisms could be important for the lifespan-expanding effect of CR because ROS generated in mitochondria may promote ageing processes (15). That CR results in increased oxidative defence was already mentioned by MITCHELL et al. (2015) (28). Moreover, Igf1-dependent inhibition of daf-16 (an orthologue of Foxo) decreases lifespan in *C. elegans* and *D. melanogaster* (12; 29).

Thus, the CR-mediated inhibition of GH/Igf1 signalling is associated with increased autophagy, reduced plasma glucose, increased hepatic insulin sensitivity, increased resistance against oxidative stress, increased expression of antioxidant genes and reduced mTOR signalling (30; 12; 20). The effect on autophagy seems to be important for mediating CR's effects as well. CR is already associated with increased autophagy, whereas ageing is associated with a decline in this process (31; 32; 33; 22). Autophagy is necessary for eliminating damaged cell organelles and dysfunctional proteins and for mobilising energy reserves (22; 32). The first mentioned effects are important for cell health and function. MORSELLI et al. (2010) showed that Sirt1 is also necessary for the induction of autophagy by CR (32).

Sirtuins and NAD⁺ metabolism

As mentioned above, sirtuins (especially Sirt1) are important for mediating CR's effects in mammals, yeast and drosophila (34). Sirtuins are class III histone deacetylases that remove acetyl groups from lysine residues (34). These processes are dependent on NAD⁺ and are inhibited by nicotinamide (NAM), a side product of these reactions (34). NAD⁺ is derived de novo from L-tryptophan, nicotinic acid (NA) or NAM (35). NAD⁺ is important for energy production, DNA repair and genomic signalling (16). Sirtuins in general are involved in circadian rhythm, cell cycle regulation and other important metabolic processes (36).

There are seven sirtuins (Sirt1 – Sirt7), but Sirt1, Sirt3 and Sirt6 have been best studied in the context of CR (34; 37; 38). Studies indicate that Sirt1 overexpression induced by CR is responsible for the lifespan-expanding effect of CR, but some investigations also associate Sirt1 with ageing under certain circumstances (34). This indicates that the effects of sirtuins in mammals seem to depend on context, tissue and sirtuin type, so it is not possible to say that sirtuins are always involved in longevity (38).

CR may activate Sirt1 by increasing the NAD⁺ level and upregulating AMPK (38; 12). On the other hand, it is assumed that Sirt1 mediates its effects via AMPK (39). Sirt1 functions as a tumour suppressor protein, enhances mitochondrial biogenesis, increases the catabolism of triglycerides and cholesterol, promotes the repair of DNA damage, inhibits chronic inflammatory response, reduces apoptosis, downregulates hypoxia inducible factor 1-alpha (HIF-1α) transcription factor, increases the expression of Sirt6 by forming a complex with Foxo3a and nuclear respiratory factor 1 (Nrf1), activates gluconeogenesis and fatty acid oxidation, inhibits glycolysis and is involved in regulating macroautophagy and mitophagy (38; 34). These functions can be explained by the many transcription factors targeted by Sirt1 that are involved in cell survival and metabolism. Best known are the tumour suppressor p53, Foxo, nuclear factor κB (NF-κB), HIF-1α, nuclear factor erythroid 2-related factor 2 (Nrf2) and PGC-1α (34).

For example, PGC-1α is assumed to be important for the Sirt1-dependent regulation of glycolysis, gluconeogenesis and fatty acid oxidation (38). Moreover, PGC-1α is involved in mitochondrial biogenesis by activating Nrf1, Nrf2 and transcription factor A, mitochondrial (TFAM). TFAM stimulates mitochondrial DNA replication and mitochondrial gene expression (34). The Sirt1-dependent effect on mitochondrial biogenesis seems to be important for mediating CR's effects (12). Mitochondrial activity depends on the ability to reduce DNA and oxidative damage, because the electron transport chain leads to enhanced production of ROS. That CR influences antioxidant defence mechanisms has already been mentioned above. An increase in mitochondrial biogenesis has already been seen for CR but some experiments have not been able to replicate this result (34; 40; 41).

The anti-inflammatory effect of Sirt1 is mediated by inhibiting tumour necrosis factor alpha (TNF-α) and NF-κB, and the inhibition of lipogenesis and of the catabolism of triglycerides are mediated by the inhibiting of sterol regulatory element binding protein 1c's (Srebp1c's) actions as well as the activation of Foxo1 (38). The effect on apoptosis seems to be mediated via p53 and Ku70 protein (17).

Sirt3, Sirt4 and Sirt5 are located in the mitochondria and are involved in regulating ATP synthesis, the electron transport chain, mitochondrial biogenesis, apoptosis and ROS formation; stimulating the mitochondrial antioxidant defence system, gluconeogenesis and fatty acid oxidation; and inhibiting lipogenesis (38; 36; 37). Moreover, Sirt3 seems to be important for the phosphorylation of AMPK (36).

Sirt6 is involved in DNA repair mechanisms and is important for maintaining genomic stability. Moreover, Sirt6 acts as a tumour suppressor protein and induces apoptosis in cancer cells but not normal cells (42). Furthermore, Sirt6 has anti-inflammatory activity,

which is probably mediated by the inhibition of TNF- α (38). Sirt6 seems to inhibit Igf signalling as well by inhibiting c-Jun activity (43). Thus, Sirt6 could also be involved in the lifespan-expanding effect of CR. Sirt6 overexpression has been shown to increase the lifespans of male mice but not female mice (44). That CR increases Sirt6 levels has already been seen but some hints indicate that this is caused not by increased expression but through an enhanced stability of Sirt6 (38; 45).

AMPK and mTOR activity

As mentioned above, AMPK could be activated by Sirt1. However, CR also increases AMPK expression by influencing the AMP/ATP ratio, which results in deprived plasma glucose (12; 46). AMPK increases mitochondrial biogenesis, promotes insulin sensitivity and is involved in shifting energy production from glycolysis to, for example, fat oxidation (39; 46). Moreover, AMPK is able to activate PGC-1 α by phosphorylation independent of Sirt1 and to inhibit the mechanistic target of rapamycin (mTOR) signalling, which is seen for CR as well (33; 46; 34). Moreover, AMPK seems to promote the expression of small heterodimer protein (SHP) and decreases the expression of phosphoenolpyruvate carboxykinase (Pepck) and glucose 6-phosphatase (G6pc), which are involved in regulating gluconeogenesis and glycolysis (47). Moreover, AMPK is assumed to promote glucose transporter type 4 (Glut4) translocation to the plasma membrane, which increases the insulin-independent glucose uptake (48). AMPK activity is positively associated with lifespan (12; 49; 50; 51). However, CR in mammals is not associated with increased AMPK activity in any investigations, indicating that other regulatory factors (besides CR) are necessary (12).

mTOR is involved in regulating metabolism and cellular growth (33). Activated mTOR signalling inhibits autophagy and stimulates cell growth, cell proliferation and protein synthesis (12). The inhibition of mTOR by CR is associated with an increased lifespan in many organisms (e.g., mice), and dysregulation of this pathway is associated with age-related diseases like cancer and diabetes mellitus (52; 53). The CR-mediated inhibition of mTOR signalling may mediate its beneficial effects by modulating protein synthesis, autophagy and ribosomal biogenesis but the exact underlying mechanism is not clear. However, targeting mTOR complex 1 (mTORC1), S6k and eukaryotic translation initiation factor 4E-binding protein 1 (4E-bp1, a translational repressor) is necessary for these effects (33). Furthermore, CAI and WEI (2016) could demonstrate that RNA polymerase III-inhibiting protein Maf1 (Maf1) is involved in the lifespan-expanding effect of mTOR inhibition in yeast but not in *C. elegans* (52). Maf1 is a stress-responsive transcriptional factor that inhibits tRNA synthesis by binding to RNA polymerase III complex (52).

Glycolysis and gluconeogenesis

The previous explanations indicate that glucose metabolism is important for lifespan regulation. Investigations show that high glucose availability is associated with a decreased lifespan, whereas the inhibition of glycolysis by 2-deoxy-D-glucose, glucosamine or glycerol is associated with an increased lifespan in *C. elegans* (54; 55). Moreover, glycolysis in *C. elegans* is increased in aged organisms, whereas gluconeogenesis is decreased (56). Glycolysis could be involved in ageing processes by producing advanced glycosylation end-products (AGEs) and methylglyoxal (15). AGEs are associated with diabetes mellitus, brain ageing and Alzheimer's disease (15). Methylglyoxal is associated with ROS production, increased cellular peroxide concentration and enhanced oxidative damage (15). How CR influences glycolysis and gluconeogenesis is not fully understood, although it is known that AMPK, mTOR and Foxo1 are involved in this process (54; 56; 27). For example, Foxo increases the expression of gluconeogenic genes like G6pc and Pepck (57).

Detoxification system

As mentioned above, ageing is associated with increased damage of DNA and macromolecules. Therefore, ageing may lead to a reduction in detoxification capabilities (10). CR increases the expression of phase I and phase II enzymes, which could also be responsible for the age-preventing effects of CR (10). This is supported by the observation that the mRNA expression of these genes is decreased in the livers of aged mice (10). In contrast, long-lived Little mice (*Ghrhr^{lit/lit}*) showed upregulation in the genes of detoxification enzymes, and JIANG et al. (2013) showed that the expression of these genes is downregulated by GH/Igf1 signalling (58). This is another hint that the inhibition of GH/Igf1 signalling is important for the beneficial effects of CR.

Circadian clock

Furthermore, CR influences the expression of circadian clock genes. Period circadian clock 1 (Per1), Per2 and cryptochrome 1 (Cry1) are upregulated in the hypothalamus under CR conditions (59). This is assumed to be important for lifespan, since Per1 and Per2 are tumour-suppressor genes (59). The reduction of the TNF- α signalling pathway triggered by CR may be responsible for the changes in clock genes' expression (59). Circadian clock disruption is associated with ageing, and CR is able to normalise and synchronise the circadian clock, which is probably responsible for the longevity effect (59).

1.2. Mimetics

The identification of these mechanisms could be useful to identify CR mimetics. This is necessary because it is very difficult for humans to reach lifelong CR. In addition to the

large availability of food which entices humans to eat, there are many side effects of CR. These are, for example a reduced satiety; higher sensitivity to cold; hypotension; reduced bone health; reduced libido; menstrual irregularities; infertility; reduced stamina and strength; slow wound healing; and psychological problems such as depression, emotional deadening and irritability (60; 25; 20; 47). Therefore, it is of interest to find substances which are able to mimic CR's effect on metabolism, stress response, healthy ageing and longevity without affecting calorie intake (47). Many substances have been analysed as potential CR mimetics, such as inhibitors of fat digestion (chitosan, orlistat and mannan-oligosaccharides), inhibitors of carbohydrate digestion (acarbose), inhibitors of glycolytic pathways (2-deoxy-d-glucose, glucosamine, 3-bromopyruvate, Iodoacetate and mannoheptulose), biguanides (phenformin, buformin, and metformin), polyamines (spermidine and thermospermine), thiazolidinediones, pegvisomant (which targets the GH receptor), sirtuin activators (resveratrol, nicotinamide and oxaloacetic acid), agmatine (a product of arginine decarboxylation), 17 β -estradiol, glycerol, ursolic acid and rapamycin (47; 54; 61; 62; 63). However, many of these substances have side effects such as digestive issues (indigestion, loose stools, excessive flatulence and faecal incontinence), suppression of the immune system and toxicity in high dosages (47). Some of the potential CR mimetics are discussed in the following.

Glycerol

Glycerol increases lifespan, enhances mitochondrial activity, improves stress resistance and prevents the accumulation of protein aggregates (54). Moreover, glycerol decreases glycolysis and increases oxidative phosphorylation which indicates that glycerol could be a CR mimetic. However, in high dosages, glycerol shows toxic effects which are associated with a decreased lifespan. If further studies confirm that glycerol is a CR mimetic, determining the best dosage in humans will be quite important.

Glucosamine

Glucosamine is known to decrease glycolysis but to a much lesser extent than does 2-deoxy-d-glucose. Moreover, glucosamine administration increases lifespan in nematodes and mice (55). Furthermore, glucosamine is associated with a decrease in overall mortality in humans as well (64; 65). D-glucosamine is used in the treatment of osteoarthritis in humans and shows no side effects besides of occasional allergic reactions in contrast to 2-deoxy-d-glucose which is cardio-toxic and increases the occurrence of tumours (55; 12). WEIMER et al. (2014) could further show that glucosamine administration is associated with AMPK activation, which leads to increased mitochondrial biogenesis and enhanced mitochondrial respiration of fatty acids and amino acids. This is followed by an increase in ROS formation and in the expression of genes of the

stress defence system. Interestingly, in nematodes, ROS formation is an important requirement for the lifespan-expanding effect of glucosamine.

Rapamycin

Until now, only rapamycin, an inhibitor of the mTOR pathway, has seemed to be able to mimic CR's effects on longevity consistently in both genders (47; 53; 60; 66; 67). Rapamycin is known to inhibit mTOR, mTOR complex 1 and mTOR complex 2 (by chronic administration of rapamycin); this inhibition is associated with increased lifespan in yeast and mammals (25; 20; 18). Studies indicate that rapamycin is very effective in expanding lifespan but that it has many side effects. This means dysregulation of glucose and insulin homeostasis, cataracts and immunosuppression (18). Further investigation is necessary to determine the dose at which beneficial effects occur without side effects. Moreover, CR and rapamycin can prevent or reverse cardiac ageing (33). Rapamycin is normally used for immunosuppression and cancer treatment (18).

Metformin

Metformin is discussed as a potential CR mimetic as well. This biguanide is normally used in the treatment of type 2 diabetes (18; 68). Metformin decreases gluconeogenesis, increases fatty acid oxidation, enhances peripheral glucose uptake and suppresses glucose uptake in the gastrointestinal tract (47). How metformin mimics CR is not clear, but metformin is theorized to activate AMPK, which inhibits mTOR signalling (18; 20; 39). AMPK activation may be mediated by the metformin-induced inhibition of complex 1 of the respiratory chain, which leads to a decrease in ATP production and an increased AMP/ATP ratio (39; 69). On the other hand, metformin activates autophagy via Sirt1 independent of AMPK (69). Both CR and metformin increase mitophagy in *ob/ob* mice (69). This seems to be mediated by a reduction in endoplasmic reticulum stress markers and the expression of cytosolic p53.

To be a CR mimetic, metformin must also influence lifespan and the occurrence of age-related diseases. Metformin decreases all-cause mortality in diabetic and cardiovascular disease patients; enhances lifespan in worms and mice; and depresses the occurrence of cancer, cardiovascular disease and chronic kidney disease (47). However, no effect on longevity is seen in drosophila and rats, and the meta-analysis of BOUS-SAGEON et al. (2012) found that metformin had no beneficial effect on overall mortality (70; 47). These different results can be explained by the doses of metformin. For example, MARTIN-MONTALVO et al. (2013) showed that a high dosage (1% of diet) decreased survival and that a low dosage (0.1% of diet) led to increased lifespan and improved health (47; 68). Further studies are necessary to clarify whether metformin is

a CR mimetic and which minimum dosage is necessary to achieve the health-improving and lifespan-expanding effect.

Dietary plant bioactives such as resveratrol

Dietary plant bioactives may also be able to mimic CR's effect because various phytonutrients protect against age-related diseases (46; 71; 72). Dietary plant bioactives are components of fruits, vegetables, nuts, cereals and most foods which are made from herbal ingredients like juice, wine, coffee and chocolate (13). Some secondary plant bioactives are already mentioned in context of CR such as resveratrol (18; 73).

Resveratrol is present in mulberries, peanuts, grapes and red wine; it has been shown to extend lifespan in yeast, worms, flies, fish and mice (73; 74; 75; 76; 77; 78; 79; 46; 80). However, in mice, the results are contradictory because PEARSON et al. (2008) cannot find an effect on longevity among mice fed a normal diet – though several beneficial effects on health are seen (81). Other experiments show no effect on lifespan as well (82; 66; 73). However, resveratrol improves health in mammals, including a reduction in cardiovascular diseases, an improvement in bone health, a reduced cataract incidence and increased insulin sensitivity (78; 81; 46). The phenotypic and molecular effects of resveratrol seem to partially overlap with those seen in CR mice (78). Resveratrol is theorized to mediate its CR-mimicking effects via sirtuins by activating AMPK, but the exact underlying mechanism is still unclear (18; 11; 73). Moreover, resveratrol downregulates glycolysis and both Igf1 and mTOR signalling, and it upregulates Stat3 signalling, just as CR does (46). However, resveratrol and CR show different effects on cell-cycle regulation and apoptosis (46). Nonetheless, resveratrol induces gene expression changes similar to CR (73; 11).

These studies indicate that resveratrol is a potential CR mimetic. However, the differences in study design (e.g., in study size, dosage and application of resveratrol) could be responsible for the conflicting results. Therefore, further studies are necessary to investigate toxicity, dose, bioavailability and the underlying mechanism (18). Other phytonutrients such as epicatechin, epigallocatechin, quercetin, curcumin and Korean mistletoe extract have also been discussed in the context of longevity and beneficial health effects (13; 83; 84). However, it is not known yet if these substances are CR mimetics.

1.3. Identification of CR mimetics

To identify CR mimetics, *in vivo* experiments could be used to investigate lifespan and metabolic changes. Gene expression analysis could also be useful because gene expression influences proteomes and metabolomes. Moreover, many genes are associated with increased or decreased lifespan (as mentioned by the GenAge database, <http://genomics.senescence.info/genes/> (85)), and CR reverses many of the mRNA

changes triggered by ageing (10). Some of the genes which are influenced by CR (e.g., sirtuins) are already mentioned above. Other genes, such as *Indy* (an acronym for ‘I am Not Dead, Yet’, a homolog in mice: *mIndy*, *Slc13A5*), have been little studied but are promising. Downregulation of *Indy* promotes longevity in *drosophila* and *C. elegans* via mechanisms similar to those in CR (e.g., increasing AMPK activity) (39; 86). *Indy* encodes for a transporter of citrate-cycle intermediates such as citrate, pyruvate and α-ketoglutarate (86). Although genetic regulation of metabolism is not completely understood, gene expression analysis could be very useful in identifying potential CR mimetics, and their impact on lifespan and health can be further investigated *in vivo*.

1.3.1. Methods for identification of relevant biological functions addressed by the differently expressed genes (enrichment tools)

Methods are available which allow investigating the expression changes of few genes in one experiment (e.g., real-time PCR). However, for this it is necessary to know which genes are influenced by CR and which ones are important for mediating the beneficial CR effects. It can be assumed that not all of the relevant genes have been identified. Thus, methods are necessary which allow for the genome-wide investigation of gene regulation to identify novel and important CR genes and to compare expression changes triggered by several interventions. High-throughput genomic, proteomic and bioinformatics scanning approaches (e.g., expression microarray, promoter microarray, proteomics, and ChIP-on-ChIPs) could be used for this (87). However, a problem with these methods is that they could provide lists of up to 1000 differently regulated genes, which would make it difficult to interpret the results. Many bioinformatics methods (e.g., Gene Ontology (GO) (88), Database for Annotation, Visualisation and Integrated Discovery (DAVID) (89), ConsensusPathDB (CPDB; <http://cpdb.molgen.mpg.de>) (92; 93), GSEA version 2.0.10 (<http://www.broadinstitute.org/gsea/downloads.jsp>) (90), BiNGO (<http://www.psb.ugent.be/cbd/papers/BiNGO/Home.html>) (91) and PANTHER (www.pantherdb.org/) (94)) have been developed to facilitate the identification of relevant and over-represented biological functions in the organism (87). Many more enrichment tools are available (reviewed by HUANG et al. (2009)), and the number may continuously increase in coming years (87).

Although these tools have made a major contribution to gene functional analysis, all tools are in the improving stage, and no method is the universal standard in analysis. Thus, it is very difficult for investigators to decide which tool is the most suitable for their analytic needs. Moreover, comparisons between the tools indicate that the provided results could be different. Therefore, users are advised to try different tools (87).

To decide which method is best, it is necessary to identify the aim of enrichment analysis. This requires identifying a group of genes which belong to a certain biological function and which are over-represented in a list of genes (87). Therefore, the interpretation of the results switched from a gene-oriented view to a relevant-gene group-based view. This will identify the most influenced biological functions of the differently regulated genes. For this analysis, three different classes of enrichment tools (which differ in their statistical methods) are available. These are singular enrichment analysis (SEA), gene set enrichment analysis (GSEA) and modular enrichment analysis (MEA) (87). They are explained in detail by HUANG et al. (2009). In SEA, a preselected list of genes (normally determined by p-value or fold change) is investigated. The results are sorted by enrichment p-value (determined by different statistical methods depending on if the genes are over-represented in the investigated gene pool or if there is a random hit). Therefore, SEA can be used to extract the major biological meaning for a list of differently expressed genes. However, one disadvantage of this method is that these tools can provide a large list of interesting biological functions which make it difficult to interpret the results and determine the function of interest. Moreover, it is possible that a biological function is mentioned several times under different terms. Finally, the results strongly depend on the quality of the preselected genes; it is possible that different statistical methods (e.g., a chi-squared test or Fischer test) might obtain a different list of interesting biological functions (87).

GSEA contains the idea of SEA, but the algorithm it uses to calculate enrichment p-value is different. Another modification is that all genes from the microarray experiment are included in the analysis, which means that genes with small expression change are considered. These genes are often ignored in SEA. This will allow an interpretation which is nearer to natural biological circumstances because small expression changes can result in large biological consequences. However, GSEA tools have some disadvantages, too. For example, a summarised biological value (e.g., fold change) for each of the investigated genes is necessary for analysis. This can be a problem when the biological study or the genomic platform is too complex. For example, an array could include several single-nucleoid polymorphisms (SNPs) of a gene with different expression changes and p-values. Another disadvantage is that it is not possible with GSEA to simultaneously investigate multiple factors in a study. Furthermore, the revealed biological functions with the highest p-value include the genes with the greatest expression change. As mentioned above, these genes cannot be assumed to be the most important ones (87).

MEA tools include SEA's basic enrichment calculations but consider the interrelationships of GO terms. This means that terms which address the same biological function

are summarised in one term. This could reveal the biological meaning of a gene list, which is closer to the true nature of the biology. However, a disadvantage of MEA is that terms or genes without strong relationships to other terms or genes could get lost (87).

The explanations above indicate that the choice of the enrichment tool can influence the results of an analysis. However, other factors influence the results of an investigation. One is the analyst. He or she determines the interesting biological functions based on his or her biological knowledge and excludes annotations with high p-values when they do not make sense based on the user's question. Another important fact that should be considered is that a high number of analysed annotations linearly increases the rate of false-positive results. To reduce this, performing multiple test corrections of enrichment p-values is recommended. Examples are Bonferroni, Benjamini-Hochberg, Holm and q-value (87). Moreover, the selection of the background has a big impact on the results because the list of interesting genes is compared with this background. Thus, it is a great benefit when a tool allows an individual background to be uploaded rather than only choosing from preselected ones. Furthermore, no tool provides all the biological aspects which can be obtained from different databases, but some tools (e.g., DAVID and CPDB) do include several of these databases. Examples are the GO (95) database, which provides information about biological processes, molecular functions and cellular components; the BIND (96) database, which provides information about protein-protein interactions; the KEGG (97) and REACTOME (98) databases, which provide information about pathways; the TRANSFAC (99) and JASPAR (100) databases, which give information about gene regulation (e.g., transcription factor binding sites (TFBs)); and the OMIM (101) database, which is used to obtain gene-disease associations. Other databases are not mentioned here (87).

Moreover, it is necessary that all genes are clearly mapped to a gene identifier. If this is not possible, the analysis is strongly impaired. Several identifiers could be available in the analyst's present gene list. Examples are the Affymetrix array identifiers (102), NCBI Entrez gene (103), UniProt (104) and PIR-NREF (105). The interconversion could be necessary when the enrichment tool does not support the present identifier. This conversion could be a great challenge because the ID which belongs to another ID is not always clear. Many tools should facilitate this interconversion (IDconverter (106), Synergizer [<http://llama.mshri.on.ca/synergizer/doc/>] (108) and DAVID ID converter (107)). However, it is possible that some of the IDs cannot be converted (87).

Another fact which influences the results is that not all of the gene interactions and physiological mechanisms have been revealed. This means that repeating the analysis

of a gene list a few years later can lead to completely different results based on the newly discovered interactions. However, gene expression analysis could be a useful tool to identify relevant biological functions in the organism.

1.3.2. Advantage of *in silico* analysis for identifying CR mimetics

Regarding the number of potential substances which can act as CR mimetics, their investigation could last decades because *in vivo* studies, which must be conducted for gene expression analysis, have a high duration (as they are highly dependent on the life expectancy of the analysed animal model), and only a few substances can be analysed at the same time in one experiment. To accelerate the process of identifying potential CR mimetics, an *in silico* analysis could be useful. This approach allows for the analysis of many substances at the same time and for the use of data from previously conducted experiments – regardless of whether they are examined in a CR context. FORTNEY et al. (2012) give an example by searching for potential CR mimetics in *silico* (109).

Databases such as ArrayExpress (<http://www.ebi.ac.uk/ArrayExpress/>) and Gene Expression Omnibus (GEO; <http://www.ncbi.nlm.nih.gov/geo/>) can be used to identify relevant experiments (110; 111; 112). One challenge of *in silico* analysis is that the results of several experiments which investigate the influence of the same substance or intervention need to be summarised to get one biological value (e.g., fold change) for a gene. A requirement for this is that the genes are represented by the same identifier. Several statistical methods are available to obtain significantly expressed genes or fold changes. An example is the RankProd R package, which reveals fold changes and p-values for the genes over many experiments (113). The genes which are significantly expressed in the different interventions can be compared to identify an overlap in gene expression. Furthermore, they can be analysed using different enrichment tools to investigate if the substances address the same biological functions as CR.

1.4. Aim of the present analysis

Our work should provide a new approach which allows for faster identification of potential CR mimetics; these potential mimetics should subsequently investigated *in vivo* to confirm their desired positive effect on ageing, health and lifespan. This analysis is limited to phytonutrients because they may show fewer side effects than pharmaceuticals. Moreover, there are hints that a diet rich in fruits, vegetables and whole grains (such as the Mediterranean diet) – in other words, a diet rich in phytochemicals – is associated with a healthier life (114; 115). Therefore, whether phytonutrients other than resveratrol can mimic CR is of great interest. To our knowledge no *in silico* analysis has investigated the role of dietary plant bioactives as CR mimetics. Therefore, it is our interest to

examine a potential transcriptional overlap between CR and different phytonutrients. In our analysis, an overlap refers to genes which are regulated in the same direction by both interventions (CR and the respective phytonutrient).

In the present thesis a few tools are tried, but it is decided at last to pre-process the raw data with different R packages, to unify the Entrez IDs (based on the annotation information mentioned for the different chips), to consistently identify up- and downregulated genes with the RankProd package and to carry out the enrichment analysis with CPDB. Supplementary information is obtained from NCBI: gene (116). The results could be used to identify new functions of CR and to compare the overlap in gene expression or function between CR and the respective dietary plant bioactive. This indicates whether a phytonutrient could be a mimetic for CR or if the dietary plant bioactive mimics at least one function.

2. Methods

To identify an overlapped gene expression between CR and a phytonutrient, only public data are used, and none of our own experiments are performed. Figure 2 summarises the methods used, which are then further explained in the following sections.

2.1. Public data collection

The databases ArrayExpress (<http://www.ebi.ac.uk/ArrayExpress/>) and GEO (<http://www.ncbi.nlm.nih.gov/geo/>) are used to find gene expression data for experiments which investigate the influence of dietary plant bioactives or CR in murine liver (112; 111; 110). The search items are ‘hepatic’ or ‘liver’ in combination with ‘diet’, ‘food’, ‘dietary’, ‘caloric’, ‘calorie’ and ‘treatment’. The organism *Mus musculus* is used.

The descriptions of all mentioned studies are analysed to identify relevant experiments. The selection criteria are that the experiments must have a control and at least one intervention group and that the intervention and collection of RNA probes must be in the same generation. All experiments in which the plant bioactives are administered by food, intraperitoneally or intravenously are included, but the types of administration are analysed separately. Furthermore, the raw experiment data must be available. If they are not public, the authors are asking for this data.

2.2. Pre-processing of raw data and standardisation of array information

Before performing enrichment analysis, the raw data must be pre-processed (background correction, normalisation, etc.). In the selected experiments, different arrays are used for examining gene expression changes, so different pre-processing methods are necessary. For a better comparison between the arrays, all are quantile normalised with the corresponding Bioconductor packages in R. For Affymetrix arrays, the ‘affy’

and ‘oligo’ packages are used (117; 118). ‘Affy’ contains the ‘rma’ function (robust multi-array average), which includes all pre-processing steps (119; 120; 121), such as background correction, quantile normalisation and summarisation. The ‘oligo’ package is specially designed for the Affymetrix Mouse Gene 1.0 ST array, so this is used instead of ‘affy’ for experiments using this array (118). The ‘oligo’ package contains the ‘rma’ function, too. The annotation information of the Affymetrix arrays is available in different R packages, which are provided by Bioconductor.




Figure 2: Data collection, preprocessing, standardization and analysis methods

Illumina's beadarrays does not work with 'affy' or 'oligo', so the 'lumi' package is used instead (122; 123; 124; 125). First, the data are background-corrected using the 'bgAdjust.affy' method. Then, the data are log2-transformed and quantile normalised. The annotation information is downloaded directly from the Illumina website (<http://www.illumina.com/>) (126). For the Agilent array the 'limma' package is used (127). Before quantile normalisation the data are background-corrected using the 'normexp' method (128; 129). The array annotation information is downloaded together with the raw data.

Before the pre-processed data can be analysed, more steps are necessary. First, all genes must be coded with the same ID system. The Entrez IDs are used because the gene names have changed often in the past. There are many synonyms, so the assignment is not always clear. In some cases, only an array's gene name is mentioned. To get the corresponding Entrez ID, the online tools Synergizer (108), NCBI:gene (116) and Ensembl mouse (http://www.ensembl.org/Mus_musculus/Info/Index) (130) are used. If no Entrez ID is revealed, the array ID is used instead. Finally, the technical replicates for one gene in one biological replicate are summarised by calculating the mean values with Excel.

2.3. Identification of significantly regulated genes observable over all data sets

To identify the genes which are up- and downregulated across all studies, four different methods are used: the t-test and moderated t-test of the limma R package and the Wilcoxon test and rank product analysis of the RankProd R package (113). The rank-product method ranks all genes by the expression levels of all experiments at the same time. Therefore, genes are identified which are probably up- or downregulated over all studies. A p-value is calculated, and, as a multiple-testing correction method, a false discovery rate (FDR, $q < 0.05$) is used.

The t-test is the standard method in gene expression analysis, so the t-test is calculated in Excel; FDR is used as a multiple-testing correction method (cut-off value = 0.05). The moderated t-test is calculated with the 'limma' R package. The moderated t-test differs from the normal t-test in that it moderates the standard errors across the genes using a simple Bayesian model (131). The p-value is also corrected using FDR (cut-off value = 0.05). One contribution of the t-test or moderated t-test is that the data are normally distributed. However, the box plots of the different experiments show that the data are skewed to the left; thus, the t-test should not be used. The Wilcoxon test seems to be the right method to analyse gene expression changes. The results are corrected using FDR as well (cut-off value = 0.05). All three methods are then combined with a fold change ≥ 1.5 .

While the rank product analysis works with all experiments at the same time, the t-test, moderated t-test and Wilcoxon test must be calculated separately for each intervention design. To get a list of up- and downregulated genes, further steps are necessary. First, a one-sided binomial-test is calculated to see if the change in gene regulation is real or random over all experiments ($p < 0.1$). The multiple-testing corrections are omitted because the data have been previously corrected often. Next, two further methods are used to see whether the levels of up- and downregulation differ significantly. The first is a chi-squared test, which is the standard test to identify the significant differences; however, this test does not work with a small number of experiments. Therefore, another method called ‘3-fold bigger’ is also used. This means that if the number of studies which show the same regulation rate is 3-fold bigger than the number of studies which show the opposite regulation rate, the gene is mentioned in the list of regulated genes.

2.4. Functional Analysis

Afterwards, the functions of the significantly regulated genes are analysed to identify the potential mechanisms which are targeted by CR and the respective phytonutrient. First, the over-represented pathways are identified, then the most strongly regulated CR genes are investigated in terms of their function and their expression by the phytonutrients, and finally, the expression regulation of known longevity genes by CR and the phytonutrients is analysed.

2.4.1. Overrepresented pathways

For pathway analysis, the online tool ConsensusPathDB (CPDB, <http://cpdb.molgen.mpg.de/MCPDB>; MM9 release 11.10.2013) is used (92; 93). CPDB uses data from 16 different public resources (e.g., Reactome, KEGG or MouseCyc) and contains interactions curated from the literature. An over-representation analysis is done to identify pathways which are influenced by both CR and dietary plant bioactives. Inclusion criteria are $q < 0.05$ (multiple-testing p-value corrected with FDR) and at least 2 genes involved in the mentioned pathway.

At first, only the significantly regulated CR genes are analysed to identify unknown CR functions. Then, CR and the respective dietary plant bioactive are examined together. The investigated genes in the different interventions are dependent on the array used, so a gene’s expression changes are not necessarily analysed by both the CR and the respective phytonutrient expressions. Therefore, only the genes which are investigated for all interventions are included in the gene pool (see Table 1). The rank product analysis is repeated for these reduced lists. To identify co-mentioned functions, two different approaches are applied (Table 1). First, all significantly regulated genes of CR and

the dietary plant bioactives are analysed separately, and the co-mentioned pathways are discovered (Table 1, separate analysis). Second, only the genes overlapping between CR and the respective dietary plant bioactive are analysed (Table 1, overlapped analysis). All pathway resources mentioned by CPDB are included (KEGG, Reactome, MouseCyc and Wikipathways).

Table 1: Overview of the approaches used in the functional analysis

gene pools (compare Figure 2)	I. all genes which are investigated by CR (bgCR) II. genes investigated by CR and R (bgR) III. genes investigated by CR and P (bgP) IV. genes investigated by CR and Qd (bgQd) V. genes investigated by CR and Qi (bgQi) VI. genes investigated by CR and Gi (bgGi)
identification of CR functions	significantly regulated genes by CR are calculated for each gene pool with rank product analysis. CR functions are identified with ConsensusPathDB a) Regulated CR genes with bgCR b) Regulated CR genes with bgR c) Regulated CR genes with bgP d) Regulated CR genes with bgQd e) Regulated CR genes with bgQi f) Regulated CR genes with bgGi g) Regulated genes which are identified in a-f, bgCR
<u>Identification of pathways that are affected by both, the CR and the respective plant bioactive treatments</u>	
separate analysis	- all significantly regulated CR genes (gene pool: I-VI or gene pool: II-VI) - all significantly regulated R genes - all significantly regulated P genes - all significantly regulated Qd genes - all significantly regulated Qi genes - all significantly regulated Gi genes
overlapped analysis	- overlapped genes between CR and R; bgR - overlapped genes between CR and P, bgP - overlapped genes between CR and Q, bgQd - overlapped genes between CR and Qi, bgQi - overlapped genes between CR and Gi, bgGi

CR= caloric restriction, R = Resveratrol, P = phloridzin, Qd=quercetin administered by diet, Qi = quercetin administered intraperitoneal, Gi= genistein administered intraperitoneal



overlapped pathways between CR and the respective phytonutrient are identified

2.4.2. Functions of the most strongly regulated CR genes identified by fold change or q-value

Next, to reduce the list of interesting CR pathways, the over-representation analysis is repeated for the CR genes with a fold change ≥ 1.5 . Moreover, PubMed reveals the functions of these genes, and Multiple Experiment Viewer (MeV 4.9.0, <https://sourceforge.net/projects/mev-tm4/files/mev-tm4/MeV%204.9.0/>) identifies and illustrates the overlap in gene expression between CR and the phytonutrients.

Next, the top 50 overlapped genes are identified by q-value for the different interventions. Genes or IDs with unknown functions are excluded. The functions of these genes are revealed using NCBI:gene.

2.4.3. Functions of the genes which are investigated for CR and all dietary phytonutrients

As mentioned above, not all genes are investigated for each phytonutrient. This impedes the identification of the best CR mimetic in our analysis. Therefore, it is decided to repeat the rank product analysis for the investigated genes using all dietary interventions. The phytonutrients administered intraperitoneally are excluded from this analysis.

Next, the overlap in gene expression is determined. Then, the Pearson correlation between the expression changes (mediated by CR) and the phytonutrients is calculated for all genes which are included in the respective CR pathways. The CR pathways are identified using CPDB. This means that all pathways for which at least one of the strongly regulated CR genes ($FC \geq 1.5$) is involved are included, independent of p- or q-value.

2.5. Investigation of genes with special regard for longevity

For a better assessment of whether a phytonutrient is able to mimic the lifespan-expanding effect of CR, it is analysed if the phytonutrients address the same longevity genes as CR. To identify genes associated with longevity, the GenAge database is used (<http://genomics.senescence.info/genes/>) (85). All longevity genes which are regulated by CR in our analysis are discovered. Afterwards, the resulting list is compared with the expression changes triggered by the phytonutrients. A high number of overlapped genes probably indicates a mimicking effect on longevity.

2.6. Analysis of transcription factor binding sites

Finally, it is analysed whether CR and the phytonutrients address the same TFBs. Several methods are investigated, but most are excluded because they are not suitable. The online tool oPOSSUM 3.0 (<http://opossum.cisreg.ca/oPOSSUM3/>) is selected because it is able to identify over-represented TFBs for a list of genes (132; 133). OPOSSUM uses the JASPAR database (100). First, using single site analysis, all significantly regulated CR genes are uploaded to determine TFBs which are overrepresented either in the upregulated genes or in the downregulated genes. Afterwards, the genes overlapping between CR and the phytonutrients are uploaded, and the co-mentioned TFBs are identified. For mice, the results contain the JASPAR collections CORE and PBM. The results can be sorted by z-score or Fischer (F)-score. The z-score compares the rate of a TFB's occurrence in the list of target genes to the expected rate, which is estimated from the background gene list. The F-score (based on one-tailed Fisher exact probability) compares the proportion of co-expressed genes containing a particular TFB to the proportion of the background set that contains the site. For further explanation, see oPOSSUM help (<http://opossum.cisreg.ca/oPOSSUM3/help.html>). In our analysis, all TFBs with either z

> 5 or $F > 5$ are included. The functions of the transcription factors (TF) belonging to the revealed TFBs are identified with the UniProt database (104).

3. Results

3.1. Public Data Collection

As of November 2014, a total of 717 experiments had been mentioned on ArrayExpress along with 174 datasets or 1679 series on GEO using the search items. After reviewing the experiment designs, nine CR and five dietary plant bioactives experiments are included in the analysis (Tables 2 and 3).

As seen in Tables 2 and 3, CR's effect and the phytonutrient treatments on gene expression are analysed with different microarrays. This means that many genes are not analysed by all treatments (Figure 3). For example, the effect of CR is analysed for 46373 Entrez IDs whereas the effect of R is analysed for 18681 Entrez IDs; 17780 IDs are analysed by both interventions. Due to the fact that the p-value and q-value calculations are based on the total number of genes in the gene pool, it is decided to perform the rank product analysis with limited gene pools (see Table 1).




Figure 3: Venn diagram of the shared gene pool between CR and the dietary plant bioactives
(CR = caloric restriction; R = resveratrol; P/Q = phloridzin/quercetin; Qi/Gi = quercetin/genistein administered intraperitoneal)

Figures 4 and 5 indicate that the number of regulated CR genes strongly depends on the used gene pool. Only 285 upregulated and 338 downregulated genes are identified with each gene pool. This is probably based on the large CR gene pool (bgCR) because over 50% of these genes are not analysed by the phytonutrients. Excluding bgCR, 812 upregulated genes and 1106 downregulated genes are identified within each gene pool.

Table 2: Overview of the characteristics of the analyzed caloric restriction studies

	Mouse strain	sex	start age	n	Standard diet	Intervention	dura-tion	Array
CR1 (134)	C57BL6/J 129S1/SvImJ C3H/HeJ CBA/J DBA/2J B6C3F1/J	male	8 wk	n=8	AIN93M 10,2 % fat	25 % Caloric restriction C: 12,7 kcal/day (88,9 kcal/wk) CR: 9,44 kcal/day (66 kcal/wk) Strains (129S1/SvImJ and C57BL6/J), food intake was subsequently decreased to 7,34 kcal/day at 16 weeks of age	14 wks	Affymetrix Mouse Genome 430 2.0 Array
CR2 (135)	B6C57/J	male	27 wk	n=5	AIN-93G 16,4% Fat	Lifelong 40 % Caloric restriction Control: 13,3 kcal per day (93,1 kcal/wk)	to age: 1 year	Illumina MouseRef-8 v2.0 expression beadchip
CR3 (136)	C57Bl/6J	male	8 wk	C n=3 CR n=4	2916 global diet (Harlan) 12 % Fat	40 % Caloric restriction	2 wk	Illumina MouseRef-8 v2.0 expression beadchip
CR4 (137)	C57BL/6J	male	5 wks	n=3	D12450B, Research Diets 10% fat	adaption phase: 1 wk 30% calorie restriction	57 wks	Affymetrix Mouse Genome 430 2.0 Array
CR5 (138)	C57BL/6J	female	15-16 month	n=3	standard chow diet (5053 PicoLab Diet) 13,2 % Fat	ad libitum (106 kcal /wk) vs. Caloric Restriction (80 kcal /wk for 2 weeks, followed by 53 kcal/d for 2 weeks)	4 wk	Affymetrix Mouse Gene 1.0 ST Array [transcript (gene) version]
CR6 (139)	long-lived B6C3F1	male	CRL: 5 mo CRS: 20 mo	n=4	AIN-93M 10,2 % fat	C: 93 kcal/wk or 13,3 kcal/day ad libitum CRS: 77 kcal/wk of CR diet for 2 wk, followed by 52,2 kcal/wk of CR diet for 6 wk.) CRL: 52,2 kcal/wk or 7,5 kcal/d	CRS: 8 wk CRL: 25 mo	Affymetrix Murine Genome U74A Version 2 Array
CR7 (140)	normal mouse (+/+ or +/df)	female	2 mo	C n=7 CR n=8	LabDiet, PMI Feeds, Inc., St Louis, MO	1 wk 10 % CR, than 1 wk 20 % CR, for the remaining time 30 % CR	4 mo	Affymetrix Murine Genome U74A Version 2 Array
CR8 (141)	C57BL/6	male	16 wk	n=12 pooled to 4 groups	2018 Global Rodent diet, Harlan Teklad, U.K. 18% fat	Ad libitum vs. 30 % caloric Restriction	48 h	Affymetrix Mouse Genome 430 2.0 Array
CR9 (142)	C57BL6/J	male	2 mo	n=11	LabDiet 5LG6-JL 12,2% Fat	Ad libitum vs. 40% Caloric Restriction	6 mo	Illumina MouseRef-8 v2.0 expression beadchip

CR = caloric restriction, C = Control, wk = week

Table 3: Overview of the characteristics of the plant bioactive studies

	Mouse strain	sex	age start	n	Standard diet	Intervention	duration	Array
Q1 (143)	C57BL/6	male	6-8 wk	n = 9 pooled in 3 groups	RM3 (E) 801710 Soya-free pow- dered diet from B.S & S; no fur- ther information	Quercetin was added separately to the semi-purified diets at a concen- tration of 6200 ppm (0.62 %; 6.2 g per kg) approximately 7 mg/mouse	adaption phase 2 wks, 4 h food with quercetin	Affymetrix Mouse Expression 430A Array
Q2 (144)	BALB/c	male	6 wk	n = 6	AIN-93G diet fat: 16,4 % (kcal/g)	diet containing 0 % (C), 0,1 %, 0,5 %, or 1,0 % quercetin, ad libitum	2 wk	Affymetrix Mouse Genome 430 2.0 Array
Q3 (145)	n.m.	male	n.m.	n = 2	_____	Quercetin: intraperitoneal (ip) injec- tion in DMSO with 0 mg/kg (C), 500 mg/kg, 1500 mg/kg or 2000 mg/kg	24 h after ad- ministration	Agilent Mouse 60K Genotypic De- signed Array
G (145)						Genistein: intraperitoneal (ip) injec- tion in DMSO with 0 mg/kg (C), 500 mg/kg, 1500 mg/kg or 2000 mg/kg		
R (146)	C57BL/ 6NIA	male	11 mo	C18 n = 4 C27 n = 3 LR n = 5 HR n = 4	AIN-93G diet fat: 16,4 % (kcal/g)	low res (LR): 0.01% resveratrol high res (HR): 0.04% resveratrol	adaption phase 1 mo 7 (C18) or 16 mo	Illumina mouseRef- 8 v1.1 expression beadchip
P (147)	BALB/c	male	6 wk	n = 6 p.g.	AIN-93G diet fat: 16,4 % (kcal/g)	diet containing 0 % (C), 0,1 %, 0,5 %, or 1,0 % phloridzin (purity > 97%)	2 wk	Affymetrix Mouse Genome 430 2.0 Array

Q=quercetin, G= genistein, R = Resveratrol, P = phloridzin, C = control, wk = week, n.m. = not mentioned, mo = month




Figure 4: Venn diagram of the upregulated CR genes identified from rank product analysis with the different gene pools
 (CR = bgCR; R = bgR; P/Q = bgP and bgQd; Qi/Gi = bgQi and bgGi)




Figure 5: Venn diagram for the downregulated CR genes identified from rank product analysis with the different gene pools
 (CR = bgCR; R = bgR; P/Q = bgP and bgQd; Qi/Gi = bgQi and bgGi)

3.2. Comparison of analysis methods used to identify significantly regulated genes/Finding the appropriate p-value test

The number of differently expressed genes varies between the analysis methods used (Table 4; for detailed information, see Appendix A), whereby the rank product analysis shows a larger number than the t-test, the moderated t-test or the Wilcoxon test. The moderated t-test shows more regulated genes than the normal t-test. The Wilcoxon test shows a significant regulation in response to CR but not to the phytonutrient treatments. A combination of the three tests with a fold change ≥ 1.5 additionally reduces the number of regulated genes. No significant up- or downregulation was detected for the phytonutrient treatments when the chi-squared test was applied. This may be explained by the limited number of repetitions per phytonutrient (three to five single experiments, which is too small for the chi-squared test). Therefore, the chi-squared test

was replaced by the 3-fold bigger method, which modestly increased the number of significantly regulated genes.

Table 4: Number of genes that are regulated by caloric restriction or dietary plant bioactives (different analysis methods)

	chi-squared-test ¹	t-test	t-test + FC	moderated t-test	moderated t-test + FC	Wilcoxon test	Wilcoxon test FC	+	rank prod	rank prod + FC
upregulated genes										
CR	yes	26	0	79	1	49	0		3503 ²	464 ²
	no	347	3	571	7	331	0			
R	yes	0	0	0	0	0	0		1304 ³	318 ³
	no	53	10	553	34	0	0			
Qd	yes	0	0	0	0	0	0		826 ³	29 ³
	no	0	0	0	0	0	0			
P	yes	0	0	0	0	0	0		1789 ³	13 ³
	no	277	0	901	0	0	0			
Gi	yes	0	0	0	0	0	0		559 ³	292 ³
	no	0	0	498	3	0	0			
Qi	yes	0	0	0	0	0	0		866 ³	564 ³
	no	0	0	2487	4	0	0			
downregulated genes										
CR	yes	54	0	81	0	64	0		4479 ²	608 ²
	no	460	1	577	2	366	1			
R	yes	0	0	0	0	0	0		1080 ³	228 ³
	no	48	2	530	7	0	0			
Qd	yes	0	0	0	0	0	0		899 ³	4 ³
	no	0	0	0	0	0	0			
P	yes	0	0	0	0	0	0		2922 ³	30 ³
	no	416	0	951	0	0	0			
Gi	yes	0	0	0	0	0	0		917 ³	872 ³
	no	0	0	440	11	0	0			
Qi	yes	0	0	0	0	0	0		1498 ³	1495 ³
	no	0	0	2451	13	0	0			

¹ no chi-squared test means that the '3-fold bigger' method is used; both methods are not used in rank prod analysis

² this number include all Entrez IDs which are significantly up- and downregulated by at least one approach based on the gene pools shown in Table 1

³ results are obtained from rank product analysis with the limited gene pools (Table 1)

FC = fold change, rank prod = rank product method, CR= caloric restriction, R = resveratrol, Qd=quercetin (diet), P = phloridzin, Qi = quercetin (intraperitoneal), Gi= genistein (intraperitoneal)

Based on these results, an overlap between the genes regulated by CR and dietary plant bioactives is calculated (Table 5, Table 6 and Figure 6; overlapped genes; see Appendix B). The results of the 3-fold bigger method are used. The percentage overlap calculation is based on the number of regulated CR genes, which is also investigated using the respective dietary plant bioactive.

Table 5: Total and percental¹ overlap in gene expression between caloric restriction and plant bioactives

	t-test	t-test + FC	moderated t-test	moderated t-test + FC	Wilcoxon test	Wilcoxon test + FC	rank prod ²	rank prod + FC
upregulated genes								
R	2 (0.9%)	0	33 (11%)	0	0	0	275 (29%)	12 (25%)
Qd	0	0	0	0	0	0	158 (7%)	2 (3%)
P	5 (1.1%)	0	15 (2.8%)	0	0	0	172 (8%)	1 (2%)
Gi	0	0	10 (3.0%)	0	0	0	75 (8%)	8 (13%)
Qi	0	0	14 (1.2%)	1 (33%)	0	0	71 (7%)	8 (13%)
downregulated genes								
R	12 (3.7%)	0	41 (11%)	0	0	0	315 (24%)	25 (25%)
Qd	0	0	0	0	0	0	274 (8%)	0 (0%)
P	28 (8.3%)	0	68 (13%)	0	0	0	1106 (34%)	3 (3%)
Gi	0	0	28 (6.8%)	0	0	0	168 (13%)	31 (25%)
Qi	0	0	88 (22%)	0	0	0	255 (19%)	38 (30%)

¹ % of the CR genes, based on number of regulated CR genes which are analyzed by the respective phytonutrient too

² the values of rank product method based on the results of limited gene pool (see Table1)

FC = fold change, rank prod = rank product method, CR= caloric restriction, R = resveratrol, Qd=quercetin (diet), P = phloridzin, Qi = quercetin (intraperitoneal), Gi= genistein (intraperitoneal)




Figure 6: Overlap and opposite regulation of up- and downregulated CR genes in presence of plant bioactives
(Dark bars overlapped regulation, light bars opposite regulation)

Table 6: Total and percental¹ overlap in gene expression regulation between caloric restriction and plant bioactives based on the results of the rank product analysis

total number of genes ¹	Number of genes regulated by CR	Number of genes regulated by phytonutrient	Number of overlapping genes ²
upregulated genes			
R	17780	941	1304
P	27671	2135	1789
Qd	27671	2135	826
Qi	20827	969	866
Gi	20827	969	559
downregulated genes			
R	17780	1328	1080
P	27671	3257	2922
Qd	27671	3257	899
Qi	20827	1337	1498
Gi	20827	1337	917
1 calculation of the percental overlap based on the number of regulated genes by CR 2 only genes from the respective shared gene pool (investigated by both treatments)			
2 overlapping genes are regulated in the same direction by both treatments CR= caloric restriction, R = resveratrol, Qd=quercetin (diet), P = phloridzin, Qi = quercetin (intraperitoneal), Gi= genistein (intraperitoneal)			

In the t-test, only phloridzin (P) and resveratrol (R) show an overlap for up- and downregulated genes, but the overlap disappears after use of the fold change. With the moderated t-test, an overlap is seen for all phytonutrients except for Q administered per diet (Qd), but this disappears again after using the fold change. No overlap is found with the Wilcoxon test, whereas more overlapped genes are identified with the rank product analysis. The biggest overlap is seen for R regarding the upregulated genes and for P and R regarding the downregulated genes. This result is consistent between the moderated t-test and the rank product analysis. After including the fold change, resveratrol is again the only phytonutrient which shows a large overlap regarding the up- and downregulated genes, whereas quercetin administered intraperitoneally (Qi) and genistein administered intraperitoneally (Gi) show a large overlap regarding the downregulated genes.

For a better determination of whether phytonutrients are able to mimic CR-induced gene expression, it is further analysed whether the phytonutrients also show opposite regulations compared to CR. The results presented in Figure 6 indicate as well that R is the phytonutrient which best mimics the influence of CR on gene expression. All other phytonutrients show a large opposite regulation.

3.3. Functional analysis

Next the results of functional analysis are shown. In this analysis only the genes identified by rank product analysis are included, because the t-test, the moderated t-test and

the Wilcoxon test reveal only a small number of overlapped genes. First, the regulated CR genes are examined to identify potential new functions of CR; then, the overlap between CR and the respective plant bioactives is identified.

3.3.1. Identification of over-represented pathways for the regulated CR genes with CPDB

To identify the functions of CR, all genes which are significantly regulated in at least one approach (see Table 1) are analysed with CPDB. In total, 3502 upregulated IDs and 4479 downregulated IDs are included (Appendix C) and 402 significantly affected pathways are mentioned by the up- and downregulated genes in over-representation analysis. Some of them are excluded because the cited functions are not important in murine liver (Appendix E). This leads to 353 pathways which are included in our analysis (Appendix D).

Cholesterol and steroid biosynthesis, bile acid metabolism

Many of the mentioned pathways indicate that CR influences cholesterol metabolism. Most of the involved genes are upregulated by CR, probably indicating that CR increases cholesterol biosynthesis. The epoxysqualene biosynthesis pathway (MouseCyc), trans trans-farnesyl diphosphate biosynthesis pathway (MouseCyc) and mevalonate pathway I (MouseCyc) are mentioned as well. Analysis of the pathways and regulated genes indicate that the steroid biosynthesis in general could be upregulated, whereas the steroid hormone synthesis (e.g., androgen, oestrogen) is mentioned by the downregulated genes.

While the cholesterol biosynthesis is probably increased, the peroxisome proliferator activated receptor (PPAR) signalling pathway (KEGG) indicates that the cholesterol metabolism is reduced. Moreover, CR increases the expression of transporters which excrete cholesterol in the bile canaliculus. In particular, the Abcg5:Abcg8-mediated export of cholesterol and phytosterols (both genes are overexpressed) is upregulated by CR. This complex transports the cytosolic sterols to the gut lumen or in the bile (148). Furthermore, the KEGG bile secretion pathway, which is mentioned by up- and downregulated genes, indicates that CR promotes the transport of cations, bile acid and water into bile canaliculus and upregulates carriers which are responsible for the back resorptions of bile acids. Carriers which transport bile acid into the circulating system are upregulated too. There are also hints that CR promotes LDL and HDL uptake in hepatocytes.

Detoxification

Moreover, the results indicate that CR is involved in the regulation of the metabolism of xenobiotics (e.g., drugs and arsen) and endogenous substances (e.g., fatty acids and

sterols) including phase I (functionalisation of compounds) and phase II reactions (conjugation). The phase I reactions make different xenobiotics more chemically reactive by exposing functional groups. The phase II reactions, especially the glutathione conjugation, cytosolic sulfonation of small molecules and glucuronidation, increase the excretory potential of different compounds. Genes of these pathways are up- and downregulated by CR.

Most regulated genes belong to the cytochrome P450 system, which is involved in phase I reactions. It is not clear which effect CR has on phase I reactions because the cytochrome P450 monooxygenases are mentioned by the up- and downregulated genes. It is interesting that most of the regulated Cyp 450 isoenzymes are of the Cyp2 family. Besides their effects in detoxification, they are also involved in eicosanoid metabolism in humans (149). Furthermore, the results indicate that CR influences the expression of glutathione S-transferases and UDP-glucuronosyltransferases which are involved in phase II reactions. Genes which are involved in cytosolic sulfonation of small molecules (3'-phosphoadenosine 5'-phosphosulfate synthase 2 (Papss2), sulfotransferase family 1E member 1 (Sult1e1), and sulfotransferase family 2A dehydroepiandrosterone (DHEA)-preferring member 3 (Sult2a3)) are upregulated as well. Moreover, CR seems to influence glutathione metabolism in general, including the γ -glutamyl cycle (MouseCyc).

Lipid metabolism

Furthermore, CR seems to regulate the metabolism of lipids and lipoproteins. Several pathways which are part of fatty acid biosynthesis are mentioned by the upregulated genes (e.g. ‘triglyceride biosynthesis’ (Reactome), ‘biosynthesis of unsaturated fatty acids’ (KEGG) and ‘biosynthesis of a long-chain fatty-acyl-CoA’ (KEGG)), but there are always genes that are downregulated, too. Moreover, it is possible that CR is involved in fatty-acid activation. The results of the PPAR signalling pathway (KEGG) indicate as well that CR probably promotes lipogenesis, whereas the beta-oxidation rate is reduced. A few genes of beta-oxidation are upregulated, but most genes are downregulated. Several pathways of beta-oxidation are mentioned for the downregulated genes in over-representation analysis.

Additionally, it is possible that CR has a limiting effect on ketogenesis and lipid transport (‘Lipid digestion, mobilisation, and transport’ – Reactome). Moreover, phospholipid metabolism (including glycerophospholipid biosynthesis and the Kennedy pathway) is shown by both forms of regulation, alpha-linolenic acid and linoleic acid metabolism are mentioned by the downregulated genes, and arachidonic acid metabolism is revealed by the upregulated genes. The last pathway indicates that CR influ-

ences the formation of lecithin, DHETs, HETEs and EETs and seems to regulate prostaglandin and leukotriene metabolism. HETEs can be produced from arachidonic acid and are involved in vasodilation/vasoconstriction and angiogenesis (150). Eicosanoids in general are involved in different biological processes including regulation of insulin sensitivity, stimulation of lipid metabolism, activation of PPAR α and PPAR γ , maintaining of blood pressure (vasodilation/vasoconstriction), anion and proton transport, and inflammation and thermoregulation.

Carbohydrate metabolism/glycolysis/gluconeogenesis/citric acid cycle/respiratory electron chain

Moreover, the pathway analysis indicates that CR is involved in the regulation of glucose metabolism (including glycolysis and gluconeogenesis). One of the upregulated genes is tuberous sclerosis 2 (Tsc2), which is known to interact with Sirt1 and is involved in the inhibition of mTOR signalling (151). Moreover, the results indicate that CR probably increases gluconeogenesis. Furthermore, pyruvate metabolism, the pentose phosphate pathway, the citric acid cycle and the electron transport chain are revealed in pathway analysis. CR upregulates the expression of several genes which are involved in the formation of the different complexes of the electron transport chain (complex I: NADH ubiquinone dehydrogenases like Ndufa5, Ndufa6, Ndufa8, Ndubf6, Ndubf10, Ndufc1, Ndufc2, Ndufs1, Ndufs2, Ndufs8, and Ndufv1; complex II: succinate dehydrogenases like Sdhb, Sdhc, and Sdhd; complex III: ubiquinol-cytochrome c reductases like Uqcrc1, Uqcr10, and Uqcrfs1, complex IV: cytochrome c oxidases like Cox5a, Cox7a1 and complex V: ATP synthases like Atp5a1, Atp5c1, Atp5f1, Atp5g1, Atp5l, Atp5o). Furthermore, the results indicate that CR upregulates Na⁺-independent, facilitative glucose transporters (GLUTs). These are Glut2 (Slc2a2), Glut4 (Slc2a4), Glut5 (Slc2a5) and Glut8 (Slc2a8).

Amino acid and protein metabolism

CR seems to be involved in the metabolism of different amino acids as well. For example, the tryptophan, phenylalanine, tyrosine, serine, glycine, threonine, arginine, glutamate, proline, cysteine, homocysteine, methionine, valine, leucine and isoleucine metabolisms have been discovered. The results indicate that phenylalanine and tyrosine catabolism could be increased by CR, whereas most genes of tyrosine metabolism in general are downregulated. Tyrosine is a substrate for different catecholamines. The 'catecholamine biosynthesis' pathway is shown by the downregulated genes as well. The 'valine, leucine and isoleucine degradation' pathway (KEGG) is mentioned for downregulated genes too. Moreover, CR regulates the expression of several genes which are involved in homocysteine metabolism.

The urea cycle seems to be influenced as well. This hypothesis is supported by the increase of carbamoyl-phosphate synthetase 1 (Cps1) expression, which is a key enzyme of the urea cycle (152). Of particular interest is that the NAD⁺ biosynthesis from tryptophan is mentioned for the upregulated genes, as are nicotinate and nicotinamide metabolism. These regulations could be important for the Sirt1 activity which is regulated by NAD⁺ and NAM (see above). Besides, CR seems to be involved in the apoptotic cleavage of cellular proteins and in ubiquitin mediated proteolysis.

Signalling pathways

Moreover, the pathway analysis indicates that CR regulates several signalling pathways, including growth hormone receptor signalling, insulin signalling, PPAR signalling, leptin receptor signalling, Jak/STAT signalling, ‘androgen receptor signalling pathway’ (Wikipathways), ‘epidermal growth factor receptor (EGFR1) signalling’ (Wikipathways), TNF-α and NF-κB signalling and the transforming growth factor (TGF) beta receptor signalling. Several of these pathways indicate that CR is involved in the regulation of insulin sensitivity and insulin resistance, which has already been mentioned as important for beneficial CR effects (see above). The influence on growth hormone and insulin signalling is an interesting observation as well.

Immune system and inflammation

CR seems to influence inflammation processes and immune function too. Most of these pathways are mentioned by the downregulated genes. These are the ‘antigen processing and presentation’ (MHC I and II) (KEGG), ‘type II interferon signalling (IFNG)’ (Wikipathways), toll like receptor cascade (especially TLR3 and TLR4), complement cascade, coagulation and interleukin signalling (5, 6, 7 and 9). The pathway ‘macrophage markers’ (Wikipathways) is mentioned on over-representation analysis too.

Translation regulation/Regulation of gene expression

Moreover, the pathway analysis indicates that CR is involved in gene expression regulation. In particular, genes that are involved in translation (including elongation, initiation and termination) are upregulated. For example, 3'-UTR mediated translation, cap-dependent translation initiation and activation of gene expression by Srebp are mentioned. Furthermore, CR upregulates genes that code for cytoplasmic ribosomal proteins (large (60S) and small (40S) subunits). They are important for translation too. Moreover, the results indicate that CR is involved in the processing of mRNA (including mRNA transport, splicing and the regulation of mRNA stability by binding of proteins) and post-translational protein modification. Transcription processes are mentioned by the downregulated genes too.

Transporter

Several transporters are regulated by CR. For example, the expression of ABC (ATP-binding cassette) transporters, which are important for the ATP-dependent transport of different compounds against the concentration gradient, is influenced (153). Furthermore, CR seems to be involved in the binding and uptake of ligands by scavenger receptors and the membrane trafficking. This function is important for vesicle-mediated transport, which itself is mentioned by the downregulated genes in over-representation analysis. One of the upregulated receptors is secreted acidic cysteine rich glycoprotein (Sparc). Sparc is involved in liver cell proliferation, migration and extracellular matrix remodelling. The expression of Sparc is increased in fibrotic livers (154). Furthermore, Sparc interacts with AMPK and could be involved in the regulation of Glut4 expression or in regulation of glucose metabolism via AMPK activation (155). Therefore, Sparc upregulation could probably be involved in mediating beneficial CR effects. Moreover, the results indicate that CR is involved in the regulation of metal ion SLC transporters. In particular, transporters that are involved in the zinc influx into cells are downregulated by CR.

Other pathways

Moreover, there are indications that CR is involved in vitamin metabolism. In particular, 'Biotin transport and metabolism' (upregulation) and retinol metabolism (up- and downregulation) are mentioned. In addition, CR upregulates genes which are involved in the circadian clock (Per1, Per2, Cry1 and Cry2). Pathways belonging to circadian regulation are mentioned by the up- and downregulated genes. The cell cycle itself seems to be influenced by CR as well (upregulation; 'G1 to S cell cycle control' – Wikipathways). Moreover, CR seems to downregulate genes of the meiotic recombination. Additionally, the results indicate that CR could be involved in purine and pyrimidine metabolism. For example, the pyrimidine salvage reactions are mentioned in over-representation analysis for upregulated genes.

Heme biosynthesis seems to be influenced as well because the pathways 'heme biosynthesis' (MouseCyc, Wikipathways, Reactome), 'porphyrin and chlorophyll metabolism' (KEGG), 'metabolism of porphyrins' (Reactome) and 'tetrapyrrole biosynthesis II' (MouseCyc) are shown by the downregulated genes. Porphyrins are a special form of tetrapyrroles which can react with metals such as iron to generate different haemoproteins (haemoglobin, cytochromes). Furthermore, the results indicate that CR is involved in phagocytosis. The Wikipathway 'oxidative stress' is also mentioned for the up- and downregulated genes. Moreover, the results indicate that CR is involved in propionate and butanoate metabolism. Whereas the propionate metabolism is mentioned for both up- and downregulated genes, the butanoate metabolism is primarily mentioned for the

downregulated genes of CR. Finally, CR seems to influence methylation processes, (deoxy)ribose phosphate degradation, selenium metabolism and formaldehyde oxidation.

3.3.2. Determining the overlap between CR and the dietary plant bioactives regarding the over-represented pathways

Knowing whether a phytonutrient is a CR mimetic is necessary to determine overlapped functions/pathways besides the transcriptional overlap. Two different approaches are used for this purpose. First, all up- and downregulated genes of the different interventions are investigated independently from each other in over-representation analysis, and co-mentioned pathways are identified (separate analysis). Because this approach does not determine whether CR and the dietary plant bioactive regulate the same genes and mechanisms in the pathway, only the overlapped genes between CR and the respective phytonutrient are analysed in the second approach (overlapped analysis).

3.3.2.1. Separate analysis

For the separate analysis, different methods can be used to identify over-represented pathways by CR. For example, it is feasible to investigate all identified Entrez IDs by the different limited gene pools against the background bgCR (see Table 1, a), which leads to 122 significantly upregulated and 97 significantly downregulated pathways. Moreover, it is possible to take all results of the limited gene pools by themselves and combine them into a list of regulated pathways (see Table 1, a-f). This leads to 211 significantly upregulated and 155 significantly downregulated pathways. Both approaches are used to identify overlapped pathways between CR and the dietary plant bioactives. To identify the over-represented pathways for the phytonutrients, the limited gene pools (bgR, bgP, bgQd, bgQi and bgGi – dependent on the phytonutrient) are used.

Table 7 shows the number of over-represented pathways revealed by the different phytonutrients and the total and percentage overlap between CR and the dietary plant bioactives. The overlap analysis indicates that R is the best mimetic for the upregulated pathways (17% or 11%), and Qi (40% or 33%) and P (28% or 28%) are the best mimetics for downregulated pathways. This coincides with the outcomes of the gene expression analysis. The observation that Qd shows no or only a small overlap with CR is based on the fact that no or only a few over-represented pathways are identified.

Combining the results of all three plant bioactives which are administered per diet shows that no benefit is seen regarding the upregulated genes compared to R alone. For the downregulated genes, a small benefit is seen compared to P alone, which is

caused by the R pathways, whereas Qd results in no further benefit. After including the plant bioactives which are administered intraperitoneally, two further overlapped pathways could be identified regarding the upregulated genes, whereas 18 or 23 further pathways were mimicked by Qi and Gi regarding the downregulated genes. Table 8 shows all overlapped pathways, which are identified in a separate analyses.

The results indicate that the investigated plant bioactives, especially R, are able to mimic the upregulating influence of CR on the biological oxidation (phase I and phase II reactions), the steroid and cholesterol biosynthesis, the fatty acid degradation, the pyruvate metabolism, the pentose and glucoronate interconversions and the binding and uptake of ligands by scavenger receptors. On the other hand, the downregulating effect of CR on the lipid and lipoprotein metabolism (beta oxidation, arachidonic acid metabolism), biological oxidation (phase 1 reactions), steroid and steroid hormone biosynthesis, bile secretion, meiotic recombination, retinol metabolism, protein processing in endoplasmic reticulum, amino acid metabolism (valine, leucine, Isoleucine, tryptophan, cysteine, homocysteine, glycine, serine and threonine), gene expression (including RNA transport, processing and splicing), complement cascade and circadian rhythm seems to be mimicked by phytonutrients as well.

Table 7: Total and percental overlap between the pathways affected by caloric restriction and plant bioactives in separate analyses

	R	P	Q	Qi	Gi	phyto-nutrients per diet*	all phytonutrients
upregulated genes							
number of regulated pathways by the phytonutrients	99	1	0	3	5		
Overlap with CR 122 pathways (total and percental overlap)	21 (17%)	1 (1%)	0 (0%)	3 (3%)	1 (1%)	21 (17%)	23 (19%)
Overlap with CR 211 pathways (summarizing all approaches total and percental overlap)	23 (11%)	1 (1%)	0 (0%)	3 (1%)	1 (1%)	23 (12%)	25 (12%)
downregulated genes							
number of regulated pathways by the phytonutrients	19	83	5	144	38		
Overlap with CR 97 pathways (total and percental overlap)	10 (10%)	27 (28%)	4 (4%)	39 (40%)	22 (23%)	31 (32%)	49 (51%)
Overlap with CR 155 pathways (summarizing all approaches total and percental overlap)	10 (7%)	43 (28%)	5 (3%)	51 (33%)	27 (17%)	47 (30%)	70 (45%)

* Pathways of the experiments with intraperitoneal administration are not considered

Table 8: Overlapping pathways between CR and plant bioactives identified in separate analysis

overlapped upregulated pathways	overlapped downregulated pathways
Metabolism – Reactome ¹	Metabolism – Reactome ^{1,2,4,5}
Biological oxidations – Reactome ^{1,4,5}	Meiotic Recombination – Reactome ¹
Cholesterol Biosynthesis – Wikipathways ^{1,2}	Tryptophan metabolism – Wikipathways ^{1,2}
Cholesterol biosynthesis – Reactome ¹	Metabolism of lipids and lipoproteins – Reactome ^{1,4,5}
Metabolism of lipids and lipoproteins – Reactome ¹	Sulfur amino acid metabolism – Reactome ^{1,4}
Drug metabolism - cytochrome P450 - Mus musculus (mouse) – KEGG ¹	Steroid hormone biosynthesis - Mus musculus (mouse) – KEGG ^{1,2,4}
Pyruvate metabolism - Mus musculus (mouse) – KEGG ¹	Drug metabolism - other enzymes - Mus musculus (mouse) – KEGG ^{1,2,4,5}
Phase 1 - Functionalization of compounds – Reactome ¹	Protein processing in endoplasmic reticulum - Mus musculus (mouse) – KEGG ^{1,2,4,5}
Cytochrome P450 - arranged by substrate type – Reactome ¹	Fatty acid, triacylglycerol, and ketone body metabolism – Reactome ^{1,2}
cholesterol biosynthesis III (via desmosterol) – MouseCyc ¹	Systemic lupus erythematosus - Mus musculus (mouse) – KEGG ^{1,4}
Metapathway biotransformation – Wikipathways ¹	Xenobiotics – Reactome ^{2,3,4,5}
superpathway of cholesterol biosynthesis – MouseCyc ¹	nicotine degradation III – MouseCyc ^{2,4,5}
Terpenoid backbone biosynthesis - Mus musculus (mouse) – KEGG ¹	Valine, leucine and isoleucine degradation - Mus musculus (mouse) – KEGG ^{2,4,5}
cholesterol biosynthesis II (via 24,25-dihydrolanosterol) – MouseCyc ¹	Retinol metabolism - Mus musculus (mouse) – KEGG ^{2,3,4,5}
Pentose and glucuronate interconversions - Mus musculus (mouse) – KEGG ¹	Exercise-induced Circadian Regulation – Wikipathways ^{2,4}
Phase II conjugation – Reactome ¹	mRNA processing – Wikipathways ²
Steroid biosynthesis - Mus musculus (mouse) – KEGG ¹	nicotine degradation II – MouseCyc ^{2,4,5}
mevalonate pathway I – MouseCyc ¹	Gene Expression – Reactome ^{2,4}
cholesterol biosynthesis I – MouseCyc ¹	bupropion degradation – MouseCyc ^{2,4,5}
Fatty acid degradation - Mus musculus (mouse) – KEGG ¹	Butanoate metabolism - Mus musculus (mouse) – KEGG ^{2,4}
Propanoate metabolism - Mus musculus (mouse) – KEGG ¹	Diurnally Regulated Genes with Circadian Orthologs – Wikipathways ^{2,4}
Diurnally Regulated Genes with Circadian Orthologs – Wikipathways ¹	Synthesis of epoxy (EET) and dihydroxyeicosatrienoic acids (DHET) – Reactome ²
Binding and Uptake of Ligands by Scavenger Receptors – Reactome ¹	Drug metabolism - cytochrome P450 - Mus musculus (mouse) – KEGG ^{2,4,5}
Chemical carcinogenesis - Mus musculus (mouse) – KEGG ⁴	Processing of Capped Intron-Containing Pre-mRNA – Reactome ^{2,4}
Leptin Insulin Overlap – Wikipathways ⁴	mRNA Processing – Reactome ^{2,4}
	mRNA Splicing – Reactome ^{2,4}
	Spliceosome - Mus musculus (mouse) – KEGG ²
	mRNA Splicing - Major Pathway – Reactome ^{2,4}
	Cytochrome P450 - arranged by substrate type – Reactome ^{2,3,4,5}
	Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) – Reactome ^{2,5}
	Chemical carcinogenesis - Mus musculus (mouse) – KEGG ^{2,4,5}
	Metabolism of xenobiotics by cytochrome P450 - Mus musculus (mouse) – KEGG ^{2,4,5}
	mRNA 3,-end processing – Reactome ²
	Post-Elongation Processing of Intron-Containing pre-mRNA – Reactome ²
	fatty acid β-oxidation I – MouseCyc ²

Tab 8/b: Overlapped pathways between CR and the plant bioactives identified in separate analysis

overlapped upregulated pathways	overlapped downregulated pathways
	Phase 1 - Functionalization of compounds – Reactome ^{2,3,4,5}
	RNA transport - Mus musculus (mouse) – KEGG ²
	Tryptophan metabolism - Mus musculus– KEGG ²
	2-methylbutyrate biosynthesis – MouseCyc ²
	Fatty acid degradation - Mus musculus – KEGG ^{2,4,5}
	Toll-Like Receptors Cascades – Reactome ²
	Mitochondrial Fatty Acid Beta-Oxidation – Reactome ²
	Biological oxidations – Reactome ^{2,3,4,5}
	Fatty Acid Beta Oxidation – Wikipathways ²
	Propanoate metabolism - Mus musculus – KEGG ^{2,4}
	Bile secretion - Mus musculus (mouse) – KEGG ²
	fatty acid β-oxidation IV (unsaturated, even number) – MouseCyc ²
	Complement cascade – Reactome ^{4,5}
	Regulation of Complement cascade – Reactome ^{4,5}
	Complement and coagulation cascades - Mus musculus (mouse) – KEGG ^{4,5}
	Bile acid and bile salt metabolism – Reactome ⁴
	Synthesis of bile acids and bile salts – Reactome ⁴
	Complement Activation, Classical Pathway - Wikipathways ^{4,5}
	Synthesis of bile acids and bile salts via 7alpha-hydroxycholesterol – Reactome ⁴
	Lipoprotein metabolism – Reactome ^{4,5}
	Retinoid metabolism and transport – Reactome ⁴
	Beta-oxidation of pristanoyl-CoA – Reactome ⁴
	Primary bile acid biosynthesis - Mus musculus (mouse) – KEGG ⁴
	Lipid digestion, mobilization, and transport – Reactome ^{4,5}
	Synthesis of bile acids and bile salts via 24-hydroxycholesterol – Reactome ⁴
	Porphyrin and chlorophyll metabolism - Mus musculus (mouse) – KEGG ⁴
	Initial triggering of complement – Reactome ^{4,5}
	Peroxisomal lipid metabolism – Reactome ⁴
	Prion diseases - Mus musculus (mouse) – KEGG ^{4,5}
	Degradation of cysteine and homocysteine – Reactome ⁵
	Steroid Biosynthesis – Wikipathways ⁴
	Statin Pathway – Wikipathways ⁴
	superpathway of cholesterol biosynthesis – MouseCyc ⁴
	Synthesis of bile acids and bile salts via 27-hydroxycholesterol – Reactome ⁴
	Glycine, serine and threonine metabolism - Mus musculus (mouse) – KEGG ⁵

¹ overlap between caloric restriction (CR) and resveratrol, ² overlap between CR and phloridzin, ³ overlap between CR and quercetin administered per diet, ⁴ overlap between CR and quercetin administered intraperitoneal, ⁵ overlap between CR and genistein administered intraperitoneal

3.3.2.2. Overlapped analysis

As mentioned above, the separate analysis does not answer the question of whether the phytonutrients address the same genes as CR in the pathways. Different gene expression changes can result in different effects in the pathway. Therefore, only the overlapped genes between CR and the respective dietary plant bioactives are next analysed with ConsensusPathDB. The overlapping genes of CR and the respective plant bioactive and the identified pathways ($q < 0.05$) are presented in Appendix F to O.

Resveratrol

A total of 275 upregulated genes and 315 downregulated genes are regulated by both CR and R. The upregulated genes are over-represented in 44 pathways, whereas the downregulated genes are over-represented in 17 pathways. They are summarised in Table 9. The results indicate that R could be able to mimic many CR functions, such as steroid and cholesterol metabolism, biological oxidation, the circadian clock, PPAR signalling, and nicotinate and nicotinamide metabolism. As mentioned above, some of them could be important for the beneficial effects of CR, including the detoxification system, the circadian clock and the nicotinamide metabolism. Table 10 shows the overlapped genes between CR and R for the pathways identified in separate and overlapped analysis. Interesting is that both interventions upregulate the expression of Cry2, Per1, Per2 and Per3, indicating an influence on circadian regulation.

Moreover, nicotinamide N-methyltransferase (Nnmt) is upregulated by CR and R. SCHMEISSER et al. (2013) indicate that Nnmt expression is important for the Sirt1-mediated effect on lifespan in *C. elegans* and that Nnmt can prolong lifespan independent of Sirt1 (35). Moreover, the production of ROS by aldehyde oxidases (Aox) is necessary for the lifespan-expanding effect of Nnmt in *C. elegans*. Our analysis indicates that CR and R downregulate the expression of Aox1 and Aox3 in murine liver. However, Nnmt may be able to increase ROS production independent of Aox, because many stress defence genes like Gstm5, Gstm7 are Gsto1 are upregulated in our analysis. Figure 7 gives an overview of the mechanisms affected by Nnmt.

Nnmt methylates NAM using S-adenosylmethionine (SAM), which leads to the formation of N1-methylnicotinamide (MNAM). NAM is known to possibly prevent the lifespan-expanding effects of CR by increasing the level of NADH, which is known to be an inhibitor of Sir2 (Sirt1) in yeast (156). MNAM, on the other hand, is known for its antithrombotic and anti-inflammatory effects and for mediating Foxo- or Nrf2-dependent increased expression of genes which are involved in detoxification and stress response

(35; 157). Thus, Nnmt could be involved in the Sirt1- and Foxo1-dependent increase of lifespan in mice.

Table 9: Summarized results of pathway analysis of the overlapping genes of CR and resveratrol

upregulated pathways	downregulated pathways
Cholesterol Biosynthesis, mevalonate pathway, Terpenoid backbone biosynthesis, steroid biosynthesis	Steroid hormone biosynthesis - <i>Mus musculus</i> (mouse) - KEGG
Metabolism of lipids and lipoproteins - Reactome	Metabolism of lipids and lipoproteins; Fatty acid, triacylglycerol, and ketone body metabolism;
Retinol metabolism	Retinol metabolism
Biological oxidations – Reactome, phase I (cytochrome P40s) and phase II reactions (Glucuronidation, Glutathione conjugation)	Biological oxidations
Pentose and glucuronate interconversions	Protein processing in endoplasmic reticulum
PPAR signaling pathway	glycerol-3-phosphate shuttle / Glycolysis
Metapathway biotransformation	Sulfur amino acid metabolism; L-cysteine degradation
Circadian clock	Drug metabolism
Glutathione metabolism	Taurine and hypotaurine metabolism
Trafficking of dietary sterols, bile secretion	Meiotic Recombination
Porphyrin and chlorophyll metabolism	glutathione biosynthesis
Pyruvate metabolism	
Starch and sucrose metabolism	
salvage pathways of pyrimidine ribonucleotides	
Nicotinate and nicotinamide metabolism	

Table 10: Genes that are regulated by caloric restriction and resveratrol in selected pathways

Pathway	upregulated genes	downregulated genes
biological oxidation	Acss2, Cyp2b9, Cyp2b13, Cyp3a13, Cyp4a10, Cyp4a14, Cyp7a1, Cyp17a1, Cyp51, Fmo3, Gstm5, Gstm7, Gst01, Gstt2, Nnmt, Papss2, Sult1a1, Sult1c2, Ugp2, Ugt1a2	Aldh1a1, Cyp2d9, Cyp2f2, Cyp4a12a, Cyp4a12b, Cyp7b1, Gclc, Gclm, Slc35d1, Sult2a8, Ugt2b1, Ugt2b34, Ugt2b35
Circadian regulation	Bhlhe40, Cebpb, Cry2, Gfra1, Gstm5, Idi1, Nr1d1, Per1, Per2, Per3, Prkab1, Stbd1, Ugp2	G0s2, Ppp1r3c, Tab2
PPAR signaling pathway	Acox1, Angptl4, Cyp4a10, Cyp4a14, Cyp7a1, Fabp5, Pck1, Ppara	Acaa1b, Cpt2, Cyp4a12a, Cyp4a12b, Rxra
Nicotinate and nicotinamide metabolism	Enpp1, Nadsyn1, Nmrk1, Nnmt	Aox1, Aox3
steroid and cholesterol biosynthesis	Cyp17a1, Cyp51, Dhcr24, Dhcr7, Fdps, Hmgcr, Hmgcs1, Idi1, Lss, Mvk, Nsdhl, Pmvk, Sqle	Hsd3b2, Hsd3b5, Lipa
bile secretion; Trafficking of dietary sterols	Abcc4; Abcg5, Abcg8, Atp1b1, Cyp7a1, Hmgcr, Slco1a4	Abcb4, Aqp8, Baat, Slc22a7, Rxra

That Nnmt is upregulated by CR has already been observed by ESTEP et al. (2009) (156). Whereas previous studies indicate that a high Nnmt expression leads to unwanted effects in adipocytes and is associated with obesity, diabetes, cancer and neu-

rodegenerative diseases, HONG et al. (2015) show that the effects of Nnmt in liver are wanted (158; 159; 160). These effects seem to be mediated by Nnmt-dependent stabilising of Sirt1 (159). Moreover, in vitro analyses indicate that Nnmt increases the expression of Sirt1, Sirt2 and Sirt3 in human neuroblastoma cells and that Sirt3 is a key mediator for enhancing the Nnmt-dependent increase in complex I activity and ATP synthesis (161). Furthermore, there are indications that Nnmt is involved in the regulation of gluconeogenesis and lipogenesis (160; 159).




Figure 7: Overview of the potential mechanism mediated by Nnmt

Moreover, R also upregulates the complex Abcg5:Abcg8, which indicates a mimicking effect on cholesterol efflux via bile acid. The upregulation of 24-dehydrocholesterol reductase (Dhcr24) seems to be important as well. Besides its effects in cholesterol biosynthesis, Dhcr24 can directly scavenge ROS, protecting cells from oxidative stress-induced apoptosis (162). Another interesting overlap is the upregulation of PPAR α . It is known that PPAR α is activated during fasting, which stimulates fatty acid oxidation, ketogenesis and gluconeogenesis through transactivation of target genes (57). Moreover, PPAR α is involved in bile acid synthesis and in modulating inflammatory responses.

Phloridzin

There were 158 upregulated and 1106 downregulated genes that overlapped between P and CR. The 19 pathways over-represented by upregulated genes and the 46 pathways over-represented by downregulated genes are summarised in Table 11. They

indicate that phloridzin could be able to mimic some effects of CR on cholesterol metabolism, circadian clock, biological oxidation and PPAR signalling as well, but the genes mentioned Table 12 indicate that P operates somewhat differently from R. Moreover, it is possible that phloridzin mimics some effects of CR on amino acid metabolism, the complement cascade, mRNA processing and fatty acid β -oxidation. It is particularly interesting that P mimics the downregulating effect of CR on insulin receptor substrate 1 (Irs1) and growth hormone receptor (Ghr). This indicates that P could be able to inhibit GH/Igf1 signalling as well. Moreover, Nnmt expression is also increased by P.

Table 11: Summarized results of pathway analysis of the overlapping genes of CR and phloridzin

upregulated pathways	downregulated pathways
Cholesterol Biosynthesis; Terpenoid backbone biosynthesis	mRNA processing; Regulation of mRNA Stability by Proteins that Bind AU-rich Elements
salvage pathways of pyrimidine ribonucleotides	nicotine degradation
Circadian Clock	Toll-Like Receptors Cascades
Metabolism of lipids and lipoproteins; Fatty Acid Biosynthesis	Valine, leucine and isoleucine degradation - Mus musculus (mouse)
Phenylalanine, tyrosine and tryptophan biosynthesis - Mus musculus (mouse)	Protein processing in endoplasmic reticulum - Mus musculus (mouse)
Regulation of mRNA Stability by Proteins that Bind AU-rich Elements	Synthesis of epoxy (EET) and dihydroxyeicosatrienoic acids (DHET) - Reactome
(deoxy)ribose phosphate degradation	Drug metabolism
	Retinol metabolism - Mus musculus (mouse)
	bupropion degradation
	Butanoate metabolism - Mus musculus (mouse)
	Circadian Clock
	Biological oxidation – Phase I (Cytochrome P 450s)
	Porphyrin metabolism
	Type II interferon signaling (IFNG)
	2-methylbutyrate biosynthesis
	fatty acid β -oxidation; Fatty acid degradation
	PPAR signaling pathway
	Complement Activation, Classical Pathway – Wikopathways

Quercetin (diet)

There are 158 upregulated and 274 downregulated overlapped genes between CR and Qd. The upregulated genes are over-represented in nine pathways, whereas the downregulated genes are over-represented in one pathway (summarised in Table 13). These pathways indicate that Qd could be able to mimic the effect of CR on biological oxidation and the circadian clock. The expression of Nnmt is increased for Qd as well (Table 14). Moreover, the downregulation of choline kinase alpha (Chka) indicates that Qd could be able to inhibit Akt signalling (163; 164). Therefore, it is possible that Qd

inhibits mTOR signalling and increases the activity of Foxos even if relevant gene expression changes are not shown in Table 14.

Table 12: Genes that are regulated by caloric restriction and phloridzin in selected pathways

Pathway	upregulated genes	downregulated genes
Growth hormone receptor signaling	Cish, Soc52	Adam17, Ghr, Irs1, Stat5a
circadian clock	Idi1, Per1, Tubb4a	Azin1, Clock, Dnaja1, G0s2, Klf9, Nr1d1, Ppp1r3c, Psma4, Stbd1, Tab2
Bile secretion		Abcb4, Abcc2, Abcc3, Abcg2, Aqp4, Aqp9, Ephx1, Ldlr, Slc22a1, Slc22a7, Slco1a1, Slco1b2
Cholesterol biosynthesis	Dhcr7, Fdps, Idi1, Nsdhl, Pmvk, Sqle	Acat1, Nudt12, Sc5d
PPAR signaling pathway	Cyp4a14, Fabp5, Scd1	Acaa1b, Acadm, Acox2, Cd36, Cpt1a, Ehhadh, Fabp2, Fabp7, Gyk, Ppara, Sorbs1
biological oxidation	Acss2, Cyp2c39, Cyp4a14, Nnmt	Adh4, Cyp2a12, Cyp2c37, Cyp2c50, Cyp2c54, Cyp2d12, Cyp2d40, Cyp2j5, Cyp2j6, Cyp3a13, Cyp3a25, Gclc, Gclm, Sult1c2, Tpm1, Ephx1, Ugt2b34, Ugt2b35

Table 13: Summarized results of pathway analysis of the overlapping genes of CR and dietary quercetin

upregulated pathways	downregulated pathways
Biological oxidations – Phase I (cytochrome P450s)	Destabilization of mRNA by AUF1 (hnRNP D0)
Macrophage markers	
Circadian Clock	
Retinol metabolism	
Hypertrophy Model	

Quercetin (administered intraperitoneally)

The overlapped analysis included 71 upregulated and 255 downregulated genes overlapping between Qi and CR. Sixty over-represented pathways could be identified for the upregulated genes, whereas 93 could be identified for the downregulated genes (summarised in Table 16). These pathways indicate that Qi could be able to mimic some effects on biological oxidation, lipid metabolism, growth hormone and insulin signalling, bile metabolism, glucose metabolism, PPAR signalling and the circadian clock. The genes, presented in Table 15, indicate that Qi seems to influence GH/Igf1 signalling as well because Ghr is downregulated by Qi too.

Table 14: Genes that are regulated by caloric restriction and dietary quercetin in selected pathways

Pathway	upregulated genes	downregulated genes
Kennedy pathway		Chka, Sgpl1
Circadian clock	Cebpb, Per1	G0s2, Nr1d1, Ppp1r3c
biological oxidation	Cyp2b10, Cyp4a14, Nnmt, Sult1e1	Cyp2d12, Cyp2d40, Slc26a1, Sult1c2

Table 15: Genes that are regulated by caloric restriction and quercetin administered intraperitoneally in selected age-related pathways

pathway	upregulated genes	downregulated genes
Methylation	Inmt, Nnmt	
Growth hormone receptor signaling	Cish, Irs2, Socs2	Ghr, Socs3, Stat5a
Adipocytokine signaling pathway, insulin signaling pathway, Leptin Insulin Overlap	Irs2, Pklr, Lepr, Slc2a4, Socs2	Cd36, Hras, Ppp1ca, Ppp1cb, Rheb, Socs3, Stat5a
Regulation of Insulin-like Growth Factor (IGF) Transport and Uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)	Igfbp1	Igfals

Moreover, CR and Qi increases the expression of insulin-like growth factor binding protein 1 (Igfbp1) and decreases the expression of insulin-like growth factor binding protein, acid labile subunit (Igfals). Then Igfbp1 binds Igf1, which results in decreased Igf1 activity (165). It is already seen that Igfbp1 expression is increased in humans after CR and in rats with a protein- and calorie-restricted diet (166; 167; 168; 169; 170). GAN et al. (2005) could further show that Sirt1 increases Igfbp1 expression by Foxo-dependent and -independent mechanisms (171). In addition, Igfbp1 mediates Igf1-dependent and independent effects on glucose metabolism, postnatal growth, and cell regeneration and survival (172), while Igfals is important for Igf1 stability (173). Thus, decreased expression of Igfals leads to reduced Igf1 levels without affecting Igf1 expression. XU and SONNTAG (1996) have already shown that CR decreases plasma Igf1 level compared to ad libitum fed mice (21). Furthermore, Qi increases the expression of Nnmt. This indicates that Qi could be able to mimic the effects of CR on NAM and NAD+ metabolism.

Genistein (administered intraperitoneally)

The analysis showed that 71 upregulated and 255 downregulated genes overlapped between Gi and CR. The upregulated genes were over-represented in 39 pathways, whereas the downregulated genes were over-represented in 57 pathways (summarised in Table 17). These pathways indicate that Gi could be able to mimic some effects of CR on biological oxidation, lipid and glucose metabolism, GH signalling, PPAR signalling, immune system and bile metabolism.

3.3.3. Analysis of the top genes up- and downregulated by CR identified with fold change (> 1.5)

As seen above, many pathways are revealed by over-representation analysis. To investigate the overlap between CR and the more purposeful phytonutrients, only the CR IDs with a fold change > 1.5 and a known gene symbol are analysed next. This leads to 81 upregulated genes and 148 downregulated genes (Appendix P). Already known

genes which are important for mediating CR effects (like sirtuins and Foxos) are not included in this list.

Table 16: Summarized results of pathway analysis of the overlapping genes of CR and quercetin administered intraperitoneally

upregulated pathways	downregulated pathways
Biological oxidations – Phase I (Cytochrome P450s) and phase II reactions (glutathione-mediated detoxification; Cytosolic sulfonation of small molecules)	Biological oxidations – Phase I (Cytochrome P450s)
Retinol metabolism	Retinol metabolism
Metabolism of lipids and lipoproteins; arachidonic acid metabolism, biosynthesis of unsaturated fatty acids; alpha-linolenic (omega3) and linoleic (omega6) acid metabolism; Fatty acid degradation; Fatty Acyl-CoA Biosynthesis; Fatty acid elongation; Triglyceride Biosynthesis	arachidonic acid metabolism; Synthesis of Prostaglandins (PG) and Thromboxanes (TX); Fatty acid degradation (Beta-oxidation; omega-oxidation); Peroxisomal lipid metabolism; sphingosine and sphingosine-1-phosphate metabolism; Lipoprotein metabolism; Fat digestion and absorption; Chylomicron-mediated lipid transport; Linoleic acid metabolism
Growth hormone receptor signaling	Growth hormone receptor signaling
Metabolism of carbohydrates; Hexose / Glucose transport; Glycolysis and Gluconeogenesis, especially glycolysis	Bile acid and bile salt metabolism; Synthesis and recycling of bile acids and bile salts, Bile salt and organic anion SLC transporters
Estrogen metabolism; biosynthesis of estrogens	Steroid hormone biosynthesis, Steroid biosynthesis
PPAR signaling pathway	PPAR signaling pathway
Adipocytokine signaling pathway	Peroxisome
Insulin signaling pathway; Regulation of Insulin-like Growth Factor (IGF) Transport and Uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)	Immune System, especially innate immune system; Cytokine Signaling; Interleukin-6 signaling; Complement and coagulation cascades; Interferon Signaling; Antigen processing and presentation
Metapathway biotransformation	heme biosynthesis, porphyrin metabolism
Methylation - Wikipathways	bupropion degradation - MouseCyc
Signaling by Leptin	Synthesis and degradation of ketone bodies
Glutathione metabolism	mevalonate pathway; Terpenoid backbone biosynthesis
Steroid hormone biosynthesis	Protein processing in endoplasmic reticulum
Selenium - Wikipathways	Androgen biosynthesis
Prolactin signaling pathway	Phagosome
bupropion degradation	Tryptophan metabolism
EPO Receptor Signaling	Proteasome
	Statin Pathway
	Circadian Clock
	One carbon metabolism and related pathways
	Valine, leucine and isoleucine degradation

Nonetheless, CR does upregulate Sirt1, Ppargc1a (codes for PGC-1α), Foxo1 and Foxo3, but with a smaller fold change. Qd is the only phytonutrient which shows an

upregulation for Sirt1 too. This is really surprising because previous experiments have indicated that resveratrol is a Sirt1 activator (80; 174). Probably these missing upregulations could be explained by study design or mouse strain. P shows a significant downregulation in Sirt1 expression. The expression of Foxo1, Foxo3 and Ppargc1a is not influenced by phytonutrients.

Table 17: Summarized results of pathway analysis of the overlapping genes of CR and genistein administered intraperitoneally

upregulated pathways	downregulated pathways
Biological oxidations – Phase I (Cytochrome P450s) and phase II reactions (Cytosolic sulfonation of small molecules; glutathione-mediated detoxification)	Biological oxidations – Phase I (Cytochrome P450s)
Retinol metabolism	Retinol metabolism
Metabolism of lipids and lipoproteins; Fatty acid, triacylglycerol, and ketone body metabolism; Arachidonic acid metabolism; Fatty acid elongation; Biosynthesis of unsaturated fatty acids; Triglyceride Biosynthesis; Fatty Acyl-CoA Biosynthesis; acyl-CoA hydrolysis	Metabolism of lipids and lipoproteins, arachidonic acid metabolism; Fatty acid degradation; Beta-oxidation; Synthesis and degradation of ketone bodies; Lipoprotein metabolism; Lipid digestion, mobilization, and transport
Growth hormone receptor signaling	Complement and Coagulation Cascades
Phenylalanine, tyrosine and tryptophan biosynthesis; Phenylalanine metabolism	Bile acid and bile salt metabolism, Primary bile acid biosynthesis; bile secretion; Bile salt and organic anion SLC transporters
Glycolysis and Gluconeogenesis, especially gluconeogenesis	Transport of glucose and other sugars, bile salts and organic acids, metal ions and amine compounds
bupropion degradation	heme biosynthesis; Porphyrin metabolism
Estrogen metabolism; biosynthesis of estrogens	cholesterol biosynthesis; Terpenoid backbone biosynthesis
Metapathway biotransformation	Destabilization of mRNA by AUF1 (hnRNP D0)
Leptin Insulin Overlap	Protein processing in endoplasmic reticulum
Prolactin signaling pathway	Abacavir transport and metabolism
EPO Receptor Signaling	SLC-mediated transmembrane transport;
Cysteine and methionine metabolism	Organic cation/anion/zwitterion transport
IL-3 Signaling Pathway	Interferon signaling
	PPAR signaling pathway
	Peroxisome
	Proteasome

For the genes with FC > 1.5, over-representation analysis with CPDB is repeated (Appendix Q). Table 18 summarises the results of this analysis. Moreover, it is analysed whether these genes are already mentioned in the context of CR, ageing, lifespan or longevity. Nnmt and Igfbp1 are strongly upregulated by CR in our analysis. As mentioned above, Igfbp1 is important for Igf1 activity, and Nnmt increases the stability of Sirt1 and is necessary for Sirt1-dependent longevity in *C. elegans*. Figure 8 shows the

expression changes of the genes which are strongly regulated by CR and which are already mentioned in the context of CR or of known CR mechanisms.

More genes indicate that CR is involved in the regulation of Igf1 signalling and insulin sensitivity. For example, CR increases the expression of Glut4 (also known as Slc2a4). Glut4 is associated with increased insulin sensitivity and increased glucose uptake (158; 17). For this, Glut4 must translocate to the cell membrane. This is mediated either by Akt or by AMPK (48). That CR increases Glut4 expression has already been seen in muscle (11). The upregulation of FK506 binding protein 5 (Fkbp5) could be important for the CR effects too because it is seen in tumour cells that Fkbp5 inhibits the activity of Akt by phosphorylation (175). Thus, the function of Fkbp5 in murine liver and its role in Igf1 signalling must be further investigated. The downregulation of phosphatidylcholine transfer protein (Pctp) could be interesting as well because genetic ablation or chemical inhibition of Pctp is associated with increased insulin sensitivity by promoting Irs2 activity (176). The expression of Irs2 is increased by CR in our analysis, but the fold change is smaller than 1.5. Interferon gamma induced GTPase (Igtp) is downregulated by CR in our analysis. There are indications that Igtp expression is necessary for PI3K/Akt signalling, which suggests that the reduced PI3K/Akt signalling mentioned in relation to CR is probably at least partially mediated by Igtp (177).

Table 18: Summarized results of pathway analysis of the top regulated genes of caloric restriction

upregulated pathways	downregulated pathways
Biological oxidations (Phase 1, Phase 2)	Biological oxidations (Phase 1)
Fatty acid and triglyceride biosynthesis, arachidonic acid metabolism, beta oxidation	Metabolism of lipids and lipoproteins, arachidonic acid metabolism, fatty acid degradation
Retinol metabolism	Retinol metabolism
Amino acid synthesis and interconversion (transamination), Amino acid transport across the plasma membrane, Alanine, aspartate and glutamate metabolism, Amino acid and oligopeptide SLC transporters,	Metabolism of amino acids and derivatives, tyrosine metabolism, sulfur amino acid metabolism, Glycine, serine and threonine metabolism
Glucose metabolism, Pyruvate metabolism, Glycolysis and Gluconeogenesis	Steroid hormone and steroid biosynthesis, androgen biosynthesis, glycoprotein hormones, Peptide hormone biosynthesis
Glutathione metabolism	Meiotic Recombination
pentose phosphate pathway	Immune system, complement and coagulation cascades, Interferon Signaling
Insulin signaling pathway	catecholamine biosynthesis
Cholesterol biosynthesis	Bile acid and bile salt metabolism, Synthesis of bile acids and bile salts
Biotin transport and metabolism	NFKB activation
	PPAR signaling pathway
	Binding and Uptake of Ligands by Scavenger Receptors




Figure 8: Heat map showing the fold change of selected genes regulated by caloric restriction and plant bioactives

Flavin containing monooxygenase 3 (Fmo3) is the most strongly upregulated gene in our analysis. A strong upregulation is also seen before by FU and KLAASSEN (2014) and in the meta-analysis from SWINDELL (2009) (10; 178). Moreover, Fmo3 expression and activity changes during ageing indicating that CR could be able to prevent these changes (179; 180). Fmos are part of phase I reactions in biological oxidation, catalyse the oxidation of endobiotics and xenobiotics and are involved in the regulation of glucose and lipid homeostasis (181; 10; 57). As mentioned above, the upregulation of genes of the xenobiotic metabolism could be important for the retardation of ageing, and studies show that Fmo3 expression, in particular, is regulated by insulin and GH/Igf1 signalling (182; 58). Moreover, Fmo3 is necessary for Foxo1 expression, which is known to be important for CR effects as well (183). These explanations indicate that Fmo3 could be important for mediating beneficial CR effects. The expression of Fmo3 is upregulated by all phytonutrients, but the strongest upregulation is seen for R, Qi and Gi.

Other upregulation genes involved in biological oxidation include for example glutathione S-transferase alpha 4 (Gsta4), glutathione S-transferase mu 3 (Gstm3), Sult1e1 and Sult2a3. That CR upregulates several genes of phase II reactions, including Gsta4, Gstm3 and Sult1e1, has already been observed (10; 184). It is proposed that besides its effects in biological oxidation, Sult1e1 could be involved in the regulation of GH/Igf1 signalling. LI et al. (2009) have shown that Sult1e1 inhibits GH-stimulated STAT5b phosphorylation and Igf1 synthesis in transfected HepG2 cells (185). Gsta4 and Gstm3 expressions are associated with reduced cellular oxidative damage, which is important for beneficial CR effects (10; 186). It is seen in *C. elegans* that increased expression of Gsta4 is associated with stress resistance and lifespan (187). The fact that CR reduces oxidative damage is confirmed by the downregulation vanin 1 (Vnn1). Vnn1 is a marker for oxidative stress (188). Alcohol dehydrogenase 4 (class II), pi polypeptide (Adh4) is linked to lifespan and oxidative stress as well. Lifespan is increased in Adh4 knockout worms, possibly mediated by increased ROS production (189). This supports the hormesis theory of ageing, which says that CR causes an increased lifespan by promoting mild stress. To our knowledge, no study is available that has investigated the impact of Adh4 on lifespan in mice. Therefore, it should be analysed whether Adh4 is involved in the lifespan-expanding effect of CR in mice too. Nox4 (NADPH oxidase 4) is another gene that is downregulated by CR in our analysis. It is known that high levels of NADPH oxidases are associated with increased oxidative stress and damage (190). Stress-resistant mouse strains show an increased lifespan, which indicates that the downregulation of Nox4 by CR could be involved in the lifespan-expanding effect of CR as well.

Furthermore, CR downregulates the expression of several major urinary proteins (Mup3, Mup4, Mup5, Mup6, Mup10, Mup20, Mup21). It is already mentioned that dietary restriction decreases Mup expression in adipose tissue and liver (191; 192; 193; 28; 194). Mups are markers of reproductive performance. Higher reproduction is associated with increased protein synthesis and enhanced oxidative damage. Therefore, reduced reproduction, which is indicated by reduced Mups, can result in an increased lifespan (28). Moreover, Mups are involved in glucose, lipid and energy metabolism, so mTOR may be involved in the regulation of Mup expression (195; 196; 197; 194). ZHOU et al. (2009) have shown that Mup1 overexpression decreases the expression of gluconeogenic and lipogenic genes (196). Therefore, it is possible that CR increases gluconeogenesis at least partly by inhibiting the expression of Mups. It is known that Mup expression is increased by growth hormone, testosterone and thyroid hormone, whereas glucocorticoids inhibit expression, all factors which are influenced by CR as well (198; 194).

The downregulation of growth differentiation factor 15 (Gdf15) could be involved in mediating CR effects as well. It is known that Gdf15 increases with ageing and is a predictor of all-cause mortality and cardiovascular diseases (199). Moreover, a high Gdf15 level functions as a biomarker for mitochondrial disorders (200). The expression of Gdf15 is increased in stress conditions (199). To our knowledge, the influence of CR on Gdf15 expression has not been investigated before but could be an important factor in CR-mediated longevity. Gdf15 expression is strongly downregulated by all phytonutrients except Gi.

Foxa3, which is downregulated by CR in our analysis, could be another interesting gene. Whereas PANOWSKI et al. (2007) have shown that PHA-4 (an orthologue of Foxa1, 2, 3) is involved in the lifespan-expanding effect of dietary restriction in *C. elegans* but is not necessary, MA et al. (2014) have shown that Foxa3 KO mice show an increased lifespan and insulin sensitivity (201; 202). Moreover, Foxa3 is a negative regulator of PGC1α, which is known to be increased under the CR condition (202). The role of Foxa3 on murine lifespan and the mechanisms that are targeted by Foxa3 should be further analysed.

Regarding the up- and downregulated genes of CR, some genes that are predominant in females, such as Fmo3, Nnmt, Cyp2b9, Cyp2b10, Cyp2b13 and Cyp4a10, are upregulated, whereas genes that are predominant in males, such as hydroxy-delta-5-steroid dehydrogenase 3 beta- and steroid delta-isomerase 5 (Hsd3b5), Cyp4a12a, Cyp4a12b, Cyp7b1, UDP glucuronosyltransferase 2 family polypeptide B1 (Ugt2b1) and solute carrier organic anion transporter family member 1a1 (Slco1a1), are downregulated (203; 179; 204; 205). Interestingly, the expression of most genes is testosterone dependent, indicating a reduction of testosterone level by CR in murine liver (203). It has already been shown that CR feminises gene expression (10).

Other genes could be interesting as well, but the connection to CR mechanisms is not really clear. For example, the upregulation of the regulator of G-protein signalling 16 (Rgs16) could be involved in the inhibition of mTOR signalling (206), the downregulation of Slco1a1 could promote the accumulation of bile acids in the serum (207), the upregulation of eukaryotic translation initiation factor 4E binding protein 3 (Eif4ebp3) could be involved in regulation of protein synthesis (208), the upregulation of neuregulin 4 (Nrg4) could mediate the downregulated effect on lipogenesis (209) and the upregulation of protein phosphatase 1 regulatory subunit 3G (Ppp1r3g) could be important for the CR effects on glucose metabolism (210; 211). Nonetheless, the other genes mentioned in Appendix P could be important for mediating CR effects as well.

3.3.4. Overlap between the top 50 regulated genes identified by q-value

To specify the functions which are mimicked by the phytonutrients, the top 50 up- and downregulated genes (sorted by q-value) of each intervention are compared, and the overlapped genes are identified. The functions of these genes are recognised with CPDB and NCBI:gene (Table 19 and 20). R shows the largest number of overlapped genes (seven upregulated genes and 11 downregulated genes) followed by Gi (six upregulated genes and four downregulated genes) and Qd (four upregulated genes and four downregulated genes). This indicates furthermore that R is the best mimetic for CR.

Interestingly, Igfpb1 and Nnmt are included in this list too. A new observation is that Igfpb1 is one of the top genes regulated by Qd but not by Qi. Previous analyses have not indicated that Qd could be a CR mimetic except of the upregulation of Sirt1. The strong upregulation of Igfpb1 could refute this assumption. Figure 8 illustrates the regulation of Igfpb1 and the mechanisms influenced by Igfpb1. The expression of Igfpb1 seems to be dependent on Sirt1 through Foxo and mitogen-activated protein kinase (MAPK) activation (172). Whereas Sirt1 is upregulated by Qd and CR in our analysis, the expression of Foxo1 is only increased by CR. However, CR and Qd reduce Irs1 expression, which probably indicates that PI3K/Akt signalling is reduced. This inhibition could lead to increased Foxo1 activity without affecting gene expression. In contrast, FARDINI et al. (2014) have shown that Foxo1 inhibits PI3K/Akt signalling itself by upregulating Igfpb1 (212). Foxa1, which is upregulated by CR and Qd, is also known to increase igfbp1 activity (213; 214).

Another indication that Qd could be involved in the inhibition of Igf1 signalling is the reduced expression of Igfals. As mentioned above, Igfals, together with Igfpb3, is important for Igf1 stability and increases the half-life of Igf1 from a few minutes up to 15 hours (173). The expression of Igfpb3 is downregulated by CR as well but is not influenced by Qd. Besides, Igfpb1 could be involved in the inhibition of mTOR signalling by reducing PI3K/Akt signalling (215).

Metallothionein 1 (Mt1) and lipin 1 (Lpin1), two genes upregulated by CR and Qd, could further indicate that Qd is a CR memetic. It has already been shown that CR increases the expression of Mt1 in liver and that overexpression in murine heart is associated with increased lifespan (220; 221; 178). Mt1 is involved in the inhibition of oxidative stress damage and in scavenging of ROS (220; 222). As mentioned above, this is supposed to be important for mediating beneficial CR effects. Another Mt1 mechanism that could be involved in CR effects is that Mt1 lowers zinc bioavailability (220; 223). This could explain the reduced immune response observed by CR, which is important for

successful ageing. The results of SWINDELL et al. (2010) indicate that Mt1 expression is increased by inhibition of GH/Igf1 signalling in murine kidney (224).




Figure 9: Interaction of the overlapped genes between caloric restriction and quercetin (administered per diet)

(Purple arrow: CR effect, green arrow: Qd effect, ↑: upregulation, ↓: downregulation, ↗: tendency in upregulation, ↘: tendency in downregulation, - no influence on gene expression)

Lpin1 is less investigated in the context of CR and lifespan, but knockdown of Lpin1 decreases PGC-1 α (225; 226). Moreover, FINCK et al. (2005) have mentioned that Lpin1 is a coactivator for PPAR α and PGC-1 α and is involved in the enhanced fatty acid beta oxidation in murine liver (227). Based on the fact that Sirt1 is important for PGC-1 α , it is possible that Lpin1 is involved in Sirt1-dependent CR effects. This is supported by the observation that ethanol-induced Sirt1 inhibition affects transcription regulation by Lpin1 in alcoholic fatty liver disease (228; 229). Moreover, troxerutin, a derivate of the bioflavonoid rutin, improves hepatic lipid homeostasis by increasing NAD $^+$ level and Sirt1 activity and reduces Lpin1 signalling (230). It is assumed that Sirt1 activates AMPK, inhibiting the mammalian target of rapamycin complex 1 (mTORC1) signalling and modified Lpin1 signalling. The influence of CR on the interaction of Lpin1 and Sirt1 in healthy liver should be further investigated.

Table 19: Pathways and biological functions of the overlapping genes between the top 50 upregulated genes of caloric restriction and the plant bioactives upregulated genes

	R	P	Qd	Qi	Gi	function / pathways
Mt1	X	X				<u>pathways</u> : oxidative stress, mineral absorption; <u>GO function</u> : copper and zinc ion binding; <u>GO process</u> : cellular zinc ion homeostasis, cellular response to zinc ion and chromate, detoxification of copper ion, negative regulation of neuron apoptotic process, nitric oxide mediated signal transduction
Cyp2b9	X					<u>pathways</u> : biological oxidation (phase I), retinol metabolism; <u>GO function</u> : arachidonic acid epoxygenase activity, aromatase activity, heme, oxygen or iron ion binding, steroid hydroxylase activity; <u>GO process</u> : epoxygenase P450 pathway, exogenous drug catabolic process, oxidation-reduction process, xenobiotic metabolic process
Cyp2a5	X		X			<u>pathways</u> : biological oxidation (phase I), retinol metabolism; <u>GO function</u> : arachidonic acid epoxygenase activity, coumarin 7-hydroxylase activity, enzyme, heme or oxygen binding, steroid hydroxylase activity; <u>GO process</u> : cellular response to cadmium ion, coumarin metabolic process, epoxygenase P450 pathway, exogenous drug catabolic process, heme metabolic process, oxidation-reduction process, xenobiotic metabolic process
Nsdhl	X					<u>pathways</u> : steroid / cholesterol biosynthesis; <u>GO function</u> : 3-beta-hydroxy-delta5-steroid dehydrogenase activity, sterol-4-alpha-carboxylate 3-dehydrogenase (decarboxylating) activity; <u>GO process</u> : lipid metabolic process, oxidation-reduction process, smoothened signaling pathway
Nipal1	X					<u>GO function</u> : magnesium ion transmembrane transporter activity
Usp2	X	X				<u>pathway</u> : TNF-alpha NF-kB signaling pathway; <u>GO function</u> : cysteine-type endopeptidase activity, ubiquitin-specific protease activity, metal ion binding; <u>GO process</u> : positive regulation of mitotic cell cycle, circadian behavior, circadian regulation of gene expression, negative regulation of transcription from RNA polymerase II promoter, proteasome-mediated ubiquitin-dependent protein catabolic process, protein deubiquitination, protein stabilization, regulation of proteasomal protein catabolic process
Rgs16	X					<u>pathways</u> : EGFR1 signaling, calcium regulation in the cardiac cell; <u>GO function</u> : GTPase activator activity; <u>GO process</u> : termination of G-protein coupled receptor signaling pathway, positive regulation of GTPase activity
Nnmt	X					<u>pathways</u> : biological oxidation (phase II) / methylation, nicotinate and nicotinamide metabolism; <u>GO function</u> : nicotinamide N-methyltransferase activity
Dhcr7	X					<u>pathways</u> : activation of gene expression by SREBF (SREBP) / regulation of cholesterol biosynthesis by SREBP (SREBF); <u>GO function</u> : 7-dehydrocholesterol reductase activity; <u>GO process</u> : blood vessel development, cell differentiation, oxidation-reduction process, regulation of cell proliferation
Lpin1		X				<u>pathways</u> : glycerophospholipid metabolism, triglyceride biosynthesis , cell cycle (M Phase); <u>GO function</u> : RNA polymerase II transcription factor binding, transcription coactivator activity, histone deacetylase binding, phosphatidate phosphatase activity, peroxisome proliferator activated receptor binding; <u>GO process</u> : actin cytoskeleton reorganization, cellular response to insulin stimulus, fatty acid catabolic process, mitochondrial fission, negative and positive regulation of transcription from RNA polymerase II promoter, positive regulation of histone deacetylation, protein dephosphorylation, ruffle organization
Nrg4	X					<u>pathways</u> : innate and adaptive immune system, cytokine signaling (interleukin signaling), involved in signal transduction, MAPK family signaling cascades
Igfbp1	X					<u>pathways</u> : regulation of Insulin-like Growth Factor (IGF) transport and uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs), metabolism of proteins; <u>GO function</u> : insulin-like growth factor I and II binding; <u>GO process</u> : regulation of cell growth, regulation of insulin-like growth factor receptor signaling
Fmo3	X	X				<u>pathway</u> : phase I of biological oxidation / detoxification; <u>GO function</u> : NADP binding, amino acid binding, flavin adenine dinucleotide binding, N,N-dimethylaniline monooxygenase activity, trimethylamine monooxygenase activity
Sult2a3	X	X				<u>pathways</u> : biological oxidation (phase II), bile secretion; <u>GO function</u> : sulfotransferase activity, 3'-phosphoadenosine 5'-phosphosulfate binding, drug binding
Cyp17a1	X					<u>pathways</u> : biological oxidation (phase I), steroid and Steroid hormone biosynthesis, metabolism of lipids and lipoproteins; <u>GO function</u> : 17-alpha-hydroxyprogesterone aldolase activity, heme or iron binding, steroid 17-alpha-monooxygenase activity; <u>GO process</u> : glucocorticoid biosynthetic process, positive regulation of steroid hormone biosynthetic process, oxidation-reduction process, progesterone metabolic process
Cyp2b13	X					<u>pathways</u> : arachidonic acid metabolism, biological oxidation (phase I), retinol metabolism, steroid hormone biosynthesis; <u>GO functions</u> : arachidonic acid epoxygenase activity, heme binding, oxidoreductase activity, steroid hydroxylase activity, oxygen binding
Mfsd2a	X					<u>GO functions</u> : phospholipid transporter activity, symporter activity; <u>GO process</u> : fatty acid / phospholipid transport, lipid transport across blood brain barrier
Slc22a7	X					<u>pathways</u> : SLC-mediated transmembrane transport / transport of glucose and other sugars, bile salts and organic acids, metal ions and amine compounds



Table 20: Pathways and biological functions of the overlapping genes between the top 50 downregulated genes of caloric restriction and the plant bioactives downregulated genes

	R	P	Qd	Qi	Gi	function / pathways
Elovl3	X			X		<u>pathways:</u> fatty acid biosynthesis/elongation, triglyceride biosynthesis, fatty acyl-CoA biosynthesis, linoleic acid (LA) and alpha-linolenic acid (ALA)metabolism; <u>GO function:</u> transferase activity
Hsd3b5	X		X	X		<u>pathways:</u> metabolism of vitamin D, steroids and steroid hormones (e.g. androgen), glucocorticoid and mineralcorticoid metabolism; <u>GO function:</u> oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor, steroid dehydrogenase activity, aldo-keto reductase (NADP) activity, NAD binding
Pdilt	X					<u>GO function:</u> protein disulfide isomerase activity, protein binding, peptidyl-proline 4-dioxygenase activity; <u>GO process:</u> spermatid development, cell migration, cell differentiation, cell redox homeostasis, multicellular organismal development, protein folding
G0s2	X	X	X			<u>pathways:</u> circadian regulation; <u>GO process:</u> positive regulation of extrinsic apoptotic signaling pathway
Wfdc2	X					<u>GeneRIFs:</u> affecting high density lipoprotein levels (216); <u>GO function:</u> aspartic-type endopeptidase inhibitor activity, cysteine-type endopeptidase inhibitor activity, serine-type endopeptidase inhibitor activity; <u>GO process:</u> negative regulation of peptidase activity
Cyp4a12a	X			X		<u>pathways:</u> phase I of biological oxidation / detoxification, arachidonic acid metabolism / synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE), retinol metabolism, PPAR signaling pathway, fatty acid degradation, inflammatory mediator regulation of TRP channels; <u>GO function:</u> iron ion binding, heme binding, aromatase activity, alkane 1-monoxygenase activity
Ddc	X					<u>pathways:</u> catecholamine biosynthesis; serotonin and melatonin biosynthesis; tyrosine, tryptophan, histidine and phenylalanine metabolism / biogenic amine synthesis; dopaminergic synapse/ dopaminergic neurogenesis; <u>GO function:</u> L-dopa decarboxylase activity, amino acid binding, enzyme binding, protein domain specific binding, pyridoxal phosphate binding; <u>GO process:</u> cellular amino acid metabolic process, dopamine biosynthetic process, response to toxic substance, serotonin biosynthetic process, synaptic vesicle amine transport
Alas2	X					<u>pathways:</u> metabolism of porphyrins / heme biosynthesis; glycine, serine and threonine metabolism; <u>GO function:</u> 5-aminolevulinate synthase activity, coenzyme binding, glycine binding, pyridoxal phosphate binding; <u>GO process:</u> heme biosynthetic process, hemoglobin biosynthetic process, cellular iron ion homeostasis, response to hypoxia, erythrocyte differentiation, cellular iron ion homeostasis
Hspf1	X					<u>pathways:</u> protein processing in endoplasmic reticulum, vesicle-mediated transport (binding and uptake of ligands by scavenger receptors), cellular response to stress; <u>GO process:</u> chaperone mediated protein folding requiring cofactor, negative regulation of intrinsic apoptotic signaling pathway in response to hydrogen peroxide, negative regulation of establishment of protein localization to mitochondrion, negative regulation of p38MAPK cascade, positive regulation of MHC class I biosynthetic process, positive regulation of NK T cell activation, positive regulation of protein tyrosine kinase activity, positive regulation of transcription from RNA polymerase II promoter, regulation of microtubule cytoskeleton organization
Car3	X	X				<u>pathways:</u> reversible hydration of carbon dioxide, nitrogen metabolism; <u>GO function:</u> metal ion binding (zinc, nickel), carbonate dehydratase activity, phosphatase activity; <u>GO process:</u> response to oxidative stress
Creld2	X					<u>GeneRIFs:</u> Probably a mediator in regulating the onset and progression of endoplasmic reticulum stress-associated diseases (217); <u>GO function:</u> calcium ion binding
Obp2a	X	X				<u>GeneRIFs:</u> Decreases lipid levels in liver → inhibits lipogenesis, stimulates beta-oxidation (218); enhances insulin signaling, suppress hepatic glucose production (219)
Hspa1b	X	X				<u>pathways:</u> metabolism of RNA / spliceosome / regulation of mRNA stability, MAPK signaling pathway, endocytosis, protein processing in endoplasmic reticulum, estrogen signaling pathway, antigen processing and presentation, cellular response to stress/ heat stress; <u>GO Process:</u> negative regulation of apoptotic process, negative regulation of cysteine-type endopeptidase activity involved in apoptotic process, response to heat
Csad	X					<u>pathways:</u> degradation of cysteine and homocysteine, sulfur amino acid metabolism, taurine and hypotaurine metabolism, mRNA processing; <u>GO function:</u> sulfinoalanine decarboxylase activity, pyridoxal phosphate binding; <u>GO process:</u> L-cysteine catabolic process to taurine
Ifit1			X			<u>pathways:</u> interferon signaling, antiviral mechanism by IFN-stimulated genes, cytokine signaling in immune system; <u>GO Process:</u> cellular response to interferon-alpha and -beta; innate immune response; defense response to virus

R = Resveratrol, P = phloridzin, Qd=quercetin diet, Qi = quercetin intraperitoneal, Gi = genistein intraperitoneal

The downregulation of G0s2 by CR, R, Qd and Qi could be interesting as well. NIELSEN et al. (2011) have shown that fasting decreases G0s2 expression in human adipose tissue, and AHN et al. 2013 have shown that short-term fasting decreases G0s2 expression in porcine adipose tissue, whereas 8-day CR does not influence G0s2 expression (231; 232). Nonetheless, this indicates that G0s2 expression depends on nutritional status. G0s2 codes for an inhibitor of adipose triglyceride lipase (ATGL), indicating that downregulation of G0s2 promotes lipolysis. Moreover, it is known that G0s2 is involved in the regulation of the cell cycle (231). ZHANG et al. (2016) have shown that G0s2 expression is inhibited by Foxo1 in liver and that this regulation is important for Foxo1's dependent effect on glucose and lipid metabolism (233). This probably indicates that G0s2 is involved in mediating CR effects and that phytonutrients are able to mimic this effect.

3.3.5. Calculation of Pearson correlation for gene expression between CR and phytonutrients in over-represented pathways for a better comparison between phytonutrients and their mimicking effect

Although there are several indications that R, Qd and Qi could be CR mimetics, it is difficult to say which is the best phytonutrient for mimicking CR effects because not every Entrez ID has been investigated in relation to every phytonutrient. For a better comparison, only the Entrez IDs which are investigated by all diet interventions are included in the rank product analysis (Qi and Gi are excluded). This leads to 15552 investigated Entrez IDs (compare Figure 2). Furthermore, it would be interesting if synergistic effects indicated that a diet rich in different phytonutrients would show a better mimicking effect than ingestion of one phytonutrient alone. For this, the results are combined whereby opposite regulations by the different phytonutrients lead to an exclusion of the gene. The mean values of the fold changes are calculated for the heat map.

Table 21: Number of regulated genes based on the q-value or the q-value in combination with a fold change (FC) > 1.5

	CR	R	P	Q	RP	RQ	PQ	RPQ
<i>number of upregulated genes based on ...</i>								
q value	1365	1185	1151	603	1973	1553	1453	2192
q-value & FC	38	304	6	23	304	315	26	316
<i>number of downregulated genes based on ...</i>								
q-value	1833	998	1793	548	2316	1347	1988	2462
q-value & FC	80	216	13	2	225	214	14	223

CR = caloric restriction, R = resveratrol, P = phloridzin, Q = quercetin, RP / RQ / PQ / RPQ = combined results of the different interventions, FC = fold change




Figure 10: Heat map showing the fold changes of all upregulated CR genes with a fold change > 1.5 in presence of the different plant bioactives (individual and synergistic analysis)

Table 21 shows the number of regulated genes either based on the q-value or on the q-value combined with a fold change > 1.5. Figure 12 shows the percentage overlap based on the results from Table 21. These results show that when only the q-value is respected, R and P show a similar overlap, although P shows a bigger overlap for downregulated genes. Combining the results of R and P indicates that both phytonutrients together have a greater effect in mimicking CR. Further inclusion of Qd shows only a very small benefit regarding the upregulated genes. After including the fold change, only R was able to mimic the gene expression of CR. Combining the results of R with P or P and Qd shows only a small benefit, whereby a combination with Qd shows a deterioration in the overlap.

On further analysis, only the CR genes with a fold change > 1.5 are included. The heat maps presented in Figures 10 and 11 show the gene expression regulation for these genes by CR and the phytonutrients including synergistic effects. A small fold change indicates an upregulation in gene expression. Next it was analysed whether a phytonutrient can mimic the gene expression over all 15552 genes. Therefore, the Pearson correlation was calculated (Table 22A). All phytonutrients show a significant weak positive correlation with CR. Although the correlation between CR and Qd is weak, the result is significant probably because of the large number of genes.




Figure 11: Heat map showing the fold changes of all downregulated CR genes with a fold change > 1.5 in presence of the different plant bioactives (individual and synergistic analysis)

Table 22: A.) Pearson correlation between CR and the plant bioactives (including synergistic analysis) based on all 15552 IDs; B.) Number of pathways with a high Pearson correlation between calorific restriction (CR) and the plant bioactives (total: 358 CR pathways)

(R = based on the fold changes of resveratrol, P = based on the fold changes of phloridzin, Q = based on the fold changes of quercetin, RP = based on the mean values of the fold changes of resveratrol and phloridzin, RQ = based on the mean values of the fold changes of resveratrol and quercetin, PQ = based on the mean values of the fold changes of phloridzin and quercetin, RPQ = based on the mean values of the fold changes of resveratrol, phloridzin and quercetin)

A.)	Pearson correlation		Pearson correlation					
		p-value	≥ 0.5	≥ 0.6	≥ 0.7	≥ 0.8	≥ 0.9	≥ 0.95
R	0.26	< 0.0001	106	82	58	40	23	17
P	0.20	< 0.0001	53	37	26	16	6	2
Q	0.02	< 0.05	45	34	26	16	4	1
RP	0.31	< 0.0001	121	93	75	45	24	18
RQ	0.24	< 0.0001	104	90	63	36	22	17
PQ	0.17	< 0.0001	56	41	32	18	8	4
RPQ	0.29	< 0.0001	120	99	77	39	25	19

Overlap based on the q-value



Overlap based on the q-value & fold change




Figure 12: Overlap between CR and plant bioactives including synergistic effects

(Calculation based on the regulated genes identified by limited gene pool (15552 IDs); significant regulated genes identified with the q-value or the q-value in combination with a fold change > 1.5)

For a better determination of which pathways are mimicked by the phytonutrients, a Pearson calculation is repeated for all pathways that include at least one CR gene with a fold change > 1.5. Small pathways that consist of one or two genes are excluded. This leads to 358 relevant pathways (Appendix L). Table 22B shows the number of pathways with a correlation of 0.5 to 0.95 for the respective intervention. R seems to be able to mimic the gene expression of the most pathways. Furthermore, a combination with P, Qd or both is able to increase the number of mimicked pathways. Table 23 shows all pathways with a Pearson correlation r > 0.9.

Nnmt and Igfpb1 are genes which seem to be of interest in our analysis. The pathways ‘Regulation of Insulin-like Growth Factor (IGF) Transport and Uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)’ (Reactome) and ‘Nicotinate and nicotina-

mide metabolism – *Mus musculus* (mouse)’ (KEGG) include these genes, so the heat maps of these pathways are presented in Figures 13 and 14. As expected, Qd is the only dietary phytonutrient that shows a positive correlation regarding the Igfbp1 pathway; the other phytonutrients show a negative correlation (see Appendix L). The results for the pathway ‘Nicotinate and nicotinamide metabolism – *Mus musculus* (mouse)’ (KEGG) indicate that P and all combinations with P (RP, PQ, RPQ) show a high correlation with CR expression.

Table 23: Pathways with a high Pearson correlation (≥ 0.9) in gene expression between caloric restriction (CR) and the plant bioactives

pathway	R	P	Q	RP	RQ	PQ	RPQ
Glycoprotein hormones - Reactome	0,86	0,89	0,91	0,89	0,89	0,94	0,91
Peptide hormone biosynthesis - Reactome	0,87	0,84	0,87	0,89	0,90	0,90	0,91
Steroid Biosynthesis - Wikipathways	0,98	0,25	0,50	0,99	0,98	0,44	0,98
Androgen biosynthesis - Reactome	0,97	0,31	0,52	0,97	0,97	0,46	0,97
Metabolism of steroid hormones and vitamin D - Reactome	0,97	0,35	0,37	0,97	0,97	0,42	0,96
Steroid hormones - Reactome	0,97	0,35	0,37	0,97	0,97	0,42	0,96
FMO oxidizes nucleophiles - Reactome	0,93	0,79	0,91	0,99	0,98	0,85	1,00
L-cysteine degradation I - MouseCyc	1,00	0,95	0,51	1,00	0,98	1,00	1,00
Serotonin and melatonin biosynthesis - Reactome	1,00	-0,38	-0,38	0,98	1,00	-0,39	0,97
Catecholamine biosynthesis - Reactome	0,99	-0,50	-0,69	0,99	0,98	-0,76	0,99
serotonin and melatonin biosynthesis - MouseCyc	1,00	-0,38	-0,38	0,98	1,00	-0,39	0,97
biosynthesis of serotonin and melatonin - MouseCyc	1,00	-0,38	-0,38	0,98	1,00	-0,39	0,97
4-hydroxyproline degradation I - MouseCyc	0,90	0,96	0,99	0,95	0,94	0,98	0,96
Organic anion transport - Reactome	0,95	0,97	0,77	0,99	0,96	0,96	0,99
tetrapyrrole biosynthesis II - MouseCyc	0,98	0,65	0,32	0,99	0,96	0,57	0,97
phenylalanine degradation III - MouseCyc	0,90	-0,26	0,84	0,65	0,93	0,26	0,75
Methionine salvage pathway - Reactome	0,98	0,78	0,78	0,93	0,97	0,94	0,97
Mineralocorticoid biosynthesis - Reactome	1,00	0,32	0,48	1,00	1,00	0,44	1,00
Glucocorticoid biosynthesis - Reactome	1,00	0,30	0,49	1,00	1,00	0,44	1,00
ADP signalling through P2Y purinoceptor 1 - Reactome	0,51	0,52	0,90	0,58	0,69	0,72	0,68
Highly calcium permeable nicotinic acetylcholine receptors - Reactome	0,70	0,94	0,80	0,90	0,77	0,96	0,91
Taurine and hypotaurine metabolism - <i>Mus musculus</i> (mouse) - KEGG	0,95	0,92	-0,71	0,98	0,92	0,84	0,99
Ovarian steroidogenesis - <i>Mus musculus</i> (mouse) - KEGG	0,96	0,21	0,40	0,96	0,96	0,33	0,96
Nicotine Activity on Dopaminergic Neurons - Wikipathways	0,93	-0,22	0,18	0,92	0,95	-0,15	0,94
Glucuronidation - Wikipathways	0,90	0,15	0,06	0,89	0,90	0,14	0,86
Reversible Hydration of Carbon Dioxide - Reactome	0,91	0,18	0,85	0,90	0,92	0,76	0,92
Glucocorticoid & Mineralcorticoid Metabolism - Wikipathways	1,00	0,34	0,53	1,00	1,00	0,48	1,00
Highly calcium permeable postsynaptic nicotinic acetylcholine receptors - Reactome	0,64	0,93	0,74	0,89	0,74	0,93	0,90
Biogenic Amine Synthesis - Wikipathways	0,90	-0,09	-0,10	0,91	0,87	-0,11	0,86
Dopminergic Neurogenesis - Wikipathways	0,90	0,01	-0,15	0,91	0,84	-0,11	0,86




Figure 13: Heat map of genes of the pathway ‘Regulation of Insulin-like Growth Factor (IGF) Transport and Uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)’ (Reactome)




Figure 14: Heat map of genes of the pathway ‘Nicotinate and nicotinamide metabolism - Mus musculus (mouse)’ (KEGG)

3.4. Analysis of longevity genes in mouse

As already mentioned, CR is known for a lifespan-expanding effect (106). Therefore, it is of interest whether phytonutrients are able to extend the lifespan as well. In *Mus musculus*, 112 genes are mentioned in context with longevity; 34 of them have anti-longevity effects, 75 have pro-longevity effects and 3 have not been classified (234). Genes in each category are up- and downregulated by the different intervention groups (Table 24). Therefore, it is impossible to say whether CR or a phytonutrient has an impact on longevity because it is difficult to say which gene has the biggest impact. According to GenAge, Ghr is mentioned as the gene with the largest impact on lifespan in mice (235). The knockout of Ghr increases lifespan by up to 40-50%. The results show that CR, P and Qi downregulate the expression of Ghr (Table 25). R, Qd and Gi do not influence gene expression.

Table 24: Influence of caloric restriction and plant bioactives on longevity genes

anti-longevity genes (n = 34)	CR	R	P	Q	Qi	Gi
upregulation	3	1	1	2	3	2
downregulation	5	2	4	2	3	2
pro-longevity genes (n = 75)	CR	R	P	Q	Qi	Gi
upregulation	8	8	5	5	5	3
downregulation	17	5	13	3	6	2

Table 25: CR longevity genes which are influenced by at least one of the plant bioactive

downregulated anti-longevity genes by CR			R	P	Q	Qi	Gi
12575	Cdkn1a	cyclin-dependent kinase inhibitor 1A (P21)		↓		↑	
66143	Eef1e1	eukaryotic translation elongation factor 1 epsilon 1					
14600	Ghr	growth hormone receptor			↓		↓
625249	Gpx4	glutathione peroxidase 4		----	↑		
20416	Shc1	src homology 2 domain-containing transforming protein C1					↓

upregulated pro-longevity genes by CR

12189	Brca1	breast cancer 1					
218294	Cdc14b	CDC14 cell division cycle 14B			↓	↓	
14824	Grn	granulin		↑			
17685	Msh2	mutS homolog 2 (E. coli)					
114774	Pawr	PRKC, apoptosis, WT1, regulator			↑		
18534	Pck1	phosphoenolpyruvate carboxykinase 1, cytosolic		↑			
22590	Xpa	xeroderma pigmentosum, complementation group A		↑			
22596	Xrcc5	X-ray repair complementing defective repair in Chinese hamster cells 5					

Next it is analysed whether the phytonutrients regulate the same genes as CR (Table 25). R could be the only phytonutrient which mimics the effect of CR on longevity because four genes are regulated in the same direction as CR and no gene is regulated in the opposite direction. Qd regulates only two of these genes, but both are regulated

in the opposite direction as compared to CR. P regulates two genes in the same direction as CR, but the other two show opposite regulation. However, it is nonetheless possible that phytonutrients have a life-extending effect without affecting the same genes as CR.

3.5. Transcription factor binding sites (TFBs)

To further clarify whether the phytonutrients are mimetics of CR, it was investigated whether they address the same transcription factor binding sites (TFBs) as CR. To identify TFBs which are important for CR, all significantly regulated CR genes which are investigated by at least one phytonutrient were uploaded in oPOSSUM separated by up- and downregulated genes. Not all downregulated genes could be analysed because oPOSSUM limits the number of potential genes to 2000. Therefore, only the top 2000 downregulated genes sorted by fold change were investigated.

All TFBs with $z > 5$ or $F > 5$ were included in the analysis. This resulted in 121 TFBs which are addressed by the upregulated CR genes and 94 TFBs which are addressed by downregulated CR genes (Appendix U and V). To determine whether the phytonutrients address the same TFBs as CR, next the overlapped genes between CR and the phytonutrients separated by up- and downregulated genes were analysed with oPOSSUM. All TFBs with $z > 5$ or $F > 5$ were included too. Table 26 shows the absolute and percentage overlaps between the TFBs mentioned by the CR genes and the TFBs revealed by the overlapped genes between CR and the respective phytonutrient. All phytonutrients address TFBs which are mentioned by CR. Regarding the upregulated genes, Qd shows the largest overlap followed by P and R. Regarding the downregulated genes P, Qd and Qi show a high overlap with CR. This confirmed that Qd could be a CR mimetic. Tables 27 and 28 show the top TFBs addressed by CR. For this, the top 10 TFBs sorted by z-score and the top 10 TFBs sorted by F-score are summarised. Table 29 shows the functions of the transcription factors (TF) which typically address the top TFBs mentioned by CR. The functions are revealed with the UniProt database. Functions and biological processes which are not important in murine liver cases are not mentioned but instead are presented in Appendix W.

Table 26: Total and percental overlap in transcription factor binding sites (TFBs) putatively affected by CR and the plant bioactives

	total (and percental) overlap				
	R	P	Qd	Qi	Gi
121 TFBs addressed by upregulated CR genes	67 (55%)	73 (60%)	78 (64%)	53 (44%)	61 (50%)
94 TFBs addressed by downregulated CR genes	17 (18%)	64 (68%)	61 (65%)	67 (71%)	13 (14%)

TFBs = transcription factor binding sites, CR = caloric restriction; R = resveratrol, P = phloridzin, Qd = quercetin (diet), Qi = quercetin (intraperitoneal), Gi = genistein (intraperitoneal)

Table 27: Top transcription factor binding sites (TFBs) putatively affected by upregulated genes of caloric restriction identified with the z- or F-score and the overlap with the plant bioactives

JASPAR ID	Transcription factor	UniProt ID	R	P	Qd	Qi	Gi
MA0006.1	Ahr - Aryl hydrocarbon receptor, Arnt - Aryl hydrocarbon receptor nuclear translocator	P30561 P53762					
MA0141.1	Esrrb - Steroid hormone receptor ERR2	Q61539					
MA0099.2	Fos - Proto-oncogene c-Fos, Jun - Transcription factor AP-1	P01101 P05627					
MA0158.1	Hoxa5 - Homeobox protein Hox-A5	P09021					
PB0142.1	Jdp2 - Jun dimerization protein 2	P97875					
MA0039.2	Klf4 - Krueppel-like factor 4	Q60793					
PB0039.1	Klf7 - Krueppel-like factor 7	Q99JB0					
MA0059.1	Max - Protein max, Myc - Myc proto-oncogene protein	P28574 P01108					
MA0104.2	Mycn - N-myc proto-oncogene protein	P03966					
MA0056.1	Mzf1 - Myeloid zinc finger 1	A1L358					
MA0057.1	Mzf1 - Myeloid zinc finger 1	A1L358					
MA0060.1	Nfyα - Nuclear transcription factor Y subunit alpha	P23708					
MA0065.2	Pparg - Peroxisome proliferator-activated receptor gamma, Rxra - Retinoic acid receptor RXR-alpha	P37238 P28700					
MA0079.2	Sp1 - Transcription factor Sp1	O89090					
MA0081.1	Spib - Transcription factor Spi-B	O35906					
MA0095.1	Yy1 - Transcriptional repressor protein YY1	Q00899					
MA0103.1	Zeb1 - Zinc finger E-box-binding homeobox 1	Q64318					
MA0146.1	Zfx - Zinc finger X-chromosomal protein	P17012					
MA0130.1	Znf354c - Zinc finger protein 354C	Q571J5					

R = resveratrol, P = phloridzin, Qd = quercetin (diet), Qi = quercetin (intraperitoneal), Gi = genistein (intrap.)

Table 28: Top transcription factor binding sites (TFBs) putatively affected by downregulated genes of caloric restriction identified with the z- or F-score and the overlap with the plant bioactives

JASPAR ID	Transcription factor	UniProt ID	R	P	Qd	Qi	Gi
PB0002.1	Arid5a - AT-rich interactive domain-containing protein 5A	Q3U108					
MA0102.2	Cebpa - CCAAT/enhancer-binding protein alpha	P53566					
MA0099.2	Fos - Proto-oncogene c-Fos, Jun - Transcription factor AP-1	P01101 P05627					
MA0041.1	Foxd3 - Forkhead box protein D3	Q61060					
PB0016.1	Foxj1 - Forkhead box protein J1	Q61660					
PB0017.1	Foxj3 - Forkhead box protein J3	Q8BUR3					
PB0123.1	Foxl1 - Forkhead box protein L1	Q64731					
MA0046.1	Hnf1a - Hepatocyte nuclear factor 1-alpha	P22361					
MA0158.1	Hoxa5 - Homeobox protein Hox-A5	P09021					
MA0056.1	Mzf1 - Myeloid zinc finger 1	A1L358					
MA0060.1	Nfyα - Nuclear transcription factor Y subunit alpha	P23708					
MA0063.1	Nkx2-5 - Homeobox protein Nkx-2.5	P42582					
MA0122.1	Nkx3-2 - Homeobox protein Nkx-3.2	P97503					
MA0081.1	Spib - Transcription factor Spi-B	O35906					
MA0084.1	Sry - Sex-determining region Y protein	Q05738					
PB0080.1	Tbp - TATA-box-binding protein	P29037					
MA0103.1	Zeb1 - Zinc finger E-box-binding homeobox 1	Q64318					
PB0093.1	Zfp105 - Putative uncharacterized protein	Q3UHV1					
MA0130.1	Znf354c - Zinc finger protein 354C	Q571J5					

R = resveratrol, P = phloridzin, Qd = quercetin (diet), Qi = quercetin (intraperitoneal), Gi = genistein (intrap.)

Table 29: Overview on the transcription factors and their biological function

TF	function / biological process
Ahr	activates the expression of multiple phase I and II enzyme genes / xenobiotic metabolic process / response to stress / toxic substance / xenobiotic stimulus, circadian regulation of gene expression, regulation of cell cycle / morphogenesis / size, cellular response to cAMP, common bile duct and liver development, regulation of gene expression and transcription
Arid5a	regulation of histone acetylation, regulation of transcription
Arnt	cell differentiation, regulation of hormone biosynthetic process, regulation of protein sumoylation, regulation of transcription (for example in response to oxidative stress), response to hypoxia and toxic substance,
Cepba	regulates gluconeogenesis and lipogenesis by interacting with Foxo1 and Srebf1, liver development, cholesterol metabolic process, cytokine-mediated signaling pathway, mitochondrion organization, regulation of cell cycle / maturation / proliferation, regulation of transcription, Notch signaling pathway, urea cycle
Esrrb	regulation of transcription
Fos	ageing, signal transduction (e.g. SMAD protein signaling), cell proliferation and differentiation, regulation of transcription, transforming growth factor (TGF) beta receptor signaling pathway
Foxd3	regulation of transcription
Foxj1	activation of GTPase activity, cell differentiation, regulation of NF-kappaB transcription factor activity, regulation of transcription
Foxj3	cell differentiation, regulation of transcription
Foxl1	cell differentiation, regulation of transcription, Wnt signaling pathway
Hnf1a	transport of bile acid, bile salt, cholesterol (reverse) and fatty acids, bile acid, fatty acid and cholesterol biosynthetic process, glucose homeostasis, response to glucose and oxidative stress, liver development, regulation of transcription / histone acetylation, regulation of hormone secretion (e.g. insulin), Wnt and SMAD signaling pathway
Hoxa5	cell migration, regulation of apoptotic process, positive regulation of receptor biosynthetic process, regulation of transcription
Jdp2	regulation of transcription, positive regulation of histone deacetylation
Jun	ageing, circadian rhythm, liver development, regulation of apoptotic process, regulation of protein autophosphorylation, regulation of transcription (e.g. in response to endoplasmic reticulum stress), regulation of cell cycle / differentiation / proliferation, regulation of ERK1 and ERK2 cascade, regulation of GTPase activity, release of cytochrome c from mitochondria, SMAD protein signal transduction, TGF beta receptor signaling pathway
Klf4	Wnt signaling pathway, cellular response to growth factor stimulus, regulation of cell migration /differentiation / proliferation, regulation of apoptotic process, regulation of ERK1 and ERK2 cascade, negative regulation of NF-kappaB transcription factor activity, negative regulation of phosphatidylinositol 3-kinase signaling, negative regulation of protein kinase B signaling, negative regulation of response to cytokine stimulus, regulation of transcription, positive regulation of cellular protein metabolic process, positive regulation of nitric oxide biosynthetic process, positive regulation of telomerase activity
Klf7	regulation of transcription
Max	regulation of transcription
Myc	Wnt signaling pathway, regulation of apoptotic process, cell cycle arrest, cellular response to DNA damage stimulus /drugs / interferon-alpha, chromatin remodeling / chromosome organization / regulation of DNA biosynthetic process and telomere maintenance, MAPK cascade, regulation of gene expression and transcription, regulation of cell proliferation, protein processing
Mycn	regulation of ROS metabolic process, positive regulation of cell death and proliferation, regulation of transcription
Mzf1	regulation of transcription
Nfyα	regulation of transcription, regulation of circadian clock genes

Table 29/b: Overview on the transcription factors and their biological function

Nkx2-5	Wnt signaling pathway, cell differentiation /proliferation, apoptosis, regulation of gene expression and transcription, positive regulation of sodium ion transport
Nkx3-2	negative regulation of apoptotic process, regulation of transcription
Pparg	regulates beta-oxidation, regulation of apoptosis, signal transduction regulation of cell maturation / proliferation, cellular response to insulin stimulus, response to dietary excess, diet induced thermogenesis, glucose homeostasis, inflammatory response, lipoprotein and long-chain fatty acid transport, low-density lipoprotein particle receptor biosynthetic process, negative regulation of cholesterol storage, negative regulation of interferon-gamma-mediated signaling pathway, negative regulation of peptide hormone secretion, regulation of transcription, peroxisome proliferator activated receptor signaling pathway, regulation of circadian rhythm,
Rxra	ageing, cellular response to insulin stimulus, inflammatory response, liver development, regulation of cell proliferation, regulation of gene expression, transcription and translation, peroxisome proliferator activated receptor signaling pathway, regulation of apoptosis, retinoic acid receptor signaling pathway
Sp1	involved in cell growth, apoptosis, differentiation and immune responses , liver development, regulation of transcription, circadian rhythm
Spib	regulation of transcription
Sry	cell differentiation, regulation of gene expression and transcription
Tbp	regulation of transcription
Yy1	cell differentiation, cellular response to DNA damage stimulus, cellular response to interleukin-1, chromosome organization, regulation of gene expression and transcription
Zeb1	cellular response to amino acid stimulus, regulation of cell proliferation, regulation of transcription, regulation of TGF beta receptor signaling pathway
Zfp105	regulation of transcription
Zfx	regulation of transcription
Znf354c	regulation of transcription

TF = transcription factor

4. Discussion

This paper introduces a new approach to identify potential mimetics of CR. The advantages of this method are fast calculation, low cost and the ability to analyse several experiments together, which limits the occurrence of random results; in addition, no protracted *in vivo* or *in vitro* experiments are necessary. Furthermore, this method limits the risk of wasting time and research funds because it reduces the probability that an experiment does not show the desired result. Therefore, it allows for more purposeful work. The only condition of this approach is that gene expression data must be available. Databases like GEO and ArrayExpress are useful to find these data. All previous experiments can be included because no control group with a CR diet is necessary. The results of all available CR experiments can be summarised to get a control. For this, it is necessary to unify the different experiments. This could be a source of error because experimental design varies in the studies (e.g., in the number of animals and replicates, age, gender, genetic strain, degree of CR, concentration of phytonutrients, contribution of the standard diet and duration of the intervention). Furthermore, different microarrays are used, which can influence the results because each gene is not ana-

lysed on each array and because the different arrays (e.g., Affymetrix and Illumina) need different R packages for pre-processing and identifying the gene expression changes. It is tried to compensate for this by pre-processing the raw data using the quantile normalisation implemented in each R package. Another difficulty is that the genes are named differently on the microarray chips. It is tried to compensate for this by using the Entrez IDs, but errors in naming could not be ruled out. Moreover, each annotation update on a chip can change the results. Nonetheless, the mentioned approach is useful in getting a first indication about both the CR function and the overlap between CR and phytonutrients; this must be confirmed by further investigation.

To identify the best method for detecting significant expression changes, several methods are compared – the t-test, the moderated t-test, the Wilcoxon test and the rank product analysis in R. The t-test is the standard method of calculating expression changes, but the experiments' box plots show that the data are not normally distributed, so the Wilcoxon test should be used instead. However, our results indicate that the Wilcoxon test does not work with a small number of replicates. Consequently, no up- and downregulated genes other than CR can be identified with the Wilcoxon test. After combining the Wilcoxon test with the fold change, no genes were identified for CR. Therefore, the Wilcoxon test was useless in our analysis. The t-test and the moderated t-test show a larger number of regulated genes. However, one disadvantage of these methods is that each experiment was analysed by itself; the results had to be combined afterward with a one-sided binomial test and a chi-squared test. The chi-squared test also does not work with a small number of experiments. For example, if all three experiments show a significant upregulation, the chi-squared test leads to a non-significant result. Therefore, the '3-fold bigger' method was used instead. A few expression changes were identified, but after combining the t-test with a fold change, the number of regulated genes declined drastically. Moreover, one disadvantage of the t-test and moderated t-test is that they only reveal genes whose expressions have obviously changed. If the expression change is very low or the p-value is a little too large, the gene is not further mentioned. This can drastically influence the results of the overlapped analyses. This could be very important because the gene with the largest expression change does not always cause the effect *in vivo*. The rank product analysis avoids the disadvantages of the previous methods because the data for all experiments can be uploaded at the same time, and the expression changes of each experiment are ranked by size. Afterward, the positions of each gene in the rankings are multiplied, which leads to a general ranking over all the experiments. Therefore, this method includes the tendencies of expression changes and shows which genes are consistently up- or downregulated over all experiments – even if no significant p-value can be iden-

tified with a t-test. Thus, the rank product analysis seems to be the most useful method. The gene lists obtained for CR and the phytonutrients can be used to identify an overlap in gene expression or in function. Many tools are available for enrichment analysis, but it is decided to use CPDB because it includes the pathways of many known databases, such as KEGG, Reactome and MouseCyc. Nonetheless, other tools could provide other results.

As mentioned above, R is known as a potential CR mimetic (236). Our results indicate as well that R shows the largest overlap in gene expression and that therefore many CR functions could be mimicked. However, our results show too that each of the investigated phytonutrients regulates some genes in the same direction and that overlapped pathways could be identified in enrichment analysis. Further studies are necessary to see if this overlap is sufficient to mimic CR *in vivo*.

For a better assumption it could be useful to investigate the expression changes of genes which are known to be important for the beneficial effects of CR. Sirt1 is the most popular gene. In contrast to the results of past investigations, R shows no upregulation in our analysis, whereas Sirt1 expression is increased by Qd. Thus, the upregulation of Sirt1 indicates that Qd could be a CR mimetic even if only a small number of overlapped genes are identified.

The presumption that Qd is a CR mimetic is supported by the upregulation of Igfbp1 and the downregulation of Irs1 and Igfals. This indicates that Qd inhibits Igf1 signalling, which is associated with many beneficial CR effects (see above). The fact that CR downregulates Irs1 was shown by MA et al. (2014) in murine hippocampus too (237). In contrast, GESING et al. (2014) have shown that CR increases Irs1 expression in long-lived Ames dwarf (df/df) mice (238). The results of WANG et al. (2013) indicate that the effect of CR on Irs1 phosphorylation in skeletal muscle of humans is strongly dependent on the composition of the diet (fat content, carbohydrate content) (239). Moreover, the ability of CR to regulate phosphorylation of Irs1 is dependent on AMPK and mTOR signalling (240). To our knowledge, nothing is known about the influence of quercetin on the expression of these genes in murine liver, but quercetin ameliorates insulin resistance by beneficial regulation of Irs1 function in human umbilical vein endothelial cells (241). Furthermore, quercetin reduces Irs1 concentration in human keloid fibroblasts and in AT6.3 cells of rats (prostate adenocarcinoma cell line) (242; 243). This is associated with an inhibited Igf1-induced proliferation.

It is assumed that Qd and CR upregulate the expression of Igfbp1 by increasing the expression and/or activity of Foxo1 and Foxa1. PANOWSKI et al. (2007) have seen that the lifespan-expanding effect of CR in *C. elegans* is mediated by Pha-4 (ortholo-

gous to Foxa1, Foxa2 and Foxa3) (201). The authors suspect that the role of Pha-4 in longevity is specific for CR and that Pha-4 overexpression can increase the lifespan in *daf-16*(Foxo)-null mutant strains. Our results indicate that Qd probably addresses the same mechanism. Whether Foxa1 is important for lifespan extension in mice as well should be further analysed.

As mentioned above, Foxo1 is associated with oxidative stress. KIM et al. (2008) have shown that PI3K/Akt activation and Foxo1 phosphorylation increase with ageing, which leads to enhanced oxidative stress (244). Several previous investigations indicate that quercetin acts as an antioxidant and prevents oxidative stress (245; 246). Moreover, quercetin modulates PI3K/Akt activation, Foxo1 phosphorylation and mTOR signalling (247; 248; 249). This indicates that quercetin reduces oxidative stress by similar mechanisms to CR.

Whether Qd is also able to affect growth hormone signalling is not really clear. However, Qi decreases Ghr expression. This probably indicates that higher dosages of Q could be more effective in influencing the GH-Igf1 axis. As mentioned above, Ghr is discussed as the gene with the biggest impact on longevity (250). In summary, our results indicate that Qd could mimic CR effects by inhibiting GH/Igf1 signalling, increasing the activity of Foxo1 and increasing the expression of Sirt1 and Nnmt. Regarding its effect on longevity, it is found that quercetin increases lifespan in *C. elegans* and *Saccharomyces cerevisiae* (251; 252). To our knowledge, no experiments have analysed the effect of Q in mice. Whether quercetin is a CR mimetic should be further analysed *in vivo*.

Regarding the analysis of longevity genes, it is probable that each phytonutrient extends the lifespan because it is difficult to say how these genes interact and which gene triggers for the anti- or pro-longevity effect. P downregulates the expression of Ghr, which indicates that P probably increases longevity even though Sirt1 expression is decreased. P expands the lifespan of *Saccharomyces cerevisiae* dose-dependently (253). In contrast to our results, superoxide dismutases (Sod1 and Sod2) and Sirt1 seem to be involved in this effect. Whereas the expression of Sod1 and Sod2 is not influenced by P in our analysis, the expression of Sod3 is decreased. It is possible that these differences are attributable to the P dosage. This should be further investigated. Thus, it is impossible to say whether P increases lifespan in mice too.

For R, a lifespan-expanding effect is seen only in mice that are fed a high-calorie diet, not in mice that are fed a standard diet (46; 81). G is shown to promote lifespan in *C. elegans*, whereas in *D. melanogaster* there is an inverse correlation between G concentration and maximal lifespan (254; 255). In humans it is discussed too whether G

could promote longevity via its antioxidant effects and by activating the NF kappa B pathways (256; 257). In summary, our results indicate that all phytonutrients could be able to increase lifespan but via mechanisms other than CR.

Our results further indicate that CR addresses many more pathways than growth hormone and Igf1 signalling. Some of them are already mentioned in the context of CR (e.g., ABC transporters (258; 259; 260), cholesterol efflux (259; 261), bile metabolism (262; 261), the cholesterol metabolism (195), HDL and LDL uptake into hepatocytes via scavenger receptor class B, member 1 (Scarb1) and low-density lipoprotein receptor (Ldlr) (263; 264), the androgen metabolism and androgen receptor signalling (265; 266; 267), PPAR signalling pathway (268; 269; 270; 271; 272; 40), AMPK signalling (240), biological oxidation (10; 273; 274; 275; 276), the circadian rhythm (277; 278; 279), apoptosis (156; 280; 281), expression of ribosomal subunits (282), phago-some/autophagy/mitophagy mechanisms (283; 284; 285; 286; 32), processing of mRNA (including splicing) (178), gene expression/translation regulation (287), lipid metabolism (288; 274; 289; 290; 19; 291; 292), carbohydrate/pyruvate metabolism (293; 269; 294; 295), the citric acid cycle (296; 297), the electron transport chain (293; 296), the pentose phosphate pathway (298; 299; 300; 301), the biotin metabolism (302; 303; 304), the regulation of immune system/inflammation/oxidative stress (305; 306; 307; 308; 309; 310), Interleukin-6 signalling (311; 312), epidermal growth factor receptor (EGFR1) signalling (313; 314), amino acid metabolism, protein synthesis/processing/turnover, and ubiquitin-dependent protein degradation (315; 293; 294; 316; 295; 317; 318; 319)). However, others have not been revealed in the context of CR before, such as the influence on oestrogen metabolism, retinol metabolism, the urea cycle, glutathione metabolism including the γ -glutamyl cycle, starch and sucrose metabolism, vesicle-mediated transport, selenium metabolism, heme biosynthesis/metabolism of porphyrins, interferon signalling, toll-like receptor signalling, complement and coagulation cascade, and the expression of metal ion SLC transporters (especially of zinc transporters). It is necessary to confirm whether CR influences these pathways *in vivo* as well because they could lead to a better understanding of the mechanisms of CR. For example, it is known that all-trans-retinoic acid (atRA) is involved in bile acid homeostasis (320). Therefore, it is possible that the influence of CR on retinol metabolism is involved in the regulation of cholesterol and bile acid. The influence in zinc SLC transporters is very interesting too. It is known that zinc is involved in immune function, apoptosis and insulin resistance (321; 322; 323; 324). Furthermore, some sirtuins are zinc-dependent (325). CABREIRO et al. (2008) have shown that ageing is associated with zinc deficiency, which leads to higher oxidative stress (326). All of these functions have already been discussed in the context of CR as well

(see above). This leads to the hypothesis that the influence of CR on zinc transporters or zinc metabolism could be involved in the beneficial effects of CR. This hypothesis is supported by the upregulation of Mt1 by CR, which is important for cellular zinc ion homeostasis. Unfortunately, no phytonutrient is able to mimic the effect on SLC transporters, but R and Qd upregulate the expression of Mt1 too. Whether zinc is important for beneficial CR effects should be further analysed.

Nonetheless, all phytonutrients are partly able to mimic the effect of CR, although Qd and R seem to be the most useful. For a better determination of which phytonutrient is the best mimetic, it is necessary to understand in detail how CR affects the lifespan and health. Limiting the analysis on these pathways and the genes included in them will be more effective for identifying mimetics.

5. Conclusion

This work shows a new approach to identify CR mimetics in *silico*. The results show that all phytonutrients could be partly able to mimic CR but that R shows the largest overlap in gene expression. Nonetheless, it is hypothesised that Qd could be a CR mimetic as well, although only a small number of overlapping genes and pathways could be identified. However, Qd was shown to upregulate the expression of Sirt1 and Nnmt and probably to inhibit Igf1 and mTOR signalling and increase Foxo1 activity. For a better assessment of whether a phytonutrient could be a CR mimetic, it is necessary to understand all mechanisms that are important for the beneficial health effects of CR. In *silico* analysis could also be useful for finding them. For example, our analysis indicates that zinc ion homeostasis could be a new mechanism that explains many CR functions. R and Qd upregulate the expression of Mt1 as well, which indicates again that both are potential CR mimetics. It is impossible to say whether other phytonutrients are more effective than R or Qd to mimic CR because only datasets from a few phytonutrients are available. Therefore, it should be a goal to motivate scientists to upload their data to make in *silico* analyses more useful. Furthermore, it is important that datasets of different plant-based diets with varying concentrations of phytonutrients are available because it is impossible to say how the phytonutrients interact in vivo.

To carry out in *silico* analysis and make it easier to find CR mimetics, a database similar to a connectivity map could be useful; this could summarise the influences of different substances (e.g., phytonutrients, vitamins and drugs) on the expression of one gene in different organisms. This could be handy to answer questions such as ‘which substances upregulate Sirt1 in murine liver?’ or ‘which substance is available to regulate a group of genes in the same direction like CR?’. The mentioned substances could further be analysed in vitro and in vivo, which would allow identifying more purposefully

potential CR mimetics. To develop this database, it would be necessary for all authors to upload their results, for example on ArrayExpress or GEO or directly to the newly developed database. Therefore, further effort should be expended to motivate scientists to upload their data and to establish an organisation that summarises the results and develop this database. Furthermore, it is important to find or develop a tool for enrichment analysis which could be used as gold standard. This would facilitate comparison of the results of different studies.

Despite all the difficulties which limited the results of our analysis, the mentioned approach gives a first insight to the overlapped functions between CR and the phytonutrients and could be useful to find potential CR mimetics.

References

1. Ribeiro AI, Krainski ET, Carvalho MS, Pina MF. Where do people live longer and shorter lives? An ecological study of old-age survival across 4404 small areas from 18 European countries. *J Epidemiol Community Health.* 2016 Feb 15. pii: jech-2015-206827.
2. Le Couteur DG, Solon-Biet S, Cogger VC, Mitchell SJ, Senior A, de Cabo R, Raubenheimer D, Simpson SJ. The impact of low-protein high-carbohydrate diets on aging and lifespan. *Cell Mol Life Sci.* 2015 Dec 30. [Epub ahead of print].
3. D, Schafer. Aging, longevity, and diet: historical remarks on calorie intake reduction. *Gerontology.* 2005 Mar-Apr;51(2):126-30.
4. Speakman JR, Mitchell SE, Mazidi M. Calories or Protein? The effect of dietary restriction on lifespan in rodents is explained by calories alone. *Exp Gerontol.* 2016 Mar 19. pii: S0531-5565(16)30069-9.
5. Osborne TB, Mendel LB, Ferry EL. THE EFFECT OF RETARDATION OF GROWTH UPON THE BREEDING PERIOD AND DURATION OF LIFE OF RATS. *Science.* 1917 Mar 23;45(1160):294-5.
6. McCay C, Crowell M, Maynard L. The effect of retarded growth upon the length of life and upon ultimate size. *J Nutr.* 1935; 10:63–79.
7. Zimmer Z, Rubin S. Life Expectancy With and Without Pain in the U.S. Elderly Population. *J Gerontol A Biol Sci Med Sci.* 2016 Mar 17. pii: glw028.
8. Sasaki T, Izawa M, Okada Y. Current Trends in Health Insurance Systems: OECD Countries vs. Japan. *Neurol Med Chir (Tokyo).* 2015;55 Suppl 1:267-75.
9. Kelley AS, McGarry K, Gorges R, Skinner JS. The burden of health care costs for patients with dementia in the last 5 years of life. *Ann Intern Med.* 2015 Nov 17;163(10):729-36.
10. Fu ZD, Klaassen CD. Short-term calorie restriction feminizes the mRNA profiles of drug metabolizing enzymes and transporters in livers of mice. *Toxicol Appl Pharmacol.* 2014 Jan 1;274(1):137-46.
11. Barger JL, Kayo T, Vann JM, Arias EB, Wang J, Hacker TA, Wang Y, Raederstorff D, Morrow JD, Leeuwenburgh C, Allison DB, Saupe KW, Cartee GD, Weindruch R, Prolla TA. A low dose of dietary resveratrol partially mimics caloric restriction and retards aging parameters in mice. *PLoS One.* 2008 Jun 4;3(6):e2264.
12. López-Lluch G, Navas P. Calorie restriction as an intervention in ageing. *J Physiol.* 2015 Nov 26.

13. Si H, Liu D. Dietary antiaging phytochemicals and mechanisms associated with prolonged survival. *J Nutr Biochem*. Jun 2014, pp. 25(6):581-91.
14. Lee C, Longo V. Dietary restriction with and without caloric restriction for healthy aging. *F1000Res*. 2016 Jan 29;5.
15. AR, Hipkiss. Does chronic glycolysis accelerate aging? Could this explain how dietary restriction works? *Ann N Y Acad Sci*. 2006 May;1067:361-8.
16. Poljsak B, Milisav I. The NAD(+) depletion theory of ageing: NAD(+) as the link between oxidative stress, inflammation, caloric restriction, exercise, DNA repair, longevity and health span. *Rejuvenation Res*. 2016 Jan 4. [Epub ahead of print].
17. DA, Sinclair. Toward a unified theory of caloric restriction and longevity regulation. *Mech Ageing Dev*. 2005 Sep;126(9):987-1002.
18. C, Handschin. Caloric restriction and exercise "mimetics": Ready for prime time? *Pharmacol Res*. 2015 Dec 2;103:158-166.
19. Okita N, Tsuchiya T, Fukushima M, Itakura K, Yuguchi K, Narita T, Hashizume Y, Sudo Y, Chiba T, Shimokawa I, Higami Y. Chronological analysis of caloric restriction-induced alteration of fatty acid biosynthesis in white adipose tissue of rats. *Exp Gerontol*. 2015 Jan 20.
20. Stenvinkel P, Kooman JP, Shiels PG. Nutrients and ageing: what can we learn about ageing interactions from animal biology? *Curr Opin Clin Nutr Metab Care*. 2016 Jan;19(1):19-25.
21. Xu X, Sonntag WE. Growth hormone-induced nuclear translocation of Stat-3 decreases with age: modulation by caloric restriction. *Am J Physiol*. 1996 Nov;271(5 Pt 1):E903-9.
22. Yang L, Licastro D, Cava E, Veronese N, Spelta F, Rizza W, Bertozzi B, Villareal DT, Hotamisligil GS, Holloszy JO, Fontana L. Long-Term Calorie Restriction Enhances Cellular Quality-Control Processes in Human Skeletal Muscle. *Cell Rep*. 2016 Jan 26;14(3):422-8.
23. Balasubramanian P, Longo VD. Growth factors, aging and age-related diseases. *Growth Horm IGF Res*. 2016 Jan 7. pii: S1096-6374(16)30001-6.
24. Babar P, Adamson C, Walker GA, Walker DW, Lithgow GJ. P13-kinase inhibition induces dauer formation, thermotolerance and longevity in *C. elegans*. *Neurobiol Aging*. 1999 Sep-Oct;20(5):513-9.

25. Calvert S, Tacutu R, Sharifi S, Teixeira R, Ghosh P, de Magalhães JP. A network pharmacology approach reveals new candidate caloric restriction mimetics in *C. elegans*. *Aging Cell*. 2015 Dec 16.
26. Furukawa-Hibi Y, Kobayashi Y, Chen C, Motoyama N. FOXO transcription factors in cell-cycle regulation and the response to oxidative stress. *Antioxid Redox Signal*. 2005 May-Jun;7(5-6):752-60.
27. Gross DN, Wan M, Birnbaum MJ. The role of FOXO in the regulation of metabolism. *Curr Diab Rep*. 2009 Jun;9(3):208-14.
28. Mitchell SE, Delville C, Konstantopoulos P, Hurst J, Derous D, Green C, Chen L, Han JJ, Wang Y, Promislow DE, Lusseau D, Douglas A, Speakman JR. The effects of graded levels of calorie restriction: II. Impact of short term calorie and protein restriction on circulating hormone levels, glucose homeostasis and oxidative stress in male C57BL/6 mice. *Oncotarget*. 2015 Sep 15;6(27):23213-37.
29. CJ, Kenyon. The genetics of ageing. *Nature*. 2010 Mar 25;464(7288):504-12.
30. VN, Anisimov. Conservative growth hormone/IGF-1 and mTOR signaling pathways as a target for aging and cancer prevention: do we really have an antiaging drug? *Interdiscip Top Gerontol*. 2015;40:177-88.
31. Kume S, Koya D. Autophagy: A Novel Therapeutic Target for Diabetic Nephropathy. *Diabetes Metab J*. 2015 Dec;39(6):451-60.
32. Morselli E, Maiuri MC, Markaki M, Megalou E, Pasparaki A, Palikaras K, Criollo A, Galluzzi L, Malik SA, Vitale I, Michaud M, Madeo F, Tavernarakis N, Kroemer G. Caloric restriction and resveratrol promote longevity through the Sirtuin-1-dependent induction of autophagy. *Cell Death Dis*. 2010;1:e10.
33. Dai DF, Karunadharma PP, Chiao YA, Basisty N, Crispin D, Hsieh EJ, Chen T, Gu H, Djukovic D, Raftery D, Beyer RP, MacCoss MJ, Rabinovitch PS. Altered proteome turnover and remodeling by short-term caloric restriction or rapamycin rejuvenate the aging heart. *Aging Cell*. 2014 Jun;13(3):529-39.
34. BL, Tang. Sirt1 and the Mitochondria. *Mol Cells*. 2016 Feb 29;39(2):87-95.
35. Schmeisser K, Mansfeld J, Kuhlow D, Weimer S, Priebe S, Heiland I, Birringer M, Groth M, Segref A, Kanfi Y, Price NL, Schmeisser S, Schuster S, Pfeiffer AF, Guthke R, Platzer M, Hoppe T, Cohen HY, Zarse K, Sinclair DA, Ristow M. Role of sirtuins in lifespan regulation is linked to methylation of nicotinamide. *Nat Chem Biol*. 2013 Nov;9(11):693-700.

36. Covington JD, Bajpeyi S. The sirtuins: Markers of metabolic health. *Mol Nutr Food Res.* 2016 Jan;60(1):79-91.
37. Dikalov SI, Dikalova AE. Contribution of mitochondrial oxidative stress to hypertension. *Curr Opin Nephrol Hypertens.* 2015 Dec 29. [Epub ahead of print].
38. Wątroba M, Szukiewicz D. The role of sirtuins in aging and age-related diseases. *Adv Med Sci.* 2016 Mar;61(1):52-62.
39. Schwarz F, Karadeniz Z, Fischer-Rosinsky A, Willmes DM, Spranger J, Birkenfeld AL. Knockdown of Indy/CeNac2 extends *Caenorhabditis elegans* life span by inducing AMPK/aak-2. *Aging (Albany NY).* 2015 Aug;7(8):553-67.
40. Nisoli E, Tonello C, Cardile A, Cozzi V, Bracale R, Tedesco L, Falcone S, Valerio A, Cantoni O, Clementi E, Moncada S, Carruba MO. Calorie restriction promotes mitochondrial biogenesis by inducing the expression of eNOS. *Science.* 2005 Oct 14;310(5746):314-7.
41. Hancock CR, Han DH, Higashida K, Kim SH, Holloszy JO. Does calorie restriction induce mitochondrial biogenesis? A reevaluation. *FASEB J.* 2011 Feb;25(2):785-91.
42. Van Meter M, Mao Z, Gorbunova V, Seluanov A. SIRT6 overexpression induces massive apoptosis in cancer cells but not in normal cells. *Cell Cycle.* 2011 Sep 15;10(18):3153-8.
43. Sundaresan NR, Vasudevan P, Zhong L, Kim G, Samant S, Parekh V, Pillai VB, Ravindra PV, Gupta M, Jeevanandam V, Cunningham JM, Deng CX, Lombard DB, Mostoslavsky R, Gupta MP. The sirtuin SIRT6 blocks IGF-Akt signaling and development of cardiac hypertrophy by targeting c-Jun. *Nat Med.* 2012 Nov;18(11):1643-50.
44. Kanfi Y, Naiman S, Amir G, Peshti V, Zinman G, Nahum L, Bar-Joseph Z, Cohen HY. The sirtuin SIRT6 regulates lifespan in male mice. *Nature.* 2012 Feb 22;483(7388):218-21.
45. Kanfi Y, Shalman R, Peshti V, Pilosof SN, Gozlan YM, Pearson KJ, Lerrer B, Moazed D, Marine JC, de Cabo R, Cohen HY. Regulation of SIRT6 protein levels by nutrient availability. *FEBS Lett.* 2008 Mar 5;582(5):543-8.
46. Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, Kalra A, Prabhu VV, Allard JS, Lopez-Lluch G, Lewis K, Pistell PJ, Poosala S, Becker KG, Boss O, Gwinn D, Wang M, Ramaswamy S, Fishbein KW, Spencer RG, Lakatta EG, Le Couteur D, Shaw RJ, Navas P, Puigserver P, Ingram DK, de Cabo R, Sinclair DA. Resveratrol improves

- health and survival of mice on a high-calorie diet. *Nature*. 2006 Nov 16;444(7117):337-42.
47. Ingram DK, Roth GS. Calorie restriction mimetics: can you have your cake and eat it, too? *Ageing Res Rev*. 2015 Mar;20:46-62.
48. Kim N, Lee JO, Lee HJ, Lee YW, Kim HI, Kim SJ, Park SH, Lee CS, Ryoo SW, Hwang GS, Kim HS. AMPK, a metabolic sensor, is involved in isoeugenol-induced glucose uptake in muscle cells. *J Endocrinol*. 2016 Feb;228(2):105-14.
49. Apfeld J, O'Connor G, McDonagh T, DiStefano PS, Curtis R. The AMP-activated protein kinase AAK-2 links energy levels and insulin-like signals to lifespan in *C. elegans*. *Genes Dev*. 2004 Dec 15;18(24):3004-9.
50. Harkness TA, Shea KA, Legrand C, Brahmania M, Davies GF. A functional analysis reveals dependence on the anaphase-promoting complex for prolonged life span in yeast. *Genetics*. 2004 Oct;168(2):759-74.
51. Tohyama D, Yamaguchi A. A critical role of SNF1A/dAMPKalpha (*Drosophila* AMP-activated protein kinase alpha) in muscle on longevity and stress resistance in *Drosophila melanogaster*. *Biochem Biophys Res Commun*. 2010 Mar 26;394(1):112-8.
52. Cai Y, Wei YH. Stress resistance and lifespan are increased in *C. elegans* but decreased in *S. cerevisiae* by mafr-1/maf1 deletion. *Oncotarget*. 2016 Feb 26. [Epub ahead of print].
53. Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, Nadon NL, Wilkinson JE, Frenkel K, Carter CS, Pahor M, Javors MA, Fernandez E, Miller RA. Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature*. 2009 Jul 16;460(7253):392-5.
54. Snell TW, Johnston RK. Glycerol extends lifespan of *Brachionus manjavacas* (Rotifera) and protects against stressors. *Exp Gerontol*. 2014 Sep;57:47-56.
55. Weimer S, Priebs J, Kuhlow D, Groth M, Priebe S, Mansfeld J, Merry TL, Dubuis S, Laube B, Pfeiffer AF, Schulz TJ, Guthke R, Platzer M, Zamboni N, Zarse K, Ristow M. D-Glucosamine supplementation extends life span of nematodes and of ageing mice. *Nat Commun*. 2014 Apr 8;5:3563.
56. Yuan Y, Hakimi P, Kao C, Kao A, Liu R, Janocha A, Boyd-Tressler A, Hang X, Alhoraibi H, Slater E, Xia K, Cao P, Shue Q, Ching TT, Hsu AL, Erzurum SC, Dubyak GR, Berger NA, Hanson RW, Feng Z. Reciprocal Changes in Phosphoenolpyruvate Carboxykinase and Pyruvate Kinase with Age Are a Determinant of Aging in *Caenorhabditis elegans*. *J Biol Chem*. 2016 Jan 15;291(3):1307-19.

57. Shih DM, Wang Z, Lee R, Meng Y, Che N, Charugundla S, Qi H, Wu J, Pan C, Brown JM, Vallim T, Bennett BJ, Graham M, Hazen SL, Lusis AJ. Flavin containing monooxygenase 3 exerts broad effects on glucose and lipid metabolism and atherosclerosis. *J Lipid Res.* 2015 Jan;56(1):22-37.
58. Jiang Y, Jin J, Iakova P, Hernandez JC, Jawanmardi N, Sullivan E, Guo GL, Timchenko NA, Darlington GJ. Farnesoid X receptor directly regulates xenobiotic detoxification genes in the long-lived Little mice. *Mech Ageing Dev.* 2013 Sep;134(9):407-15.
59. Derois D, Mitchell SE, Green CL, Chen L, Han JJ, Wang Y, Promislow DE, Lusseau D, Speakman JR, Douglas A. The effects of graded levels of calorie restriction: VI. Impact of short-term graded calorie restriction on transcriptomic responses of the hypothalamic hunger and circadian signaling pathways. *Aging (Albany NY).* 2016 Feb 23. [Epub ahead of print].
60. Dirks AJ, Leeuwenburgh C. Caloric restriction in humans: potential pitfalls and health concerns. *Mech Ageing Dev.* 2006 Jan;127(1):1-7.
61. Nissim I, Horyn O, Daikhin Y, Chen P, Li C, Wehrli SL, Nissim I, Yudkoff M. The molecular and metabolic influence of long term agmatine consumption. *J Biol Chem.* 2014 Apr 4;289(14):9710-29.
62. Stout MB, Steyn FJ, Jurczak MJ, Camporez JG, Zhu Y, Hawse JR, Jurk D, Palmer AK, Xu M, Pirtskhalava T, Evans GL, de Souza Santos R, Frank AP, White TA, Monroe DG, Singh RJ, Casaclang-Verzosa G, Miller JD, Clegg DJ, LeBrasseur NK, von Zglinicki T, and Shulman GI, Tchkonia T, Kirkland JL. 17 α -Estradiol Alleviates Age-related Metabolic and Inflammatory Dysfunction in Male Mice Without Inducing Feminization. *J Gerontol A Biol Sci Med Sci.* 2016 Jan 24. pii: gtv309. [Epub ahead of print].
63. Bakhtiari N, Hosseinkhani S, Soleimani M, Hemmati R, Noori-Zadeh A, Javan M, Tashakor A. Short-term ursolic acid promotes skeletal muscle rejuvenation through enhancing of SIRT1 expression and satellite cells proliferation. *Biomed Pharmacother.* 2016 Mar;78:185-96.
64. Pocobelli G, Kristal AR, Patterson RE, Potter JD, Lampe JW, Kolar A, Evans I, White E. Total mortality risk in relation to use of less-common dietary supplements. *Am J Clin Nutr.* 2010 Jun;91(6):1791-800.
65. Bell GA, Kantor ED, Lampe JW, Shen DD, White E. Use of glucosamine and chondroitin in relation to mortality. *Eur J Epidemiol.* 2012 Aug;27(8):593-603.

66. Miller RA, Harrison DE, Astle CM, Baur JA, Boyd AR, de Cabo R, Fernandez E, Flurkey K, Javors MA, Nelson JF, Orihuela CJ, Pletcher S, Sharp ZD, Sinclair D, Starnes JW, Wilkinson JE, Nadon NL, Strong R. Rapamycin, but not resveratrol or simvastatin, extends life span of genetically heterogeneous mice. *J Gerontol A Biol Sci Med Sci*. 2011 Feb;66(2):191-201.
67. Miller RA, Harrison DE, Astle CM, Fernandez E, Flurkey K, Han M, Javors MA, Li X, Nadon NL, Nelson JF, Pletcher S, Salmon AB, Sharp ZD, Van Roekel S, Winkleman L, Strong R. Rapamycin-mediated lifespan increase in mice is dose and sex dependent and metabolically distinct from dietary restriction. *Aging Cell*. 2014 Jun;13(3):468-77.
68. Martin-Montalvo A, Mercken EM, Mitchell SJ, Palacios HH, Mote PL, Scheibye-Knudsen M, Gomes AP, Ward TM, Minor RK, Blouin MJ, Schwab M, Pollak M, Zhang Y, Yu Y, Becker KG, Bohr VA, Ingram DK, Sinclair DA, Wolf NS, Spindler SR, Bernier M, de Cabo R. Metformin improves healthspan and lifespan in mice. *Nat Commun*. 2013;4:2192.
69. Song YM, Lee WK, Lee YH, Kang ES, Cha BS, Lee BW. Metformin Restores Parkin-Mediated Mitophagy, Suppressed by Cytosolic p53. *Int J Mol Sci*. 2016 Jan 16;17(1).
70. Boussageon R, Supper I, Bejan-Angoulvant T, Kellou N, Cucherat M, Boissel JP, Kassai B, Moreau A, Gueyffier F, Cornu C. Reappraisal of metformin efficacy in the treatment of type 2 diabetes: a meta-analysis of randomised controlled trials. *PLoS Med*. 2012;9(4):e1001204.
71. Witte AV, Kerti L, Margulies DS, Flöel A. Effects of resveratrol on memory performance, hippocampal functional connectivity, and glucose metabolism in healthy older adults. *J Neurosci*. 2014 Jun 4;34(23):7862-70.
72. Lagouge M, Argmann C, Gerhart-Hines Z, Meziane H, Lerin C, Daussin F, Messadeq N, Milne J, Lambert P, Elliott P, Geny B, Laakso M, Puigserver P, Auwerx J. Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1alpha. *Cell*. 2006 Dec 15;127(6):1109-22.
73. Mercken EM, Carboneau BA, Krzysik-Walker SM, de Cabo R. Of mice and men: the benefits of caloric restriction, exercise, and mimetics. *Ageing Res Rev*. 2012 Jul;11(3):390-8.
74. Gruber J, Tang SY, Halliwell B. Evidence for a trade-off between survival and fitness caused by resveratrol treatment of *Caenorhabditis elegans*. *Ann N Y Acad Sci*. Apr;1100:530-42, 2007.

75. Bauer JH, Goupil S, Garber GB, Helfand SL. An accelerated assay for the identification of lifespan-extending interventions in *Drosophila melanogaster*. *Proc Natl Acad Sci U S A*. Aug 31;101(35):12980-5, 2004.
76. Valenzano DR, Terzibasi E, Genade T, Cattaneo A, Domenici L, Cellerino A. Resveratrol prolongs lifespan and retards the onset of age-related markers in a short-lived vertebrate. *Curr Biol*. Feb 7;16(3):296-300, 2006.
77. Wood JG, Rogina B, Lavu S, Howitz K, Helfand SL, Tatar M, Sinclair D. Sirtuin activators mimic caloric restriction and delay ageing in metazoans. *Nature*. 2004 Aug 5;430(7000):686-9.
78. Hu P, Zhao L, Chen J. Physiologically achievable doses of resveratrol enhance 3T3-L1 adipocyte differentiation. *Eur J Nutr*. 2014 Jul 17.
79. Kulkarni SS, Cantó C. The molecular targets of resveratrol. *Biochim Biophys Acta*. 2014 Oct 12; S0925-4439(14)00311-1.
80. Howitz KT, Bitterman KJ, Cohen HY, Lamming DW, Lavu S, Wood JG, Zipkin RE, Chung P, Kisielewski A, Zhang LL, Scherer B, Sinclair DA. Small molecule activators of sirtuins extend *Saccharomyces cerevisiae* lifespan. *Nature*. 2003 Sep 11;425(6954):191-6.
81. Pearson KJ, Baur JA, Lewis KN, Peshkin L, Price NL, Labinsky N, Swindell WR, Kamara D, Minor RK, Perez E, Jamieson HA, Zhang Y, Dunn SR, Sharma K, Pleshko N, Woollett LA, Csiszar A, Ikeno Y, Le Couteur D, Elliott PJ, Becker KG, Navas P, Ingram DK, Wolf NS, Ungvari Z, Sinclair DA, de Cabo R. Resveratrol delays age-related deterioration and mimics transcriptional aspects of dietary restriction without extending life span. *Cell Metab*. 2008 Aug;8(2):157-68.
82. Strong R, Miller RA, Astle CM, Baur JA, de Cabo R, Fernandez E, Guo W, Javors M, Kirkland JL, Nelson JF, Sinclair DA, Teter B, Williams D, Zaveri N, Nadon NL, Harrison DE. Evaluation of resveratrol, green tea extract, curcumin, oxaloacetic acid, and medium-chain triglyceride oil on life span of genetically heterogeneous mice. *J Gerontol A Biol Sci Med Sci*. 2013 Jan;68(1):6-16.
83. Lee SH, An HS, Jung YW, Lee EJ, Lee HY, Choi ES, An SW, Son H, Lee SJ, Kim JB, Min KJ. Korean mistletoe (*Viscum album coloratum*) extract extends the lifespan of nematodes and fruit flies. *Biogerontology*. 2014 Apr;15(2):153-64.
84. Huang CH, Tsai SJ, Wang YJ, Pan MH, Kao JY, Way TD. EGCG inhibits protein synthesis, lipogenesis, and cell cycle progression through activation of AMPK in p53

positive and negative human hepatoma cells. *Mol Nutr Food Res.* 2009 Sep;53(9):1156-65.

85. Gene Age: The Aging Gene Database. [Online] [Cited: 01 23, 2015.] <http://genomics.senescence.info/genes/>.

86. Rogina B, Helfand SL. Indy mutations and Drosophila longevity. *Front Genet.* 2013 Apr 8;4:47.

87. Huang da W, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 2009 Jan;37(1):1-13.

88. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. . *Nat Genet.* 2000 May;25(1):25-9.

89. Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc.* 2009;4(1):44-57.

90. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A.* 2005 Oct 25;102(43):15545-50.

91. Maere S, Heymans K, Kuiper M. BiNGO: a Cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks. *Bioinformatics.* 2005 Aug 15;21(16):3448-9.

92. Kamburov A, Pentchev K, Galicka H, Wierling C, Lehrach H, Herwig R. ConsensusPathDB: toward a more complete picture of cell biology. *Nucleic Acids Res.* 2011 Jan;39(Database issue):D712-7.

93. Kamburov A, Wierling C, Lehrach H, Herwig R. ConsensusPathDB--a database for integrating human functional interaction networks. *Nucleic Acids Res.* 2009 Jan;37(Database issue):D623-8.

94. Mi H, Muruganujan A, Thomas PD. PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucleic Acids Res.* 2013 Jan;41(Database issue):D377-86.

95. Harris MA, Clark J, Ireland A, Lomax J, Ashburner M, Foulger R, Eilbeck K, Lewis S, Marshall B, Mungall C, Richter J, Rubin GM, Blake JA, Bult C, Dolan M, Drabkin H,

- Eppig JT, Hill DP, Ni L, Ringwald M, Balakrishnan R, Cherry JM, Christie KR, Costanzo MC, Dwight SS, Engel S, Fisk DG, Hirschman JE, Hong EL, Nash RS, Sethuraman A, Theesfeld CL, Botstein D, Dolinski K, Feierbach B, Berardini T, Mundodi S, Rhee SY, Apweiler R, Barrell D, Camon E, Dimmer E, Lee V, Chisholm R, Gaudet P, Kibbe W, Kishore R, Schwarz EM, Sternberg P, Gwinn M, Hannick L, Wortman J, Berriman M, Wood V, de la Cruz N, Tonellato P, Jaiswal P, Seigfried T, White R and Consortium, Gene Ontology. The Gene Ontology (GO) database and informatics resource. *Nucleic Acids Res.* 2004 Jan 1;32(Database issue):D258-61.
96. Bader GD, Betel D, Hogue CW. BIND: the Biomolecular Interaction Network Database. *Nucleic Acids Res.* 2003 Jan 1;31(1):248-50.
97. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* 2000 Jan 1;28(1):27-30.
98. Croft D, Mundo AF, Haw R, Milacic M, Weiser J, Wu G, Caudy M, Garapati P, Gillespie M, Kamdar MR, Jassal B, Jupe S, Matthews L, May B, Palatnik S, Rothfels K, Shamovsky V, Song H, Williams M, Birney E, Hermjakob H, Stein L, D'Eustachio P. The Reactome pathway knowledgebase. *Nucleic Acids Res.* 2014 Jan;42(Database issue):D472-7.
99. Wingender E. The TRANSFAC project as an example of framework technology that supports the analysis of genomic regulation. *Brief Bioinform.* 2008 Jul;9(4):326-32.
100. Mathelier A, Fornes O, Arenillas DJ, Chen CY, Denay G, Lee J, Shi W, Shyr C, Tan G, Worsley-Hunt R, Zhang AW, Parcy F, Lenhard B, Sandelin A, Wasserman WW. JASPAR 2016: a major expansion and update of the open-access database of transcription factor binding profiles. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D110-5.
101. Hamosh A, Scott AF, Amberger JS, Bocchini CA, McKusick VA. Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders. *Nucleic Acids Res.* 2005 Jan 1;33(Database issue):D514-7.
102. Liu G, Loraine AE, Shigeta R, Cline M, Cheng J, Valmeekam V, Sun S, Kulp D, Siani-Rose MA. NetAffx: Affymetrix probesets and annotations. *Nucleic Acids Res.* 2003 Jan 1;31(1):82-6.
103. Maglott D, Ostell J, Pruitt KD, Tatusova T. Entrez Gene: gene-centered information at NCBI. *Nucleic Acids Res.* 2007 Jan;35(Database issue):D26-31.
104. Consortium, UniProt. The universal protein resource (UniProt). *Nucleic Acids Res.* 2008 Jan;36(Database issue):D190-5.

105. Wu CH, Yeh LS, Huang H, Arminski L, Castro-Alvear J, Chen Y, Hu Z, Kourtesis P, Ledley RS, Suzek BE, Vinayaka CR, Zhang J, Barker WC. The Protein Information Resource. *Nucleic Acids Res.* 2003 Jan 1;31(1):345-7.
106. Alibés A, Yankilevich P, Cañada A, Díaz-Uriarte R. IDconverter and IDClight: conversion and annotation of gene and protein IDs. *BMC Bioinformatics*. 2007 Jan 10;8:9.
107. Huang DW, Sherman BT, Tan Q, Kir J, Liu D, Bryant D, Guo Y, Stephens R, Baseler MW, Lane HC, Lempicki RA. DAVID Bioinformatics Resources: expanded annotation database and novel algorithms to better extract biology from large gene lists. *Nucleic Acids Res.* 2007 Jul;35(Web Server issue):W169-75.
108. Berriz GF, Roth FP. The Synergizer service for translating gene, protein and other biological identifiers. *Bioinformatics*. 2008 Oct 1;24(19):2272-3.
109. Fortney K, Morgen EK, Kotlyar M, Jurisica I. In silico drug screen in mouse liver identifies candidate calorie restriction mimetics. *Rejuvenation Res.* 2012 Apr;15(2):148-52.
110. Kolesnikov N, Hastings E, Keays M, Melnichuk O, Tang YA, Williams E, Dylag M, Kurbatova N, Brandizi M, Burdett T, Megy K, Pilicheva E, Rustici G, Tikhonov A, Parkinson H, Petryszak R, Sarkans U, Brazma A. ArrayExpress update--simplifying data submissions. *Nucleic Acids Res.* 2015 Jan;43(Database issue):D1113-6.
111. Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA, Phillippy KH, Sherman PM, Holko M, Yefanov A, Lee H, Zhang N, Robertson CL, Serova N, Davis S, Soboleva A. NCBI GEO: archive for functional genomics data sets--update. *Nucleic Acids Res.* 2013 Jan;41(Database issue):D991-5.
112. Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res.* 2002 Jan 1;30(1):207-10.
113. Breitling R, Armengaud P, Amtmann A, Herzyk P. Rank products: a simple, yet powerful, new method to detect differentially regulated genes in replicated microarray experiments. *FEBS Lett.* 2004 Aug 27;573(1-3):83-92.
114. Alissa EM, Ferns GA. Dietary Fruits and Vegetables and Cardiovascular Diseases Risk. *Crit Rev Food Sci Nutr.* 2015 Jul 20:0. [Epub ahead of print].
115. Salas-Salvadó J, Guasch-Ferré M, Lee CH, Estruch R, Clish CB, Ros E. Protective Effects of the Mediterranean Diet on Type 2 Diabetes and Metabolic Syndrome. *J Nutr.* 2016 Mar 9. pii: jn218487.

116. National Center for Biotechnology Information - Gene. [Online] <http://www.ncbi.nlm.nih.gov/gene>.
117. Gautier L, Cope L, Bolstad BM, Irizarry RA. affy--analysis of Affymetrix GeneChip data at the probe level. *Bioinformatics*. 2004 Feb 12;20(3):307-15.
118. Carvalho BS, Irizarry RA. A framework for oligonucleotide microarray preprocessing. *Bioinformatics*. 2010 Oct 1;26(19):2363-7.
119. Irizarry RA, Bolstad BM, Collin F, Cope LM, Hobbs B, Speed TP. Summaries of Affymetrix GeneChip probe level data. *Nucleic Acids Res*. 2003 Feb 15;31(4):e15.
120. Bolstad BM, Irizarry RA, Astrand M, Speed TP. A comparison of normalization methods for high density oligonucleotide array data based on variance and bias. *Bioinformatics*. 2003 Jan 22;19(2):185-93.
121. Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, Speed TP. Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics*. 2003 Apr;4(2):249-64.
122. Du P, Kibbe WA, Lin SM. lumi: a pipeline for processing Illumina microarray. *Bioinformatics*. 2008 Jul 1;24(13):1547-8.
123. Du P, Zhang X, Huang CC, Jafari N, Kibbe WA, Hou L, Lin SM. Comparison of Beta-value and M-value methods for quantifying methylation levels by microarray analysis. *BMC Bioinformatics*. 2010 Nov 30;11:587.
124. Lin SM, Du P, Huber W, Kibbe WA. Model-based variance-stabilizing transformation for Illumina microarray data. *Nucleic Acids Res*. 2008 Feb;36(2):e11.
125. Du P, Kibbe WA, Lin SM. nuld: a universal naming scheme of oligonucleotides for illumina, affymetrix, and other microarrays. *Biol Direct*. 2007 May 31;2:16.
126. <http://www.illumina.com/>. [Online] [Cited: 05 15, 2014.]
127. GK, Smyth. Limma: linear models for microarray data. [book auth.] Carey V, Dudoit S, Irizarry R, Huber W Gentleman R. *Bioinformatics and Computational Biology Solutions using R and Bioconductor*. New York : Springer, 2005.
128. Smyth GK, Speed T. Normalization of cDNA microarray data. *Methods*. 2003 Dec;31(4):265-73.
129. Ritchie ME, Silver J, Oshlack A, Holmes M, Diyagama D, Holloway A, Smyth GK. A comparison of background correction methods for two-colour microarrays. *Bioinformatics*. 2007 Oct 15;23(20):2700-7.
130. Ensembl mouse. [Online] http://www.ensembl.org/Mus_musculus/Info/Index.

131. Smyth GK, Ritchie M, Thorne N, Wettenhal JI, Shi W, Hu Y. Bioconductor - Open source software of bioinformatics. *limma: Linear Models for Microarray and RNA-Seq Data* *User's Guide.* [Online] <http://www.bioconductor.org/packages/release/bioc/vignettes/limma/inst/doc/usersguide.pdf>.
132. Ho Sui SJ, Mortimer JR, Arenillas DJ, Brumm J, Walsh CJ, Kennedy BP, Wasserman WW. oPOSSUM: identification of over-represented transcription factor binding sites in co-expressed genes. *Nucleic Acids Res.* 2005 Jun 2;33(10):3154-64.
133. Kwon AT, Arenillas DJ, Worsley Hunt R, Wasserman WW. oPOSSUM-3: advanced analysis of regulatory motif over-representation across genes or ChIP-Seq datasets. *G3 (Bethesda)*. 2012 Sep;2(9):987-1002.
134. NCBI: GEO. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE50789>.
135. NCBI: GEO. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE40936>.
136. NCBI: GEO. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE39313>.
137. NCBI: GEO. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE36838>.
138. NCBI: GEO. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE26267>.
139. NCBI: GEO. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE2431>.
140. NCBI: GEO. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE1093>.
141. Arrayexpress. [Online] <http://www.ebi.ac.uk/arrayexpress/experiments/E-MEXP-748/>.
142. NCBI: GEO. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE40977>.
143. NCBI: Gene. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE4259>.
144. NCBI: GEO. [Online] <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE38136>.

145. NCBI: GEO. [Online]
<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE23523>.
146. NCBI: GEO. [Online]
<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE11845>.
147. NCBI: GEO. [Online]
<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE37415>.
148. Graf GA, Yu L, Li WP, Gerard R, Tuma PL, Cohen JC, Hobbs HH. ABCG5 and ABCG8 are obligate heterodimers for protein trafficking and biliary cholesterol excretion. *J Biol Chem.* 2003 Nov 28;278(48):48275-82.
149. Nebert DW, Wikvall K, Miller WL. Human cytochromes P450 in health and disease. *Philos Trans R Soc Lond B Biol Sci.* 2013 Jan 6;368(1612):20120431.
150. Cheng J, Edin ML, Hoopes SL, Li H, Bradbury JA, Graves JP, DeGraff LM, Lih FB, Garcia V, Shaik JS, Tomer KB, Flake GP, Falck JR, Lee CR, Poloyac SM, Schwartzman ML, Zeldin DC. Vascular characterization of mice with endothelial expression of cytochrome P450 4F2. *FASEB J.* 2014 Jul;28(7):2915-31.
151. Ghosh HS, McBurney M, Robbins PD. SIRT1 negatively regulates the mammalian target of rapamycin. *PLoS One.* 2010 Feb 15;5(2):e9199.
152. Ogura M, Nakamura Y, Tanaka D, Zhuang X, Fujita Y, Obara A, Hamasaki A, Hosokawa M, Inagaki N. Overexpression of SIRT5 confirms its involvement in deacetylation and activation of carbamoyl phosphate synthetase 1. *Biochem Biophys Res Commun.* 2010 Feb 26;393(1):73-8.
153. Rees DC, Johnson E, Lewinson O. ABC transporters: the power to change. *Nat Rev Mol Cell Biol.* 2009 Mar;10(3):218-27.
154. Frizell E, Liu SL, Abraham A, Ozaki I, Eghbali M, Sage EH, Zern MA. Expression of SPARC in normal and fibrotic livers. *Hepatology.* 1995 Mar;21(3):847-54.
155. Song H, Guan Y, Zhang L, Li K, Dong C. SPARC interacts with AMPK and regulates GLUT4 expression. *Biochem Biophys Res Commun.* 2010 Jun 11;396(4):961-6.
156. Estep PW 3rd, Warner JB, Bulyk ML. Short-term calorie restriction in male mice feminizes gene expression and alters key regulators of conserved aging regulatory pathways. *PLoS One.* 2009;4(4):e5242.
157. Real AM, Hong S, Pissios P. Nicotinamide N-oxidation by CYP2E1 in human liver microsomes. *Drug Metab Dispos.* 2013 Mar;41(3):550-3.

158. Kraus D, Yang Q, Kong D, Banks AS, Zhang L, Rodgers JT, Pirinen E, Pulinkunnil TC, Gong F, Wang YC, Cen Y, Sauve AA, Asara JM, Peroni OD, Monia BP, Bhanot S, Alhonen L, Puigserver P, Kahn BB. Nicotinamide N-methyltransferase knockdown protects against diet-induced obesity. *Nature*. 2014 Apr 10;508(7495):258-62.
159. Hong S, Moreno-Navarrete JM, Wei X, Kikukawa Y, Tzameli I, Prasad D, Lee Y, Asara JM, Fernandez-Real JM, Maratos-Flier E, Pissios P. Nicotinamide N-methyltransferase regulates hepatic nutrient metabolism through Sirt1 protein stabilization. *Nat Med*. 2015 Aug;21(8):887-94.
160. Trammell SA, Brenner C. NNMT: A Bad Actor in Fat Makes Good in Liver. *Cell Metab*. 2015 Aug 4;22(2):200-1.
161. Liu KY, Mistry RJ, Aguirre CA, Fasouli ES, Thomas MG, Klamt F, Ramsden DB, Parsons RB. Nicotinamide N-methyltransferase increases complex I activity in SH-SY5Y cells via sirtuin 3. *Biochem Biophys Res Commun*. 2015 Nov 20;467(3):491-6.
162. Lu X, Kambe F, Cao X, Kozaki Y, Kaji T, Ishii T, Seo H. 3beta-Hydroxysteroid-delta24 reductase is a hydrogen peroxide scavenger, protecting cells from oxidative stress-induced apoptosis. *Endocrinology*. 2008 Jul;149(7):3267-73.
163. Clem BF, Clem AL, Yalcin A, Goswami U, Arumugam S, Telang S, Trent JO, Chesney J. A novel small molecule antagonist of choline kinase- α that simultaneously suppresses MAPK and PI3K/AKT signaling. *Oncogene*. 2011 Jul 28;30(30):3370-80.
164. Yalcin A, Clem B, Makoni S, Clem A, Nelson K, Thornburg J, Siow D, Lane AN, Brock SE, Goswami U, Eaton JW, Telang S, Chesney J. Selective inhibition of choline kinase simultaneously attenuates MAPK and PI3K/AKT signaling. *Oncogene*. 2010 Jan 7;29(1):139-49.
165. Jones JI, Clemons DR. Insulin-like growth factors and their binding proteins: biological actions. *Endocr Rev*. 1995 Feb;16(1):3-34.
166. Goya L, García-Segura LM, Ramos S, Pascual-Leone AM, Argente J, Martín MA, Chowen JA. Interaction between malnutrition and ovarian hormones on the systemic IGF-I axis. *Eur J Endocrinol*. 2002 Sep;147(3):417-24.
167. Musey VC, Goldstein S, Farmer PK, Moore PB, Phillips LS. Differential regulation of IGF-1 and IGF-binding protein-1 by dietary composition in humans. *Am J Med Sci*. 1993 Mar;305(3):131-8.

168. Rasmussen MH, Juul A, Kjems LL, Skakkebaek NE, Hilsted J. Lack of stimulation of 24-hour growth hormone release by hypocaloric diet in obesity. *J Clin Endocrinol Metab.* 1995 Mar;80(3):796-801.
169. Kang HS, Kim MY, Kim SJ, Lee JH, Kim YD, Seo YK, Bae JH, Oh GT, Song DK, Ahn YH, Im SS. Regulation of IGFBP-2 expression during fasting. *Biochem J.* 2015 May 1;467(3):453-60.
170. Lewitt MS, Hall K, Bang P, Brismar K. Altered response of insulin-like growth factor-binding protein 1 to nutritional deprivation in type 2 diabetes mellitus. *Metabolism.* 2005 Mar;54(3):275-80.
171. Gan L, Han Y, Bastianetto S, Dumont Y, Unterman TG, Quirion R. FoxO-dependent and -independent mechanisms mediate SirT1 effects on IGFBP-1 gene expression. *Biochem Biophys Res Commun.* 2005 Dec 2;337(4):1092-6.
172. —. FoxO-dependent and -independent mechanisms mediate SirT1 effects on IGFBP-1 gene expression. *Biochem Biophys Res Commun.* 2005 Dec 2;337(4):1092-6.
173. Domené HM, Hwa V, Argente J, Wit JM, Camacho-Hübner C, Jasper HG, Pozo J, van Duyvenvoorde HA, Yakar S, Fofanova-Gambetti OV, Rosenfeld RG and Group, International ALS Collaborative. Human acid-labile subunit deficiency: clinical, endocrine and metabolic consequences. *Horm Res.* 2009;72(3):129-41.
174. Hu Y, Liu J, Wang J, Liu Q. The controversial links among calorie restriction, SIRT1, and resveratrol. *Free Radic Biol Med.* 2011 Jul 15;51(2):250-6.
175. Pei H, Li L, Fridley BL, Jenkins GD, Kalari KR, Lingle W, Petersen G, Lou Z, Wang L. FKBP51 affects cancer cell response to chemotherapy by negatively regulating Akt. . *Cancer Cell.* . 2009 Sep 8;16(3):259-66.
176. Ersoy BA, Tarun A, D'Aquino K, Hancer NJ, Ukomadu C, White MF, Michel T, Manning BD, Cohen DE. Phosphatidylcholine transfer protein interacts with thioesterase superfamily member 2 to attenuate insulin signaling. . *Sci Signal.* . 2013 Jul 30;6(286):ra64.
177. Liu Z, Zhang HM, Yuan J, Lim T, Sall A, Taylor GA, Yang D. Focal adhesion kinase mediates the interferon-gamma-inducible GTPase-induced phosphatidylinositol 3-kinase/Akt survival pathway and further initiates a positive feedback loop of NF-kappaB activation. *Cell Microbiol.* 2008 Sep;10(9):1787-800.
178. WR., Swindell. Genes and gene expression modules associated with caloric restriction and aging in the laboratory mouse. *BMC Genomics.* 2009 Dec 7;10:585.

179. Fu ZD, Csanaky IL, Klaassen CD. Effects of aging on mRNA profiles for drug-metabolizing enzymes and transporters in livers of male and female mice. *Drug Metab Dispos.* 2012 Jun;40(6):1216-25.
180. Okamoto M, Kita T, Okuda H, Tanaka T, Nakashima T. Effects of aging on acute toxicity of nicotine in rats. *Pharmacol Toxicol.* 1994 Jul;75(1):1-6.
181. Hines RN, Cashman JR, Philpot RM, Williams DE, Ziegler DM. The mammalian flavin-containing monooxygenases: molecular characterization and regulation of expression. *Toxicol Appl Pharmacol.* 1994 Mar;125(1):1-6.
182. Amador-Noguez D, Dean A, Huang W, Setchell K, Moore D, Darlington G. Alterations in xenobiotic metabolism in the long-lived Little mice. *Aging Cell.* 2007 Aug;6(4):453-70.
183. Miao J, Ling AV, Manthena PV, Gearing ME, Graham MJ, Crooke RM, Croce KJ, Esquejo RM, Clish CB and Morbid Obesity Study Group, Vicent D, Biddinger SB. Flavin-containing monooxygenase 3 as a potential player in diabetes-associated atherosclerosis. *Nat Commun.* 2015 Apr 7;6:6498.
184. Zhang YK, Saupe KW, Klaassen CD. Energy restriction does not compensate for the reduced expression of hepatic drug-processing genes in mice with aging. *Drug Metab Dispos.* 2010 Jul;38(7):1122-31.
185. Li L, He D, Wilborn TW, Falany JL, Falany CN. Increased SULT1E1 activity in HepG2 hepatocytes decreases growth hormone stimulation of STAT5b phosphorylation. *Steroids.* 2009 Jan;74(1):20-9.
186. Marcus NY, Blomenkamp K, Ahmad M, Teckman JH. Oxidative stress contributes to liver damage in a murine model of alpha-1-antitrypsin deficiency. *Exp Biol Med (Maywood).* 2012 Oct;237(10):1163-72.
187. Ayyadevara S, Engle MR, Singh SP, Dandapat A, Lichti CF, Benes H, Shmookler Reis RJ, Liebau E, Zimniak P. Lifespan and stress resistance of *Caenorhabditis elegans* are increased by expression of glutathione transferases capable of metabolizing the lipid peroxidation product 4-hydroxynonenal. *Aging Cell.* 2005 Oct;4(5):257-71.
188. Berruyer C, Martin FM, Castellano R, Macone A, Malergue F, Garrido-Urbani S, Millet V, Imbert J, Duprè S, Pitari G, Naquet P, Galland F. Vanin-1/- mice exhibit a glutathione-mediated tissue resistance to oxidative stress. *Mol Cell Biol.* 2004 Aug;24(16):7214-24.

189. Yizhak K, Gabay O, Cohen H, Ruppin E. Model-based identification of drug targets that revert disrupted metabolism and its application to ageing. *Nat Commun.* 2013;4:2632.
190. Chick WS, Ludwig M, Zhao X, Kitzenberg D, Williams K, Johnson TE. Screening for stress-resistance mutations in the mouse. *Front Genet.* 2014 Sep 8;5:310.
191. van Schoorst EM, Keijer J, Pennings JL, Opperhuizen A, van den Brom CE, Kohl T, Franssen-van Hal NL, Hoebee B. Adipose gene expression response of lean and obese mice to short-term dietary restriction. *Obesity (Silver Spring)*. 2006 Jun;14(6):974-9.
192. Miller RA, Chang Y, Galecki AT, Al-Regaiey K, Kopchick JJ, Bartke A. Gene expression patterns in calorically restricted mice: partial overlap with long-lived mutant mice. *Mol Endocrinol.* 2002 Nov;16(11):2657-66.
193. Dhahbi JM, Kim HJ, Mote PL, Beaver RJ, Spindler SR. Temporal linkage between the phenotypic and genomic responses to caloric restriction. *Proc Natl Acad Sci U S A.* 2004 Apr 13;101(15):5524-9.
194. Giller K, Huebbe P, Doering F, Pallauf K, Rimbach G. Major urinary protein 5, a scent communication protein, is regulated by dietary restriction and subsequent re-feeding in mice. *Proc Biol Sci.* 2013 Feb 27;280(1757):20130101.
195. Schloesser A, Campbell G, Glüer CC, Rimbach G, Huebbe P. Restriction on an energy-dense diet improves markers of metabolic health and cellular aging in mice through decreasing hepatic mTOR activity. *Rejuvenation Res.* 2015 Feb;18(1):30-9.
196. Zhou Y, Jiang L, Rui L. Identification of MUP1 as a regulator for glucose and lipid metabolism in mice. *J Biol Chem.* 2009 Apr 24;284(17):11152-9.
197. Chen CC, Lee TY, Kwok CF, Hsu YP, Shih KC, Lin YJ, Ho LT. Major urinary protein 1 interacts with cannabinoid receptor type 1 in fatty acid-induced hepatic insulin resistance in a mouse hepatocyte model. *Biochem Biophys Res Commun.* 2015 May 15;460(4):1063-8.
198. Kuhn NJ, Woodworth-Gutai M, Gross KW, Held WA. Subfamilies of the mouse major urinary protein (MUP) multi-gene family: sequence analysis of cDNA clones and differential regulation in the liver. *Nucleic Acids Res.* 1984 Aug 10;12(15):6073-90.
199. Eggers KM, Kempf T, Wallentin L, Wollert KC, Lind L. Change in growth differentiation factor 15 concentrations over time independently predicts mortality in community-dwelling elderly individuals. *Clin Chem.* 2013 Jul;59(7):1091-8.

200. Yatsuga S, Fujita Y, Ishii A, Fukumoto Y, Arahata H, Kakuma T, Kojima T, Ito M, Tanaka M, Saiki R, Koga Y. Growth differentiation factor 15 as a useful biomarker for mitochondrial disorders. *Ann Neurol.* 2015 Nov;78(5):814-23.
201. Panowski SH, Wolff S, Aguilaniu H, Durieux J, Dillin A. PHA-4/Foxa mediates diet-restriction-induced longevity of *C. elegans*. *Nature*. 2007 May 31;447(7144):550-5.
202. Ma X, Xu L, Gavrilova O, Mueller E. Role of forkhead box protein A3 in age-associated metabolic decline. *Proc Natl Acad Sci U S A.* 2014 Sep 30;111(39):14289-94.
203. Dkhil MA, Al-Quraishy S, Abdel-Baki AA, Ghanjati F, Arauzo-Bravo MJ, Delic D, Wunderlich F. Epigenetic modifications of gene promoter DNA in the liver of adult female mice masculinized by testosterone. *J Steroid Biochem Mol Biol.* 2015 Jan;145:121-30.
204. Ramirez MC, Zubeldía-Brenner L, Wargon V, Ornstein AM, Becu-Villalobos D. Expression and methylation status of female-predominant GH-dependent liver genes are modified by neonatal androgenization in female mice. *Mol Cell Endocrinol.* 2014 Feb 15;382(2):825-34.
205. Zhang Y, Klaassen CD. Hormonal regulation of Cyp4a isoforms in mouse liver and kidney. *Xenobiotica.* 2013 Dec;43(12):1055-63.
206. Wei FZ, Cao Z, Wang X, Wang H, Cai MY, Li T, Hattori N, Wang D, Du Y, Song B, Cao LL, Shen C, Wang L, Wang H, Yang Y, Xie D, Wang F, Ushijima T, Zhao Y, Zhu WG. Epigenetic regulation of autophagy by the methyltransferase EZH2 through an MTOR-dependent pathway. *Autophagy.* 2015 Nov;11(12):2309-22.
207. Lu YF, Wan XL, Xu Y, Liu J. Repeated oral administration of oleanolic acid produces cholestatic liver injury in mice. *Molecules.* 2013 Mar 7;18(3):3060-71.
208. Hu A, Sun M, Yan D, Chen K. Clinical significance of mTOR and eIF4E expression in invasive ductal carcinoma. *Tumori.* 2014 Sep-Oct;100(5):541-6.
209. Wang GX, Zhao XY, Meng ZX, Kern M, Dietrich A, Chen Z, Cozacov Z, Zhou D, Okunade AL, Su X, Li S, Blüher M, Lin JD. The brown fat-enriched secreted factor Nrg4 preserves metabolic homeostasis through attenuation of hepatic lipogenesis. *Nat Med.* 2014 Dec;20(12):1436-43.
210. Zhang Y, Xu D, Huang H, Chen S, Wang L, Zhu L, Jiang X, Ruan X, Luo X, Cao P, Liu W, Pan Y, Wang Z, Chen Y. Regulation of glucose homeostasis and lipid metabolism by PPP1R3G-mediated hepatic glycogenesis. *Mol Endocrinol.* 2014 Jan;28(1):116-26.

211. Luo X, Zhang Y, Ruan X, Jiang X, Zhu L, Wang X, Ding Q, Liu W, Pan Y, Wang Z, Chen Y. Fasting-induced protein phosphatase 1 regulatory subunit contributes to postprandial blood glucose homeostasis via regulation of hepatic glycogenesis. *Diabetes*. 2011 May;60(5):1435-45.
212. Fardini Y, Masson E, Boudah O, Ben Jouira R, Cosson C, Pierre-Eugene C, Kuo MS, Issad T. O-GlcNAcylation of FoxO1 in pancreatic β cells promotes Akt inhibition through an IGFBP1-mediated autocrine mechanism. *FASEB J*. 2014 Feb;28(2):1010-21.
213. Unterman TG, Fareeduddin A, Harris MA, Goswami RG, Porcella A, Costa RH, Lacson RG. Hepatocyte nuclear factor-3 (HNF-3) binds to the insulin response sequence in the IGF binding protein-1 (IGFBP-1) promoter and enhances promoter function. *Biochem Biophys Res Commun*. 1994 Sep 30;203(3):1835-41.
214. O'Brien RM, Noisin EL, Suwanichkul A, Yamasaki T, Lucas PC, Wang JC, Powell DR, Granner DK. Hepatic nuclear factor 3- and hormone-regulated expression of the phosphoenolpyruvate carboxykinase and insulin-like growth factor-binding protein 1 genes. *Mol Cell Biol*. 1995 Mar;15(3):1747-58.
215. Inoki K, Li Y, Zhu T, Wu J, Guan KL. TSC2 is phosphorylated and inhibited by Akt and suppresses mTOR signalling. *Nat Cell Biol*. 2002 Sep;4(9):648-57.
216. Langfelder P, Castellani LW, Zhou Z, Paul E, Davis R, Schadt EE, Lusis AJ, Horvath S, Mehrabian M. A systems genetic analysis of high density lipoprotein metabolism and network preservation across mouse models. *Biochim Biophys Acta*. 2012 Mar;1821(3):435-47.
217. Oh-hashi K, Koga H, Ikeda S, Shimada K, Hirata Y, Kiuchi K. CRELD2 is a novel endoplasmic reticulum stress-inducible gene. *Biochem Biophys Res Commun*. 2009 Sep 25;387(3):504-10.
218. Sheng L, Cho KW, Zhou Y, Shen H, Rui L. Lipocalin 13 protein protects against hepatic steatosis by both inhibiting lipogenesis and stimulating fatty acid β -oxidation. *J Biol Chem*. 2011 Nov 4;286(44):38128-35.
219. Cho KW, Zhou Y, Sheng L, Rui L. Lipocalin-13 regulates glucose metabolism by both insulin-dependent and insulin-independent mechanisms. *Mol Cell Biol*. 2011 Feb;31(3):450-7.
220. WR, Swindell. Comparative analysis of microarray data identifies common responses to caloric restriction among mouse tissues. *Mech Ageing Dev*. 2008 Mar;129(3):138-53.

221. Yang X, Doser TA, Fang CX, Nunn JM, Janardhanan R, Zhu M, Sreejayan N, Quinn MT, Ren J. Metallothionein prolongs survival and antagonizes senescence-associated cardiomyocyte diastolic dysfunction: role of oxidative stress. *FASEB J.* 2006 May;20(7):1024-6.
222. Miles AT, Hawksworth GM, Beattie JH, Rodilla V. Induction, regulation, degradation, and biological significance of mammalian metallothioneins. *Crit Rev Biochem Mol Biol.* 2000;35(1):35-70.
223. Mocchegiani E, Giacconi R, Muti E, Rogo C, Bracci M, Muzzioli M, Cipriano C, Malavolta M. Zinc, immune plasticity, aging, and successful aging: role of metallothionein. *Ann N Y Acad Sci.* 2004 Jun;1019:127-34.
224. Swindell WR, Masternak MM, Bartke A. In vivo analysis of gene expression in long-lived mice lacking the pregnancy-associated plasma protein A (PappA) gene. *Exp Gerontol.* 2010 May;45(5):366-74.
225. Ryu D, Oh KJ, Jo HY, Hedrick S, Kim YN, Hwang YJ, Park TS, Han JS, Choi CS, Montminy M, Koo SH. TORC2 regulates hepatic insulin signaling via a mammalian phosphatidic acid phosphatase, LIPIN1. *Cell Metab.* 2009 Mar;9(3):240-51.
226. Sugden MC, Caton PW, Holness MJ. PPAR control: it's SIRTainly as easy as PGC. *J Endocrinol.* 2010 Feb;204(2):93-104.
227. Finck BN, Gropler MC, Chen Z, Leone TC, Croce MA, Harris TE, Lawrence JC Jr, Kelly DP. Lipin 1 is an inducible amplifier of the hepatic PGC-1alpha/PPARalpha regulatory pathway. *Cell Metab.* 2006 Sep;4(3):199-210.
228. You M, Jogasuria A, Taylor C, Wu J. Sirtuin 1 signaling and alcoholic fatty liver disease. *Hepatobiliary Surg Nutr.* 2015 Apr;4(2):88-100.
229. Bi L, Jiang Z, Zhou J. The role of lipin-1 in the pathogenesis of alcoholic fatty liver. *Alcohol Alcohol.* 2015 Mar;50(2):146-51.
230. Zhang ZF, Fan SH, Zheng YL, Lu J, Wu DM, Shan Q, Hu B. Troxerutin improves hepatic lipid homeostasis by restoring NAD(+) depletion-mediated dysfunction of lipin 1 signaling in high-fat diet-treated mice. *Biochem Pharmacol.* 2014 Sep 1;91(1):74-86.
231. Ahn J, Oh SA, Suh Y, Moeller SJ, Lee K. Porcine G₀/G₁ switch gene 2 (G0S2) expression is regulated during adipogenesis and short-term in-vivo nutritional interventions. *Lipids.* 2013 Mar;48(3):209-18.
232. Nielsen TS, Vendelbo MH, Jessen N, Pedersen SB, Jørgensen JO, Lund S, Møller N. Fasting, but not exercise, increases adipose triglyceride lipase (ATGL)

- protein and reduces G(0)/G(1) switch gene 2 (G0S2) protein and mRNA content in human adipose tissue. *J Clin Endocrinol Metab.* 2011 Aug;96(8):E1293-7.
233. Zhang W, Bu SY, Mashek MT, O-Sullivan I, Sibai Z, Khan SA, Ilkayeva O, Newgard CB, Mashek DG, Unterman TG. Integrated Regulation of Hepatic Lipid and Glucose Metabolism by Adipose Triacylglycerol Lipase and FoxO Proteins. *Cell Rep.* 2016 Mar 29. pii: S2211-1247(16)30262-5.
234. GenAge: The Aging Gene Database. [Online] [Cited: 01 23, 2015.] <http://genomics.senescence.info/genes/search.php?search.php?show=1&sort=1&page=1&organism=Mus%20musculus>.
235. GenAge: The Aging Gene Database. [Online] [Cited: january 23, 2015.] <http://genomics.senescence.info/genes/stats.php>.
236. JA, Baur. Resveratrol, sirtuins, and the promise of a DR mimetic. *Mech Ageing Dev.* 2010 Apr;131(4):261-9.
237. Ma L, Zhao Z, Wang R, Zhang X, Zhang J, Dong W, Xu B, Zhang J. Caloric restriction can improve learning ability in C57/BL mice via regulation of the insulin-PI3K/Akt signaling pathway. *Neurol Sci.* 2014 Sep;35(9):1381-6.
238. Gesing A, Al-Regaiey KA, Bartke A, Masternak MM. Growth hormone abolishes beneficial effects of calorie restriction in long-lived Ames dwarf mice. *Exp Gerontol.* 2014 Oct;58:219-29.
239. Wang CC, Adochio RL, Leitner JW, Abeyta IM, Draznin B, Cornier MA. Acute effects of different diet compositions on skeletal muscle insulin signalling in obese individuals during caloric restriction. *Metabolism.* 2013 Apr;62(4):595-603.
240. Wang P, Zhang RY, Song J, Guan YF, Xu TY, Du H, Viollet B, Miao CY. Loss of AMP-activated protein kinase- α 2 impairs the insulin-sensitizing effect of calorie restriction in skeletal muscle. *Diabetes.* 2012 May;61(5):1051-61.
241. Guo XD, Zhang DY, Gao XJ, Parry J, Liu K, Liu BL, Wang M. Quercetin and quercetin-3-O-glucuronide are equally effective in ameliorating endothelial insulin resistance through inhibition of reactive oxygen species-associated inflammation. *Mol Nutr Food Res.* 2013 Jun;57(6):1037-45.
242. Wang S, DeGroff VL, Clinton SK. Tomato and soy polyphenols reduce insulin-like growth factor-I-stimulated rat prostate cancer cell proliferation and apoptotic resistance in vitro via inhibition of intracellular signaling pathways involving tyrosine kinase. *J Nutr.* 2003 Jul;133(7):2367-76.

243. Phan TT, See P, Tran E, Nguyen TT, Chan SY, Lee ST, Huynh H. Suppression of insulin-like growth factor signalling pathway and collagen expression in keloid-derived fibroblasts by quercetin: its therapeutic potential use in the treatment and/or prevention of keloids. *Br J Dermatol.* 2003 Mar;148(3):544-52.
244. Kim DH, Kim JY, Yu BP, Chung HY. The activation of NF-kappaB through Akt-induced FOXO1 phosphorylation during aging and its modulation by calorie restriction. *Biogerontology.* 2008 Feb;9(1):33-47.
245. Chiş IC, Mureşan A, Oros A, Nagy AL, Clichici S. Protective effects of Quercetin and chronic moderate exercise (training) against oxidative stress in the liver tissue of streptozotocin-induced diabetic rats. *Acta Physiol Hung.* 2016 Mar;103(1):49-64.
246. Zou W, Liu W, Yang B, Wu L, Yang J, Zou T, Liu F, Xia L, Zhang D. Quercetin protects against perfluorooctanoic acid-induced liver injury by attenuating oxidative stress and inflammatory response in mice. *Int Immunopharmacol.* 2015 Sep;28(1):129-35.
247. Li JM, Wang W, Fan CY, Wang MX, Zhang X, Hu QH, Kong LD. Quercetin Preserves β -Cell Mass and Function in Fructose-Induced Hyperinsulinemia through Modulating Pancreatic Akt/FoxO1 Activation. *Evid Based Complement Alternat Med.* 2013;2013:303902.
248. Huang CY, Chan CY, Chou IT, Lien CH, Hung HC, Lee MF. Quercetin induces growth arrest through activation of FOXO1 transcription factor in EGFR-overexpressing oral cancer cells. *J Nutr Biochem.* 2013 Sep;24(9):1596-603.
249. A, Bruning. Inhibition of mTOR signaling by quercetin in cancer treatment and prevention. *Anticancer Agents Med Chem.* 2013 Sep;13(7):1025-31.
250. Coschigano KT, Holland AN, Riders ME, List EO, Flyvbjerg A, Kopchick JJ. Deletion, but not antagonism, of the mouse growth hormone receptor results in severely decreased body weights, insulin, and insulin-like growth factor I levels and increased life span. *Endocrinology.* Sep;144(9):3799-810, 2003.
251. Belinha I, Amorim MA, Rodrigues P, de Freitas V, Moradas-Ferreira P, Mateus N, Costa V. Quercetin increases oxidative stress resistance and longevity in *Saccharomyces cerevisiae*. *J Agric Food Chem.* 2007 Mar 21;55(6):2446-51.
252. Saul N, Pietsch K, Menzel R, Steinberg CE. Quercetin-mediated longevity in *Caenorhabditis elegans*: is DAF-16 involved? *Mech Ageing Dev.* 2008 Oct;129(10):611-3.

253. Xiang L, Sun K, Lu J, Weng Y, Taoka A, Sakagami Y, Qi J. Anti-aging effects of phloridzin, an apple polyphenol, on yeast via the SOD and Sir2 genes. *Biosci Biotechnol Biochem*. 2011;75(5):854-8.
254. Lee EB, Ahn D, Kim BJ, Lee SY, Seo HW, Cha YS, Jeon H, Eun JS, Cha DS, Kim DK. Genistein from Vigna angularis Extends Lifespan in *Caenorhabditis elegans*. *Biomol Ther (Seoul)*. 2015 Jan;23(1):77-83.
255. Altun D, Uysal H, Aşkın H, Ayar A. Determination of the effects of genistein on the longevity of *Drosophila melanogaster* meigen (Diptera; Drosophilidae). *Bull Environ Contam Toxicol*. 2011 Jan;86(1):120-3.
256. Vina J, Gambini J, Lopez-Grueso R, Abdelaziz KM, Jove M, Borras C. Females live longer than males: role of oxidative stress. *Curr Pharm Des*. 2011 Dec 1;17(36):3959-65.
257. Borrás C, Gambini J, Gómez-Cabrera MC, Sastre J, Pallardó FV, Mann GE, Viña J. Genistein, a soy isoflavone, up-regulates expression of antioxidant genes: involvement of estrogen receptors, ERK1/2, and NFκB. *FASEB J*. 2006 Oct;20(12):2136-8.
258. Edgel KA, McMillen TS, Wei H, Pamir N, Houston BA, Caldwell MT, Mai PO, Oram JF, Tang C, Leboeuf RC. Obesity and weight loss result in increased adipose tissue ABCG1 expression in db/db mice. *Biochim Biophys Acta*. 2012 Mar;1821(3):425-34.
259. Sabeva NS, Rouse EJ, Graf GA. Defects in the leptin axis reduce abundance of the ABCG5-ABCG8 sterol transporter in liver. *J Biol Chem*. 2007 Aug 3;282(31):22397-405.
260. Kulkarni SR, Xu J, Donepudi AC, Wei W, Slitt AL. Effect of caloric restriction and AMPK activation on hepatic nuclear receptor, biotransformation enzyme, and transporter expression in lean and obese mice. *Pharm Res*. 2013 Sep;30(9):2232-47.
261. Ferland G, Tuchweber B, Perea A, Yousef IM. Effect of aging and dietary restriction on bile acid metabolism in rats. *Lipids*. 1989 Oct;24(10):842-8.
262. Fu ZD, Klaassen CD. Increased bile acids in enterohepatic circulation by short-term calorie restriction in male mice. *Toxicol Appl Pharmacol*. 2013 Dec 15;273(3):680-90.
263. Thomàs-Moyà E, Nadal-Casellas A, Gianotti M, Lladó I, Proenza AM. Time-dependent modulation of rat serum paraoxonase 1 activity by fasting. *Pflugers Arch*. 2007 Mar;453(6):831-7.

264. Martini C, Pallottini V, De Marinis E, Marino M, Cavallini G, Donati A, Straniero S, Trentalance A. Omega-3 as well as caloric restriction prevent the age-related modifications of cholesterol metabolism. *Mech Ageing Dev.* 2008 Dec;129(12):722-7.
265. Cangemi R, Friedmann AJ, Holloszy JO, Fontana L. Long-term effects of calorie restriction on serum sex-hormone concentrations in men. *Aging Cell.* 2010 Apr;9(2):236-42.
266. Troyer DA, Venkatraman JT, Fernandes G. Effects of calorie restriction and ω-3 dietary fat on aging in short-and long-lived rodents. *Age (Omaha).* 1998 Oct;21(4):175-82.
267. Echchgadda I, Song CS, Oh TS, Cho SH, Rivera OJ, Chatterjee B. Gene regulation for the senescence marker protein DHEA-sulfotransferase by the xenobiotic-activated nuclear pregnane X receptor (PXR). *Mech Ageing Dev.* 2004 Oct-Nov;125(10-11):733-45.
268. Corton JC, Apte U, Anderson SP, Limaye P, Yoon L, Latendresse J, Dunn C, Everitt JI, Voss KA, Swanson C, Kimbrough C, Wong JS, Gill SS, Chandraratna RA, Kwak MK, Kensler TW, Stulnig TM, Steffensen KR, Gustafsson JA, Mehendale HM. Mimetics of caloric restriction include agonists of lipid-activated nuclear receptors. *J Biol Chem.* 2004 Oct 29;279(44):46204-12.
269. Rocha JS, Bonkowski MS, Masternak MM, França LR, Bartke A. Effects of adult onset mild caloric restriction on weight of reproductive organs, plasma parameters and gene expression in male mice. *Anim Reprod.* 2012 Jan 1;9(1):40-51.
270. Zhu M, Miura J, Lu LX, Bernier M, DeCabo R, Lane MA, Roth GS, Ingram DK. Circulating adiponectin levels increase in rats on caloric restriction: the potential for insulin sensitization. *Exp Gerontol.* 2004 Jul;39(7):1049-59.
271. Khraiwesh H, López-Domínguez JA, Fernández del Río L, Gutierrez-Casado E, López-Lluch G, Navas P, de Cabo R, Ramsey JJ, Burón MI, Villalba JM, González-Reyes JA. Mitochondrial ultrastructure and markers of dynamics in hepatocytes from aged, calorie restricted mice fed with different dietary fats. *Exp Gerontol.* 2014 Aug;56:77-88.
272. HS, Ranhotra. Long-term caloric restriction up-regulates PPAR gamma co-activator 1 alpha (PGC-1alpha) expression in mice. *Indian J Biochem Biophys.* 2010 Oct;47(5):272-7.
273. SR, Burzynski. Aging: gene silencing or gene activation? *Med Hypotheses.* 2005;64(1):201-8.

274. Giller K, Huebbe P, Hennig S, Dose J, Pallauf K, Doering F, Rimbach G. Beneficial effects of a 6-month dietary restriction are time-dependently abolished within 2 weeks or 6 months of refeeding-genome-wide transcriptome analysis in mouse liver. *Free Radic Biol Med.* 2013 Aug;61:170-8.
275. Leakey JA, Cunny HC, Bazare J Jr, Webb PJ, Lipscomb JC, Slikker W Jr, Feuers RJ, Duffy PH, Hart RW. Effects of aging and caloric restriction on hepatic drug metabolizing enzymes in the Fischer 344 rat. II: Effects on conjugating enzymes. *Mech Ageing Dev.* 1989 May;48(2):157-66.
276. Wen H, Yang HJ, An YJ, Kim JM, Lee DH, Jin X, Park SW, Min KJ, Park S. Enhanced phase II detoxification contributes to beneficial effects of dietary restriction as revealed by multi-platform metabolomics studies. *Mol Cell Proteomics.* 2013 Mar;12(3):575-86.
277. Froy O, Chapnik N, Miskin R. Long-lived alphaMUPA transgenic mice exhibit pronounced circadian rhythms. *Am J Physiol Endocrinol Metab.* 2006 Nov;291(5):E1017-24.
278. Resuehr D, Olcese J. Caloric restriction and melatonin substitution: effects on murine circadian parameters. *Brain Res.* 2005 Jun 28;1048(1-2):146-52.
279. Bonaconsa M, Malpeli G, Montaruli A, Carandente F, Grassi-Zucconi G, Bentivoglio M. Differential modulation of clock gene expression in the suprachiasmatic nucleus, liver and heart of aged mice. *Exp Gerontol.* 2014 Jul;55:70-9.
280. Yamaza H, Komatsu T, Wakita S, Kijogi C, Park S, Hayashi H, Chiba T, Mori R, Furuyama T, Mori N, Shimokawa I. FoxO1 is involved in the antineoplastic effect of calorie restriction. *Aging Cell.* 2010 Jun;9(3):372-82.
281. Selman C, Kendaiah S, Gredilla R, Leeuwenburgh C. Increased hepatic apoptosis during short-term caloric restriction is not associated with an enhancement in caspase levels. *Exp Gerontol.* 2003 Aug;38(8):897-903.
282. García-Matas S, Paul RK, Molina-Martínez P, Palacios H, Gutierrez VM, Corpas R, Pallas M, Cristòfol R, de Cabo R, Sanfeliu C. In vitro caloric restriction induces protective genes and functional rejuvenation in senescent SAMP8 astrocytes. *Aging Cell.* 2015 Feb 25.
283. Donati A, Cavallini G, Bergamini E. Methods for inducing and monitoring liver autophagy relative to aging and antiaging caloric restriction in rats. *Methods Enzymol.* 2009;452:441-55.

284. Donati A, Recchia G, Cavallini G, Bergamini E. Effect of aging and anti-aging caloric restriction on the endocrine regulation of rat liver autophagy. *J Gerontol A Biol Sci Med Sci*. 2008 Jun;63(6):550-5.
285. Mariño G, Pietrocola F, Madeo F, Kroemer G. Caloric restriction mimetics: natural/physiological pharmacological autophagy inducers. *Autophagy*. 2014 Nov 2;10(11):1879-82.
286. Richard VR, Leonov A, Beach A, Burstein MT, Koupaki O, Gomez-Perez A, Levy S, Pluska L, Mattie S, Rafesh R, Iouk T, Sheibani S, Greenwood M, Vali H, Titorenko VI. Macromitophagy is a longevity assurance process that in chronologically aging yeast limited in calorie supply sustains functional mitochondria and maintains cellular lipid homeostasis. *Aging (Albany NY)*. 2013 Apr;5(4):234-69.
287. Ding F, Gil MP, Franklin M, Ferreira J, Tatar M, Helfand SL, Neretti N. Transcriptional response to dietary restriction in *Drosophila melanogaster*. *J Insect Physiol*. 2014 Oct;69:101-6.
288. Yu Z, Wang R, Fok WC, Coles A, Salmon AB, Pérez VI. Rapamycin and Dietary Restriction Induce Metabolically Distinctive Changes in Mouse Liver. *J Gerontol A Biol Sci Med Sci*. 2014 Apr 22. [Epub ahead of print].
289. Kuhla A, Hahn S, Butschkau A, Lange S, Wree A, Vollmar B. Lifelong caloric restriction reprograms hepatic fat metabolism in mice. *J Gerontol A Biol Sci Med Sci*. 2014 Aug;69(8):915-22.
290. Okita N, Hayashida Y, Kojima Y, Fukushima M, Yuguchi K, Mikami K, Yamauchi A, Watanabe K, Noguchi M, Nakamura M, Toda T, Higami Y. Differential responses of white adipose tissue and brown adipose tissue to caloric restriction in rats. *Mech Ageing Dev*. 2012 May;133(5):255-66.
291. Davis LJ, Biagi PL, Spanò C, Walford R, Licastro F. Dietary restriction maintains linoleic and arachidonic acid levels in murine liver during aging. *Aging (Milano)*. 1991 Dec;3(4):405-7.
292. Davis LJ, Tadolini B, Biagi PL, Walford R, Licastro F. Effect of age and extent of dietary restriction on hepatic microsomal lipid peroxidation potential in mice. *Mech Ageing Dev*. 1993 Dec 15;72(2):155-63.
293. Jové M, Naudí A, Ramírez-Núñez O, Portero-Otín M, Selman C, Withers DJ, Pamplona R. Caloric restriction reveals a metabolomic and lipidomic signature in liver of male mice. *Aging Cell*. 2014 Oct;13(5):828-37.

294. Hagopian K, Ramsey JJ, Weindruch R. Caloric restriction increases gluconeogenic and transaminase enzyme activities in mouse liver. *Exp Gerontol.* 2003 Mar;38(3):267-78.
295. Dhahbi JM, Mote PL, Wingo J, Tillman JB, Walford RL, Spindler SR. Calories and aging alter gene expression for gluconeogenic, glycolytic, and nitrogen-metabolizing enzymes. *Am J Physiol.* 1999 Aug;277(2 Pt 1):E352-60.
296. Hagopian K, Soo Hoo R, López-Domínguez JA, Ramsey JJ. Calorie restriction influences key metabolic enzyme activities and markers of oxidative damage in distinct mouse liver mitochondrial sub-populations. *Life Sci.* 2013 Dec 5;93(24):941-8.
297. Hagopian K, Ramsey JJ, Weindruch R. Krebs cycle enzymes from livers of old mice are differentially regulated by caloric restriction. *Exp Gerontol.* 2004 Aug;39(8):1145-54.
298. Whitaker R, Gil MP, Ding F, Tatar M, Helfand SL, Neretti N. Dietary switch reveals fast coordinated gene expression changes in *Drosophila melanogaster*. *Aging (Albany NY)*. 2014 May;6(5):355-68.
299. Rezzi S, Martin FP, Shanmuganayagam D, Colman RJ, Nicholson JK, Weindruch R. Metabolic shifts due to long-term caloric restriction revealed in nonhuman primates. *Exp Gerontol.* 2009 May;44(5):356-62.
300. Lee CK, Klopp RG, Weindruch R, Prolla TA. Gene expression profile of aging and its retardation by caloric restriction. *Science.* 1999 Aug 27;285(5432):1390-3.
301. Lee CK, Weindruch R, Prolla TA. Gene-expression profile of the ageing brain in mice. *Nat Genet.* 2000 Jul;25(3):294-7.
302. Xu C, Cai Y, Fan P, Bai B, Chen J, Deng HB, Che CM, Xu A, Vanhoutte PM, Wang Y. Calorie restriction prevents metabolic ageing caused by abnormal SIRT1 function in adipose tissues. *Diabetes.* 2014 Dec 4. pii: DB_141180.
303. Y, Wang. Molecular Links between Caloric Restriction and Sir2/SIRT1 Activation. *Diabetes Metab J.* 2014 Oct;38(5):321-9.
304. Xu C, Bai B, Fan P, Cai Y, Huang B, Law IK, Liu L, Xu A, Tung C, Li X, Siu FM, Che CM, Vanhoutte PM, Wang Y. Selective overexpression of human SIRT1 in adipose tissue enhances energy homeostasis and prevents the deterioration of insulin sensitivity with ageing in mice. *Am J Transl Res.* 2013 May 24;5(4):412-26.
305. Rahman M, Halade GV, Bhattacharya A, Fernandes G. The fat-1 transgene in mice increases antioxidant potential, reduces pro-inflammatory cytokine levels, and

- enhances PPAR-gamma and SIRT-1 expression on a calorie restricted diet. *Oxid Med Cell Longev.* 2009 Nov-Dec;2(5):307-16.
306. Libert S, Chao Y, Zwiener J, Pletcher SD. Realized immune response is enhanced in long-lived puc and chico mutants but is unaffected by dietary restriction. *Mol Immunol.* 2008 Feb;45(3):810-7.
307. Tung BT, Rodríguez-Bies E, Talero E, Gamero-Estévez E, Motilva V, Navas P, López-Lluch G. Anti-inflammatory effect of resveratrol in old mice liver. *Exp Gerontol.* 2015 Apr;64:1-7.
308. Christadoss P, Talal N, Lindstrom J, Fernandes G. Suppression of cellular and humoral immunity to T-dependent antigens by calorie restriction. *Cell Immunol.* 1984 Oct 1;88(1):1-8.
309. Fu C, Hickey M, Morrison M, McCarter R, Han ES. Tissue specific and non-specific changes in gene expression by aging and by early stage CR. *Mech Ageing Dev.* 2006 Dec;127(12):905-16.
310. Effros RB, Walford RL, Weindruch R, Mitcheltree C. Influences of dietary restriction on immunity to influenza in aged mice. *J Gerontol.* 1991 Jul;46(4):B142-7.
311. Kitada M, Kume S, Takeda-Watanabe A, Tsuda S, Kanasaki K, Koya D. Calorie restriction in overweight males ameliorates obesity-related metabolic alterations and cellular adaptations through anti-aging effects, possibly including AMPK and SIRT1 activation. *Biochim Biophys Acta.* 2013 Oct;1830(10):4820-7.
312. Paulson QX, Hong J, Holcomb VB, Nunez NP. Effects of body weight and alcohol consumption on insulin sensitivity. *Nutr J.* 2010 Mar 22;9:14.
313. Moore T, Beltran L, Carbajal S, Hursting SD, DiGiovanni J. Energy balance modulates mouse skin tumor promotion through altered IGF-1R and EGFR crosstalk. *Cancer Prev Res (Phila).* 2012 Oct;5(10):1236-46.
314. Moore T, Checkley LA, DiGiovanni J. Dietary energy balance modulation of epithelial carcinogenesis: a role for IGF-1 receptor signaling and crosstalk. *Ann N Y Acad Sci.* 2011 Jul;1229:7-17.
315. Dahabi JM, Cao SX, Tillman JB, Mote PL, Madore M, Walford RL, Spindler SR. Chaperone-mediated regulation of hepatic protein secretion by caloric restriction. *Biochem Biophys Res Commun.* 2001 Jun 8;284(2):335-9.
316. Richards SE, Wang Y, Claus SP, Lawler D, Kochhar S, Holmes E, Nicholson JK. Metabolic phenotype modulation by caloric restriction in a lifelong dog study. *J Proteome Res.* 2013 Jul 5;12(7):3117-27.

317. SR, Spindler. Calorie restriction enhances the expression of key metabolic enzymes associated with protein renewal during aging. *Ann N Y Acad Sci.* 2001 Apr;928:296-304.
318. Scrofano MM, Shang F, Nowell TR Jr, Gong X, Smith DE, Kelliher M, Dunning J, Mura CV, Taylor A. Aging, calorie restriction and ubiquitin-dependent proteolysis in the livers of Emory mice. *Mech Ageing Dev.* 1998 Apr 1;101(3):277-96.
319. Riis B, Rattan SI, Palmquist K, Nilsson A, Nygård O, Clark BF. Elongation factor 2-specific calcium and calmodulin dependent protein kinase III activity in rat livers varies with age and calorie restriction. *Biochem Biophys Res Commun.* 1993 May 14;192(3):1210-6.
320. Yang F, He Y, Liu HX, Tsuei J, Jiang X, Yang L, Wang ZT, Wan YJ. All-trans retinoic acid regulates hepatic bile acid homeostasis. *Biochem Pharmacol.* 2014 Oct 15;91(4):483-9.
321. Pae M, Meydani SN, Wu D. The role of nutrition in enhancing immunity in aging. *Aging Dis.* 2012 Feb;3(1):91-129.
322. Finamore A, Devirgiliis C, Panno D, D'Aquino M, Polito A, Venneria E, Raguzzini A, Coudray C, Mengheri E. Immune response in relation to zinc status, sex and antioxidant defence in Italian elderly population: the ZENITH study. *Eur J Clin Nutr.* 2005 Nov;59 Suppl 2:S68-72.
323. Fraker PJ, Lill-Elghanian DA. The many roles of apoptosis in immunity as modified by aging and nutritional status. *J Nutr Health Aging.* 2004;8(1):56-63.
324. Simon SF, Taylor CG. Dietary zinc supplementation attenuates hyperglycemia in db/db mice. *Exp Biol Med (Maywood).* 2001 Jan;226(1):43-51.
325. Trapp J, Jung M. The role of NAD⁺ dependent histone deacetylases (sirtuins) in ageing. *Curr Drug Targets.* 2006 Nov;7(11):1553-60.
326. Cabreiro F, Perichon M, Jatje J, Malavolta M, Moccagiani E, Friguet B, Petropoulos I. Zinc supplementation in the elderly subjects: effect on oxidized protein degradation and repair systems in peripheral blood lymphocytes. *Exp Gerontol.* 2008 May;43(5):483-7.

Appendix

A: Number of regulated genes by the different interventions dependent on the used test

	upregulation				downregulation			
	in at least one exper.	binom. test	chi squared test	3-fold	in at least one exper.	binom. test	chi squared test	3-fold
CR								
ttest	15185	403	26	347	6649	508	54	460
ttestfc	34	3	0	3	62	1	0	1
modttest	16712	690	79	571	7128	742	81	577
modt-testfc	40	8	1	7	90	2	0	2
Wilcoxon	15017	428	49	331	6278	423	41	369
wilcox-onfc	30	3	0	3	58	0	0	0
R								
ttest	267	53	0	53	478	48	0	48
ttestfc	39	10	0	10	3	2	0	2
modttest	1415	561	0	553	1808	531	0	530
modt-testfc	150	34	0	34	12	7	0	7
Wilcoxon	0	0	0	0	0	0	0	0
wilcox-onfc	0	0	0	0	0	0	0	0
Q								
ttest	0	0	0	0	0	0	0	0
ttestfc	0	0	0	0	0	0	0	0
modttest	10	0	0	0	15	0	0	0
modt-testfc	0	0	0	0	0	0	0	0
Wilcoxon	0	0	0	0	0	0	0	0
wilcox-onfc	0	0	0	0	0	0	0	0
P								
ttest	1977	277	0	277	2050	417	0	416
ttestfc	0	0	0	0	0	0	0	0
modttest	4383	902	0	901	3890	952	0	951
modt-testfc	1	0	0	0	0	0	0	0
Wilcoxon	0	0	0	0	0	0	0	0
wilcox-onfc	0	0	0	0	0	0	0	0
Gi								
ttest	0	0	0	0	0	0	0	0
ttestfc	0	0	0	0	0	0	0	0
modttest	1879	503	0	498	1634	452	0	440
modt-testfc	19	3	0	3	15	11	0	11
Wilcoxon	0	0	0	0	0	0	0	0
wilcox-onfc	0	0	0	0	0	0	0	0
Qi								
ttest	0	0	0	0	0	0	0	0
ttestfc	0	0	0	0	0	0	0	0
modttest	4866	2504	0	248	4688	2460	0	245
modt-testfc	37	4	0	4	19	13	0	13
Wilcoxon	0	0	0	0	0	0	0	0
wilcox-onfc	0	0	0	0	0	0	0	0

B: Overlapping genes between C and the plant bioactives identified by the different analysis methods

	upregulation	downregulation
CR and resveratrol		
t-test	Gpx7, Rras	
moderated t-test	1810046K07Rik, 2610002J02Rik, Acpp, Aldoc, Cyp17a1, Cyp51, Dclk3, Fam167b, Fam3c, Flot1, Fmo4, Got1, Gpx7, H2-DMA, Hcls1, Ifi27l2b, Il18, Il1b, Lacc1, Mmd2, Mvk, Mycl, Ngef, Nipal1, Nrg4, Nsdhl, Orm1, Per1, Pla1a, Slc38a3, Tmem50a, Tox, Usp2	Acbd4, Ankrd46, Clock, Clpx, Cyp2d10, Fam98a, Gpd2, Hdhd3, Hspa5, Hsph1, Sec14l2, Stip1
rank product method	0610007P14Rik, 1110034G24Rik, 2610002J02Rik, 4933422H20Rik, 8430419L09Rik, Abcc4, Abcg5, Abcg8, Acot3, Acox1, Acpp, Acss2, Acvr2b, Adprm, Adrb2, Aim1, Akr1b3, Akr1b7, Alas1, Alb, Aldh1b1, Aldoc, Angptl4, Anxa5, Apbb3, Aqp11, Arhgap18, Arhgap21, Arrdc4, As3mt, Asah1, Atp10d, Atp1b1, Avpi1, Baiap2l1, BC089597, Blhhe40, Casp3, Cbx7, Ccrn4l, Cd55, Cd63, Cd9, Cdkn2c, Cebpb, Cela2a, Chac1, Chp1, Chrna4, Cib3, Cirbp, Cish, Cnn3, Col4a2, Commd10, Coq10b, Cpsf4, Crcp, Crip1, Cry2, Crym, Ctgf, Cux2, Cyp17a1, Cyp2b13, Cyp2b9, Cyp2g1, Cyp4a10, Cyp4a14, Cyp51, Dbp, Dclk3, Dhcr7, Egr1, Eif4ebp3, Emp2, Enpp1, Ephb4, Eppk1, Etnk2, Fabp5, Fam134b, Fam167b, Fam180a, Fam76b, Fbxo21, Fbxo31, Fdps, Fkbp5, Flcn, Flot1, Fmo3, Fmo4, Gal3st1, Gfra1, Gja4, Glrx, Gnat1, Gne, Gng10, Got1, Gpcpd1, Gpihbp1, Gpx6, Gpx7, Grn, Gstm5, Gstm7, Gsto1, Gstt2, Gtdc1, Gusb, Gycp, H2-K1, Hjurp, Hmgcr, Hmgcs1, Hmox1, Hnrnpa1, Hook2, Id1, Idh1, Idh3g, Idi1, Ier3, Ifi27l2b, Il15, Il15ra, Il17rb, Il18, Il6ra, Izumo4, Kctd12, Klf1, Lacc1, Ldb1, Lgals4, Lpin2, Lss, Lyve1, Manea, Map3k6, Mapre2, Mast3, Med30, Mertk, Mfsd2a, Mknk2, Mmadhc, Mt1, Mthfd1l, Mvk, Myadm, Mycl, N4bp2l1, Naa38, Nadsyn1, Ngef, Nhp2l1, Nipal1, Nr1c5, Nme4, Nmrk1, Nnmt, Npc2, Nrg4, Nsdhl, Nudt2, Orm1, Orm2, Pafah1b3, Pank3, Papss2, Pck1, Pcolce, Pcsk4, Pcsk9, Pdk1, Pdzk1p1, Per1, Per2, Per3, Pim3, Pitpnc1, Pla1a, Pmvk, Pnpla2, Pold4, Por, Ppargc1b, Ppl, Ppp3cc, Pptc7, Prdx2, Prdx3, Prg4, Pros1, Prss8, Psmb10, PstPIP2, Rapgef5, Rbp1, Rdh11, Rdh9, Rgs16, Ripk4, Rnase4, Rnaseh2b, Rnf145, Rpa3, Rpp38, Rras, S100a6, Sbk1, Sc4mol, Sds, Selenbp1, Setd4, Sgk1, Sh3d21, Slc15a4, Slc19a1, Slc20a1, Slc22a26, Slc38a3, Slc3a1, Slc46a3, Slc5a6, Slc7a2, Slc8b1, Slco1a4, Slpi, Smoc2, Snai2, Soc52, Sorbs3, Spata13, Spata2l, Spg21, Sqle, St3gal5, Stap2, Stard8, Sult1d1, Sun2, Tapbp, Tat, Tef, Timm8a1, Timp3, Tk1, Tm4sf4, Tmc7, Tmem41a, Tmem9,	4931406C07Rik, Alas2, Alg12, Alg2, Atf5, Bbx, Bhmt, Clock, Creld2, Ddc, Ddx6, Dnase2a, Fgf21, G0s2, Gmppb, Gpd1, Gphn, Hsd3b2, Hspa4, Hspa5, Lipa, Lonp2, Lrrc59, March8, Mlec, Nat8, Nsmf, Pdilt, Pds5a, Pknox1, Ppp1r12a, S100a10, Sdr9c7, Sox6os, Srprb, Tex2, Tmcc1, Tmed9, Tsc22d1, Tubb4b, Wfdc2

	Tmem97, Tox, Trib3, Tubb5, Uaca, Ugp2, Ugt1a2, Ulk1, Unc5b, Usp2, Vcam1, Vim, Wsb1, Zbtb48, Zfp275, Zfp36, Zfp933, Zkscan17	Stat2, Stip1, Stk16, Stt3b, Sucnr1, Swsap1, Syvn1, Tars, Tbl1x, Tesk1, Tex2, Tjp3, Tlr5, Tmcc1, Tmed9, Tmie, Tnnt1, Tnnt3, Tpgs1, Trappc13, Trim39, Trmt10a, Trp53inp2, Tsacc, Tsc22d1, Tspan33, Tssc4, Tstd2, Ttc7b, Tuba1c, Tubb4b, Tysnd1, Ugt2b1, Ugt2b34, Ugt2b35, Uso1, Vcp, Vps37b, Wdtc1, Wfdc2, Wrap73, Xylb, Zbtb20, Zfand2a, Zfp322a, Zfp361, Zfp809, Zswim1
CR and phloridzin		
t-test	Alb, Chrna4, Nipal1, Tbc1d30, Tox	1415788_at, 1415789_a_at, 1417907_at, 1418520_at, 1423309_at, 1429272_a_at, 1450007_at, Acad11, Acp2, Aldh3a2, Amd1, Apol7a, Arf2, Ces1e, Chrna2, Decr1, Eaf1, Eif2b3, Entpd5, Fabp2, Gdf15, Gid8, Glo1, Hsd3b5, Ldha, Sdr42e1, Tmem33, Vcp
moderated t-test	1415895_at, 1415896_x_at, 1440212_at, 1441539_at, 1444973_at, 1449787_at, BC089597, Chrna4, Ggt6, Igsf8, Lrrc24, Nipal1, Slpi, Tbc1d30, Tox	1418520_at, 1420867_at, 1420868_s_at, 1423263_at, 1423308_at, 1423309_at, 1424185_a_at, 32842_s_at, 1439463_x_at, 1440325_at, 1443505_at, 1810058I24Rik, 4833417J20Rik, 4930506M07Rik, 4931406C07Rik, Abcb4, Acad11, Akr1c20, Alad, Aldh1a1, Aldh3a2, Amd1, Apol7a, Aptx, Aqp4, Arf2, Arpp19, Bcap29, Ccdc162, Cd1d1, Ces1e, Chrna2, Chuk, Cpsf2, Csad, Decr1, Dhx29, Dleu2, Dpys, Dsp, Eif2b3, Fabp2, Gdf15, Hsd3b2, Hsdl2, Ikbkg, Inhbc, Mal2, Mki67ip, Nadk2, Ndraf1, Nras, Nsfl1c, Pbl2, Pde12, Pex11a, Pik3c3, Pla2g12a, Pus7, Sco1, Sdr42e1, Sdr9c7, Slc16a7, Slc25a33, Smurf2, Tmem123, Tsc22d1, Ugt3a1
rank product method	1416021_a_at, 1418536_at, 1420570_x_at, 1428014_at, 1429466_s_at, 1430172_a_at, 1433107_at, 1434362_at, 1438009_at, 1438959_x_at, 1440016_at, 1440304_at, 1440587_at, 1441415_at, 1441539_at, 1442182_at, 1442725_at, 1443166_at, 1443377_at, 1443393_at, 1443719_x_at, 1444909_at, 1445001_at, 1445062_at, 1445492_at, 1445534_at, 1445612_at, 1445676_at, 1445728_at, 1446567_at, 1446614_at, 1449745_at, 1451626_x_at, 1452590_a_at, 1455930_at, 1456032_x_at, 1456973_at, 1457652_x_at, 1458382_a_at, 1458829_at, 1459557_at, 1459695_at, 1460132_at, 1500011K16Rik, 1810008I18Rik, 4930455C13Rik, 6030422H21Rik, Acacb, Acot3, Acss2, AFFX-18SRNAMur/X00686_3_at, Alas1, Ap1g2, Ap2a1, Arid5b, Asns, Atn1, BC094435, Bdkrb1, Ccl8, Ccne1, Ccrn4l, Cdk1, Cirbp, Cish, Coq10b, Crybb3, Ctrl, Cyp2c39, Cyp2g1, Cyp4a14, D530037H12Rik, D7Ert59e, Dbp, Defb1, Dhcr7, Egfl8, Erbb3, Fabp5, Fads1, Fbxl15, Fdps, Foxa1, Frat2, Gal3st1, Gale, Gapdhs, Ggt6, Gm129, Gm19313, Gnat1, Got1, Gpcpd1, Gpx3, Grem2, Guca1a, H2afz, Hhex, Hspb1, Id1, Idi1, Ifi27l2b, Igsf8, Il15, Isyna1, Lce3b, Lefty1, Lrrc24, Mettl20, Mt1, Myc, Nab2, Nipal1, Nrcc5, Nme4, Nmrk1, Nnmt, Nsdhl, Orm1, Orm2, Orm3, Pawr, Pcsk4, Pcsk9, Pdzk1ip1, Per1, Per3, Pfkfb1, Pgd, Plekhj1, Plxnb1, Pmvk, Pnpla5, Por, Prss8, Rhob, Rnd2, Rps13, Rundc3b, Sbk1,	0610012H03Rik, 1110004F10Rik, 1415788_at, 1415789_a_at, 1415790_at, 1416336_s_at, 1416461_at, 1416706_at, 1416811_s_at, 1416946_a_at, 1416947_s_at, 1417057_a_at, 1417714_x_at, 1417907_at, 1418274_at, 1418520_at, 1419443_at, 1420136_a_at, 1420867_at, 1420868_s_at, 1421534_at, 1422414_a_at, 1422660_at, 1422993_s_at, 1422997_s_at, 1423262_a_at, 1423263_at, 1423308_at, 1423309_at, 1423397_at, 1424300_at, 1424752_x_at, 1425665_a_at, 1426266_s_at, 1426394_at, 1426395_s_at, 1426419_at, 1427660_x_at, 1427932_s_at, 1428658_at, 1429040_at, 1429169_at, 1429252_at, 1429272_a_at, 1429468_at, 1429554_at, 1429712_at, 1429784_at, 1431158_at, 1433592_at, 1433622_at, 1434054_at, 1434161_at, 1434328_at, 1434568_at, 1434861_at, 1435137_s_at, 1435384_at, 1435574_at, 1435792_at, 1436247_at, 1436338_at, 1436933_at, 1436944_x_at, 1437026_at, 1437354_at, 1437547_s_at, 1438221_at, 1438398_at, 1438415_s_at, 1438596_at, 1438714_at, 1438786_a_at, 1439290_at, 1439305_at, 1439426_x_at, 1439463_x_at, 1439616_at, 1439831_at, 1440038_at, 1440219_at, 1440264_at, 1440284_at, 1440325_at, 1440373_at, 1440388_at, 1440490_at, 1440533_at, 1440551_at, 1440841_at, 1440896_at, 1441052_at, 1441108_at, 1441141_at, 1441328_at, 1441498_at, 1441505_at, 1441547_at, 1441765_at,

Scd1, Sdf2l1, Serpinh1, Sertad1, Slc23a2, Slc3a1, Slpi, Smad9, Soc52, Speer3, Sprr1a, Sqle, Tbc1d30, Tcf19, Thrsp, Tifa, Tmc5, Tmem159, Tmem71, Trib3, Ttc39a, Tubb2a, Tubb4a, Unc5b, Usp2, Vpreb3, Wbp2nl, Wfdc15b, Wnt5b, Zfp579, Zfp647, Zkscan17	1442049_at, 1442331_at, 1442916_at, 1443505_at, 1444333_at, 1444735_at, 1445966_at, 1446850_at, 1447502_at, 1448037_at, 1448462_at, 1449009_at, 1449855_s_at, 1450007_at, 1452052_s_at, 1452463_x_at, 1455418_at, 1455961_at, 1456433_at, 1457263_at, 1458637_x_at, 1459360_at, 1600002H07Rik, 1700066M21Rik, 1810058I24Rik, 2010315B03Rik, 2610002M06Rik, 2700089E24Rik, 2900026A02Rik, 4833417J20Rik, 4930506M07Rik, 4933411K20Rik, 5033411D12Rik, 6330564D18Rik, 9130221J18Rik, 9230114K14Rik, A130040M12Rik, A430035B10Rik, AA408650, AA987161, Abat, Abca1, Abcb4, Abcc2, Abcc3, Abcc9, Abce1, Abcg2, Abi3bp, Ablim3, Acaa1b, Acad11, Acad9, Acadm, Acat1, Acot1, Acot12, Acox2, Acsm3, Acsm5, Acy3, Adam17, Adap2, Adh4, Adra1b, Adss, Aggf1, Agl, Ahcyl1, Al314180, Al428301, Al464131, Aif1, Ak3, Akap1, Akr1c14, Akr1c20, Akr1d1, Alad, Alas2, Aldh3a2, Aldh5a1, Alg13, Alkbh7, Amd1, Amdhd1, Amy1, Anapc2, Anapc4, Angptl3, Anks4b, Anp32a, Anp32b, Aox1, Aox3, Apobec1, Apol7a, Aptx, Aqp4, Aqp9, Are1, Arf2, Arl1, Arl5a, Arl6ip1, Arl8b, Arpp19, Arrdc3, Arsg, Asb13, Asic5, Asna1, Asnsd1, Atf2, Atf6, Atf6b, Athl1, Atoh8, Atp11c, Atp6v1a, Atp7a, Atp8b1, AU041133, Avpr1a, AW209491, AW987390, Azin1, B630005N14Rik, Bag3, Bbox1, Bbs5, BC003331, BC018507, Bcap29, Bcl3, Bcl6, Bd1, Bet1, Bpgm, C1galt1, C1galt1c1, C1qb, C1s, C330007P06Rik, C4bp, C6, C730027H18Rik, C730036E19Rik, C8a, C8b, Caap1, Cab39, Cacybp, Caln1, Cand1, Canx, Cap1, Capn2, Capn7, Caprin1, Capza2, Car3, Car8, Casp8, Ccdc104, Ccdc115, Ccdc117, Ccdc126, Ccdc162, Ccdc59, Ccdc90b, Ccne2, Ccnf, Ccpq1, Ccs, Cd1d1, Cd24a, Cd2ap, Cd300lb, Cd36, Cd48, Cd68, Cd74, Cd86, Cdc123, Cdh2, Cela1, Cenpb, Ces1e, Ces1g, Ces2a,
--	---

Cetn3, Cfdp1, Chac2, Chdh, Chka, Chkb, Chmp2b, Chn2, Chordc1, Chpt1, Chrna2, Chtop, Chuk, Cideb, Cited2, Cks1b, Clcn2, Cldn1, Cldn2, Clec14a, Clec4f, Clock, Clpx, Cltc, Cmas, Cml2, Cml5, Cmtm4, Cnot7, Col27a1, Commd6, Copb1, Cops2, Cox15, Cox18, Cp, Cpeb4, Cpsf2, Cpt1a, Crip2, Crnk1, Csad, Csde1, Cse1l, Csf1r, Csgalnact2, Ctcf, Ctla2b, Ctnnb1, Ctnnbl1, Cts, Cul2, Cul3, Cul4a, Cul4b, Cxcl10, Cxcl9, Cybb, Cyp2a12, Cyp2c37, Cyp2c50, Cyp2c54, Cyp2d12, Cyp2d13, Cyp2d40, Cyp2j5, Cyp2j6, Cyp3a13, Cyp3a25, Cyp4v3, D10Ertd709e, D1Ertd448e, D5Ertd505e, D5Ertd798e, D830012I24Rik, D930016D06Rik, Daam1, Dancr, Dars, Dcaf11, Dcakd, Dct, Ddx10, Ddx3y, Ddx52, Ddx60, Decr1, Dennd4c, Derl2, Dhrs4, Dhx29, Dhx36, Dis3l, Dkc1, Dleu2, Dnaja1, Dnajb1, Dnajb4, Dnajb9, Dnajc14, Dnajc5, Dnase2a, Dnttip2, Dpm1, Dpys, Dr1, Dsg2, Dsp, Dtx4, Dync1i2, E130311K13Rik, Eaf1, Ear11, Eci2, Eea1, Efemp1, Efhd2, Efna1, Efr3a, Egln3, Ehhadh, Eif1a, Eif1b, Eif2ak2, Eif2b3, Eif2s3x, Eif3m, Eif4a2, Eif4g1, Eif5, Eif5b, Ell2, Elovl3, Emr4, Enpp2, Enpp5, Epc2, Ephb2, Ephx1, Eps8l2, Ergic2, Esf1, Etfdh, Eva1a, Exoc3, F9, Fabp2, Fabp7, Fahd1, Fam107b, Fam126b, Fam149a, Fam149b, Fam151b, Fam20a, Fam210a, Fam214a, Fam35a, Fam98a, Farsb, Fastkd5, Fbxo11, Fbxo30, Fcer1g, Fcho2, Fcna, Fez2, Fgf21, Fgfr1op, Fgfr1op2, Ficd, Fitm1, Fkbp3, Foxn2, Fpgt, Fpr2, Ftsj3, Fus, Fzd8, G0s2, G3bp1, G6pc, Gabarapl1, Gabpa, Gadd45a, Galnt1, Galnt2, Gas5, Gatc, Gbp2, Gbp2b, Gbp3, Gbp6, Gbp8, Gck, Gclc, Gclm, Gdf15, Gemin2, Ggcx, Ghr, Gimap9, Gjb1, Glcd, Glo1, Glt28d2, Gm17586, Gm19696, Gm19966, Gm20300, Gm20319, Gm2701, Gm4354, Gm6484, Gmfb, Gna12, Gna14, Gpatch11, Gpbp1, Gphn, Gpr155, Gpr97, Grpel2, Gspt1, Gtf2f2, Gtf2h1, Gtf2h2, Gtpbp10, Gtpbp4, Guk1, Gyk, H2-Aa, H2-Ab1, H2-Eb1, H2-T24, Hadha, Hccs, Hdac2, Hdhd3, Herc6, Hes1, Hexim1, Hif1a, Hmgb3, Hnf4a, Hnrnpa3, Hnrnpu, Hsd17b12, Hsd17b2, Hsd3b2, Hsd3b3, Hsdl2, Hsp90aa1, Hsp90ab1, Hsp90b1, Hspa13, Hspa1b, Hspb6, Hspb8, Hsph1, Ide, Ifi202b, Ifih1, Ifit1, Ifit2, Ift20, Igf5, ligp1, Ik, Ikbke, Ikbkg, Immt, Inhbc, Inhbe, Inpp5a, Insig2, Ints12, Ints2, Ints4, Irak2, Irf6, Irgm1, Irs1, Itfg1, Itga8, Jak1, Jmjd8, Jtb, Jun, Kat8, Kcnk5, Kctd15, Kdm1b, Kif5b, Klf10, Klf9, Klkb1, Kpna1, Kpna2, Kpna3, Kpnb1, Krr1, Lactb, Lamb3, Lamp2, Larp1, Larp7, Lars, Lcp1, Ldhd, Ldrl, Leprotl1, Lgals3bp, Lgals8, Lgals9, Lhpp, Lig4, Lima1, Lin7c, Lipg, Lonp2, Lpar6, Lrif1, Lrp5, Lrrc59, Lrrc8d, Lsm2, Ltb, Ltv1, Luc7l3, Lum, Ly86, Lyz2, Mad2l2, Mafb, Magix, Magt1, Mal2, Mamdc2, Map2k4, Mapk1, Mapk1ip1l, March8, Mars, Mbd2, Med19, Meiob, Metap2, Mettl3, Mia2, Mir22hg, Mis12, Mki67ip, Mlf2, Mlxipl, Mmd, Mme, Mmgt1, Mmp19, Mn1, Mob4, Mocs2, Moxd1, Mpdz, Mpzl2, Mrpl50, Mrs2, Msh6,

Mtap, Mtif2, Mtm1, Mtmr9, Mtpn, Mtrf1l, Mtus1, Mup10, Mx2, Naa15, Naa16, Naa50, Nadk2, Nae1, Nap111, Nbn, Ncbp2, Ncl, Ndnl2, Ndufaf1, Ndufaf3, Neat1, Necab1, Nek4, Nek6, Nelfcd, Nenf, Nfe2l2, Nfil3, Nhfp2, Nicn1, Nipa2, Nit2, Nln, Nlrp12, Nlrx1, Nmi, Nol8, Nolc1, Nop9, Nox4, Npr2, Nr1d1, Nr3c1, Nr5a2, Nras, Nrbf2, Nsf1c, Nt5dc1, Nuak2, Nudcd2, Nudt12, Nup35, Nxt2, Oasl2, Obp2a, Omd, Oplah, Osbp18, Pabpc1, Paip2, Papola, Pard6b, Parn, Parp9, Parva, Pbld1, Pbld2, Pbrm1, Pcbd2, Pcbp4, Pcdh1, Pcgf6, Pcsk6, Pde12, Pdiit, Pdlim5, Pdzk1, Pelo, Pex11a, Pex13, Pex26, Pfnd1, Pgap1, Pgap2, Pggt1b, Pgm3, Phf17, Phkb, Phldb2, Picalm, Pigr, Pik3r1, Pim1, Pitpna, Pitpn, Pja2, Pkp4, Pla2g12a, Pla2g15, Plekhb1, Plekhf1, Plin5, Plrg1, Plscr2, Pnn, Pno1, Pnpla7, Pnrc2, Polb, Ppap2b, Ppara, Ppcs, Ppp1r11, Ppp1r3b, Ppp1r3c, Ppp2ca, Ppp2r1b, Ppp2r5e, Ppp3ca, Ppp4r1, Ppp4r2, Ppp6c, Pprc1, Prkd3, Prkrir, Prorsd1, Prr16, Psap, Psma4, Psmc5, Psmc6, Psmd1, Psmd12, Psmd14, Ptp4a2, Ptpp12, Pum1, Pus7, Qpct, Rab10, Rab11a, Rab18, Rab24, Rab30, Rab6a, Rab7, Rad17, Rad21, Ranbp1, Ranbp6, Rap1b, Rars, Rbbp9, Rbm8a, Rbmxl1, Rchy1, Rdh16, Rdx, Rel11, Retsat, Rfwd2, Rgs5, Rhou, Rnf121, Rnf125, Rnf138, Rnf152, Rnf186, Rnf43, Rnf5, Rnf6, Rnft1, Rpp14, Rpp40, Rps4l, Rrp12, Rrp36, Rrp8, Rsad2, Rtfdc1, Rtn4rl1, Rtp3, S1pr1, Saa4, Sc5d, Scaf11, Scnm1, Sco1, Scoc, Sdr42e1, Sdr9c7, Sec16b, Sec62, Sec63, Senp7, Serinc1, Serpina12, Serpina3a, Serpina3k, Serpina4-ps1, Serpina9, Sfr1, Sgk2, Sgms2, Sgp1, Sgpp1, Siae, Slc15a3, Slc16a12, Slc16a13, Slc16a7, Slc17a2, Slc17a3, Slc22a1, Slc22a5, Slc22a7, Slc25a13, Slc25a16, Slc25a25, Slc25a33, Slc30a10, Slc33a1, Slc35a1, Slc35a3, Slc35b4, Slc39a8, Slc41a2, Slc45a3, Slco1a1, Slco1b2, Smad6, Smarca2, Smek2, Smg7, Smim8, Smn1, Smurf2, Smyd2, Snap23, Snhg12, Snhg6, Sntb1, Snx24, Snx5, Sorbs1, Sos2, Sp3, Spast, Spg20, Spop, Srd5a1, Srp68, Srr, Srsf1, Srsf10, Srsf3, Srsf7, Ssb, Stag1, Stap1, Stard13, Stat2, Stat5a, Stbd1, Stip1, Stk16, Strn3, Stx4a, Stxbp6, Sub1, Suchr1, Sult1c2, Sumf1, Suox, Syncrip, Tab2, Taf1a, Tardbp, Tars, Tbcel, Tbk1, Tbl1x, Tcaim, Tcea1, Tceb1, Tcerg1, Tefm, Tenc1, Tenm3, Tfec, Thap11, Thrap3, Timd4, Tipin, Tk2, Tm9sf3, Tmbim1, Tmcc1, Tmem123, Tmem184b, Tmem230, Tnfaip8l1, Tnpo3, Top2b, Tpmt, Tpp2, Trappc10, Treml4, Triap1, Trim39, Trmt10a, Trmt6, Trp53inp2, Tsc22d1, Ttc27, Ttc30b, Ttc32, Tvp23b, Twsg1, Txndc9, Tyw5, Uba3, Uba5, Ube2d1, Ube2e3, Ube2g2, Ube2l6, Ube2v2, Ubxn2a, Ubxn4, Ufd1l, Ufl1, Ugt2b34, Ugt2b35, Ugt3a1, Ulk2, Uso1, Usp18, Usp33, Utp6, Vcl, Vma21, Vprbp, Vps35, Vps41, Vwa8, Wbscr16, Wdr43, Wdr77, Wnt2, Xrn2, Xylb, Yap1, Ythdc2, Ythdf2, Ythdf3, Zbtb20, Zcchc10, Zcchc6, Zfand1,

	Zfand2a, Zfp110, Zfp141, Zfp146, Zfp260, Zfp263, Zfp326, Zfp361, Zfp386, Zfp445, Zfp51, Zfp58, Zfp617, Zfp62, Zfp825, Zfp866, Zfp932, Zfp935, Zfp946, Zfp955b, Znhit6, Znrd1, Zranb2, Zscan29, Zswim7, Zwint
CR and quercetin diet	
rank	1425222_x_at, 1427820_at, 1430172_a_at,
product	1437867_at, 1439816_at, 1440587_at,
method	1440815_x_at, 1442025_a_at, 1442026_at, 1442420_at, 1443322_at, 1445591_at, 1447901_x_at, 1453023_at, 1456973_at, 1457549_at, 1810062G17Rik, 3110043O21Rik, 3110045C21Rik, 4933426M11Rik, 5330406M23Rik, 9130221J18Rik, Adora1, AFFX-18SRNAMur/X00686_3_at, AI195470, Akap12, Alas1, Apbb3, Arhgef26, Arid5b, Atp10d, Baiap2l1, BC027231, Bnip3, Btg2, Ccrn4l, Cd14, Cd163, Cebpb, Cers6, Chac1, Cib3, Cish, Col6a1, Coq10b, Ctgf, Cyp2b10, Cyp4a14, D030063E12, D1Ert622e, Dusp14, Egr1, Epas1, Epcam, Errfi1, Fam134b, Fbxl12, Fbxo31, Fkbp5, Fn1, Foxa1, Frat2, Fst, Gadd45g, Gfpt2, Gja4, Gm129, Gm15698, Gnat1, Got1, Gpcpd1, Gps2, Grem2, Hhex, Hspb1, Id1, Ier3, Ifi2712b, Igfbp1, Il1r1, Il1rn, Il6ra, Ivns1abp, Kctd12, Klf2, Lonrf1, Lpin1, Lyve1, Map3k5, Mast3, Mettl20, Mfsd2a, Mfsd7c, Mgp, Mknk2, Mlxip, Mt1, Myadm, Myc, Nab2, Nlrc5, Nnmt, Nop56, Nrg4, Orm2, Orm3, Osmr, P2ry2, Pcfg1, Pcsk4, Per1, Plin4, Plk3, Por, Ppip5k2, Ppl, Rab20, Rasgef1b, Rhbdf1, Ripk4, Sds, Serinh1, Sertad1, Sgk1, Sirt1, Slc25a22, Slc25a32, Slc25a51, Slc43a1, Slc7a2, Slc8b1, Sorbs3, Sprr1a, Sqle, Srsf5, St3gal5, Sult1e1, Susd2, Tacc2, Tagln, Tbc1d8, Tcp11l2, Thrsp, Tk1, Tns1, Tpm4, Trib3, Trim47, Ttc39a, Tubb2a, Tuft1, Usp2, Vcam1, Vim, Vsig10, Wsb1, Zbtb16, Zfand4, Zfp275
	1418188_a_at, 1418189_s_at, 1420310_at, 1424853_s_at, 1425918_at, 1426327_s_at, 1434280_at, 1435267_at, 1436389_at, 1437162_at, 1437600_at, 1437821_at, 1438265_at, 1438403_s_at, 1438543_at, 1438714_at, 1439175_at, 1439305_at, 1439307_at, 1439600_at, 1439832_at, 1440314_at, 1440790_x_at, 1440841_at, 1441413_at, 1441498_at, 1442049_at, 1442109_at, 1442470_at, 1443003_at, 1443056_at, 1443075_at, 1443147_at, 1443505_at, 1443673_x_at, 1443949_at, 1444001_at, 1444326_at, 1445267_at, 1445402_at, 1446850_at, 1447227_at, 1447329_at, 1451477_at, 1452433_at, 1455961_at, 1456159_at, 1456386_at, 1456851_at, 1456960_at, 1457020_at, 1457367_at, 1457988_at, 1457991_at, 1458134_at, 1458205_at, 1458637_x_at, 1458832_at, 1459008_at, 1459061_at, 1459238_at, 1459948_at, 1700023H06Rik, 2310007O11Rik, 2310067E19Rik, 2610507B11Rik, 2900064B18Rik, 3110005L24Rik, 4633401B06Rik, 4930459C07Rik, 4930480G23Rik, 5730414N17Rik, 9030622O22Rik, A930004J17Rik, Adck3, Adh6-ps1, Adipor2, Agpat2, Al464131, Anks4b, Apob, Aqp4, Aqp8, Arrdc3, Arsg, B230114P17Rik, Bag3, Bbs5, BC038331, Bcl6, C430003N24Rik, C730029A08Rik, Car3, Ccdc117, Ccdc58, Cd300lb, Cda, Cflar, Chka, Chordc1, Chrna2, Cited2, Cks1b, Cldn1, Cldn3, Ctcf, Cyp2d12, Cyp2d13, Cyp2d40, Cyr61, D5Wsu152e, Daam1, Ddx6, Decr2, Dhdh, Dhx36, Dhx58, Dnajb1, Dnajb14, Dnajb4, Dnajb9, Dusp6, E130110O22Rik, Ear11, Efna1, Eph2, Evi5, Fahd1, Fam214a, Fam26f, Fam89a, Fermt2, Fgf21, Ficd, Fpgs, G0s2, G6pc, Gadd45a, Gck, Gdap10, Gdf15, Ggact, Gm11772, Gm19696, Gm2788, Gm4354, Gm6484, Gna14, Gpr97, Gprin3, H2-T24, Hacl1, Hectd1, Hes1, Hexim1, Hist1h1c, Hist1h4i, Hps5, Hsd3b3, Hspa13, Hspa1b, Hyou1, Ift20, Igfals, Inhbe, Insig1, Irs1, Jun, Kcnk5, Keap1, Klf10, Lipg, Lmo2, Lnx2, LOC75771, Lppos, Lrfn3, Lrp5, Luc7l3, Lysmd3, Magix, Mis12, Mme, Mn1, Mpst, Mup10, Nbeal1, Nfatc3, Nfib, Nfil3, Nr1d1, Nrep, Obp2a, Omd, Osbp18, Osgin1, Paqr9, Parp16, Pbl1, Pcdh1, Pde6d, Pdpr, Pglyrp1, Phf17, Plcxd2, Plekhf1, Plscr2, Ppara, Ppp1r3b, Ppp1r3c, Prkd3, Prlr, Prox1, Pter, Pum1, Rab30, Rbm25, Rfc1, Rfesd, Rgs5, Rnf125, Rnf152, Rnf186, Rnf43, Rps4l, Rrp36, Rtp3, S1pr1, Sc5d, Secisbp2l, Serpina9, Sgpl1, Skiv2l, Slc1a4, Slc25a30, Slc26a1, Slc30a10, Slc40a1, Slc45a3, Smarca4, Smyd4, Srd5a1, Stap1,

		Stard13, Sucnr1, Sult1c2, Taf1a, Tbc1, Tbl1x, Temm3, Thap11, Thrb, Timd2, Tjp3, Tmcc1, Tmem98, Tmie, Trim2, Trim39, Tsc22d1, Ttc30b, Ube2d1, Ulk2, Unc119, Vnn1, Wbscr16, Wnk1, Wrap73, Zbed6, Zbtb11os1, Zfp322a, Zfp36l1, Zfp551, Zfp62
CR and quercetin administered intraperitoneal		
moderated t-test	Arg1, Cyp2b13, Etnppl, Fmo3, Fst, Gstt3, Hamp2, Sh2d4a, Slc22a27, Tat	Adh4, Alad, C9, Crot, Cyp4a12a, Dio1, Hsd3b5, Hsd3b7, Hspa5, Ifi47, Ifit1, Igtp, Irgm2, Isg15, Ldha, Manf, Mbl1, Mgst1, Nat8, Ndufaf1, Pter, Rarres1, Serpina11, Slc16a7, Tbrg1, Tm9sf3, Ttpa, Ugt2b35
rank product method	1600016N20Rik, 8430408G22Rik, Acly, Acot7, Arl4d, Atp10d, Ccdc57, Cdc42bpq, Cish, Cpeb2, Crym, Cux2, Cyp17a1, Cyp2b13, Cyp2c39, Cyp4a10, Cyp4a14, Cyp4a31, Dak, Ddb1, Elov16, Eno1, Erdr1, Fads1, Fads2, Fmo3, Fnip2, Foxa1, Gadd45g, Gm5215, Gnat2, Gng11, Gpihbp1, Gsta4, Gstm2, Hlx, Igfbp1, Inmt, Irs2, Lce1i, Lepr, LOC100044177, LOC100046136, Lor, Lrrc24, Lss, Myh14, Nnmt, Nrg4, Pklr, Pnpla5, Ppp1r3g, Rnf39, Rpl32, Sdsl, Sh2d4a, Slc22a26, Slc22a27, Slc2a4, Smoc2, Snord15a, Snord22, Snord35a, Snord82, Snord87, Soc52, Spen, Sult1e1, Sult2a3, Tacc2, Tbc1d19	0610009B22Rik, 1600012H06Rik, 1810011O10Rik, 3010026O09Rik, 4931406C07Rik, Abhd3, Acat2, Acox2, Actg1, Adh4, Adprhl2, Agpat2, Akirin1, Akr1c14, Alad, Aldh1a1, Amacr, Amy1, Angptl3, Ankrd46, Anp32b, Aox1, Ap3s1, Aqp8, Arrdc3, Arsg, Asnsd1, Atg5, Avpr1a, AW112010, Bbox1, Bcap29, Bcl6, Bhmt2, C1galt1c1, C1qa, C1s, C4b, C4bp, C8a, C8b, C9, Cacybp, Ccm2, Cd36, Cd74, Cdh2, Ces1d, Ces3a, Chac2, Cks1b, Clpx, Cmas, Cmbl, Cml5, Cmtr1, Commd3, Crot, Csad, Csrp3, Ctcf, Ctss, Cxcl10, Cxcl9, Cyp2c29, Cyp2c50, Cyp2d40, Cyp2j5, Cyp4a12a, Cyp4a12b, Cyp4v3, Cyp7b1, D630039A03Rik, Dexi, Dhrs3, Dio1, Dnaja1, Dnajb2, Dsp, E2f8, Ech1, Ei24, Eif2ak2, Elov13, Emr4, Ephx1, Ergic1, Etfdh, Eva1a, Fam98b, Fez2, G0s2, Gbp3, Gclc, Gdf15, Gfod2, Ghr, Glcd, Glyat, Gm5150, Gm561, Gm6484, Gp49a, Gstm4, Guk1, H2-Aa, H2-Ab1, Hmgb3, Hmgcs2, Hnf4aos, Hopx, Hpgd, Hsd17b2, Hsd17b4, Hsd3b5, Hsd3b7, Hspa5, Id2, Ide, Ifi35, Ifi47, Ifit1, Igfals, Igtp, Il6st, Immp1l, Inca1, Inhba, Insig2, Ints2, Irgm2, Isg15, Kdm1b, Keg1, Klf10, Klh2, Lamtor5, Lbp, Lcat, Lcn2, Ldha, Lin7a, Lonp2, Lpgat1, Lrfn3, Ly6d, Ly86, Mafb, Magoh, Mal2, Manf, March8, Mbl1, Mcts2, Med21, Mlxipl, Mmd, Mocos, Mrpl50, Mup20, Mup21, Mup4, Mup6, Ndufaf1, Nhirc1, Nubp2, Paqr7, Parg, Parp9, Pcnp, Pdia4, Perp, Pex26, Picalm, Pkp4, Plekhf1, Pno1, Pnpla7, Polb, Ppid, Psmd2, Psmd1, Psmd6, Pter, Ptpla, R3hdm2, Rab1, Rap1a, Rarres1, Rnf7, Rraga, Rtp4, S100a10, Saa3, Saa4, Scyl3, Sdc1, Sdr42e1, Sec14l2, Sec22b, Sec23b, Serbp1, Serpina11, Serpina3n, Serpina7, Serping1, Sgk2, Sgpl1, Sgpp1, Slc10a3, Slc13a3, Slc16a2, Slc16a7, Slc1a4, Slc22a7, Slc25a45, Slco1a1, Slco1b2, Smim15, Snap47, Snhg6, Snrpe, Soc53, Spon2, Spryd7, Srd5a1, Stbd1, Sucnr1, Syap1, Tbrg1, Timd4, Tipin, Tjp3, Tmem19, Tmem254a, Tmem30a, Tmem56, Tmie, Tomm20, Tomm70a, Trex1, Trp53inp2, Tsc22d1, Ttc32, Ttpa, Tyrobp, Ube2e1, Ube2l6, Ugt2b35, Ugt3a1, Usp18, Vps37b, Zbtb18, Zbtb20, Zfand2a
CR and genistein administered intraperitoneal		
moderated t-test	Acly, Cyp17a1, Eno1, Gsta4, Gstm2, Hamp2, Hlx, Igfbp1, Lrrc24, Lss, Nipal3, Nnmt, Ppp1r3g, Slc2a4	4931406C07Rik, Acaa1b, Acs1l1, Adh4, Ak2, Alad, Aldh3a2, BC025446, Bcap29, C9, Ces1d, Cfah, Crot, Cryz, Csad, Csrp3,

		Cyp2j5, Cyp4a12a, Cyp7b1, Dio1, Dnaja1, Dsp, Eif2ak2, Ephx2, Etfdh, Fgf21, G0s2, Gbp3, Gdf15, Hnf4aos, Hrsp12, Hsd17b4, Hsd3b5, Hsd3b7, Hspa5, Hspa8, Ifi35, Ifi47, Ifit1, Igtp, Inca1, Irgm2, Isg15, Lcn2, Ldha, Lin7a, Lipa, Lonp2, Lpgat1, Mal2, Manf, March8, Mbl1, Mgst1, Mien1, Mmd, Nadk2, Ndufaf1, Ostc, Pafah1b1, Papola, Paqr7, Phf17, Psma5, Pter, Rarres1, Rtp4, S100a10, Sdc1, Sdr42e1, Serbp1, Serpina11, Serpinb1a, Sh3bg12, Skp1a, Slc16a1, Slc16a7, Slc25a33, Supt16, Tbrg1, Tm9sf3, Tmem56, Tsc22d1, Ttpa, Ugt2b35, Ugt2b5, Ugt3a1, Usp18
moderated t-test, fold change > 1.5	Fmo3	
rank product method	1600016N20Rik, Acly, Acot3, Agpat9, Arg1, Atp10d, Casp3, Ccdc57, Cdc42bpg, Cish, Clmn, Cpeb2, Crym, Cux2, Cyp17a1, Cyp2b13, Cyp2b9, Cyp2c39, Cyp4a14, Cyp4a31, Dbp, Dntt, E130012A19Rik, Eif4ebp3, Elovl6, Eno1, Erdr1, Etnppl, Fbxo31, Fmo3, Fnip2, Fst, Fzd4, Gadd45g, Gm129, Gm5215, Gnat2, Gng11, Got1, Gpcpd1, Gstt3, Id1, Igfbp1, Inmt, Irs2, LOC100046136, Lss, Map3k5, Mfsd2a, Nrg4, Pdk4, Pdxp, Plin4, Ptgds, Rarb, Rpl32, Sdsl, Sh2d4a, Slc16a5, Slc22a26, Slc22a27, Slc37a4, Slc43a1, Snora21, Snord22, Snord82, Soc5s2, Sptssa, Sult1e1, Sult2a3, Tacc2, Tamm41, Tat, Tbc1d19, Unc93a	3010026O09Rik, Acat2, Adh4, Adprhl2, Aes, Akirin1, Alad, Alas2, Amacr, Amy1, Ankrd46, Anp32b, Aox1, Ap3s1, Apon, Avpr1a, Bag3, Bbox1, Bcl6, Bhmt2, C1galt1c1, C1qa, C1s, C4b, C4bp, C9, Cacybp, Ccdc86, Cd5l, Cdhd2, Ces1d, Ces2a, Ces3a, Cfd, Chac2, Clpx, Cml5, Cmtr1, Colec10, Cris1, Crot, Csrp3, Ctss, Cxcl9, Cyp2c29, Cyp2d40, Cyp4a12a, Cyp4a12b, Cyr61, Dcakd, Dhrs3, Dio1, Dnaic1, Dnaja1, Dnajb1, Dnajb2, Ech1, Ei24, Elovl3, Ephx1, Fez2, Foxa3, G0s2, Gfod2, Glcd, Glyat, Gm5150, Gm6484, Gstm4, Guk1, Hmgb3, Hmgcs2, Hsd3b5, Hsd3b7, Hspa13, Hspa5, Hsph1, Ifi27l2a, Ifi35, Ifi47, Ifit1, Igfals, Igtp, Il6st, Inca1, Irgm2, Isg15, Keg1, Lamtor5, Lbp, Lcat, Lcn2, Ldha, Leap2, Lipa, Lrfn3, Lum, Ly6d, Ly6e, Mad2l2, Magoh, Manf, Mbl1, Med21, Mmp15, Mocos, Mpst, Mup20, Mup21, Mup6, Nat8, Ndufaf1, Oasl1, Ocel1, Paqr7, Pcnp, Pex26, Plcxd2, Pnpla7, Polb, Ppp1r3b, Psmc2, Psmd1, Psmd6, Pter, Rarres1, Reep6, Rhbg, Rraga, Sdc1, Sec14l2, Serpina11, Serpina3n, Serpina7, Serping1, Sgk2, Sgpp1, Slc13a3, Slc16a7, Slc1a4, Slc22a1, Slc22a7, Slc25a45, Slc6a13, Slco1b2, Snap47, Spon2, St3gal3, Stbd1, Sucnr1, Syap1, Tbrg1, Timd4, Tjp3, Tmem254a, Tmie, Tomm20, Trex1, Tsc22d1, Ttpa, Tyrobp, Ube2e1, Ube2l6, Ugt2b35, Ugt3a1, Usp18, Wdtc1, Zbtb20

C: All significantly regulated genes by caloric restriction which are identified in at least one approach sorted by alphabet

Upregulated CR genes (gene symbols known)

0610007P14Rik, 1110001J03Rik, 1110007C09Rik, 1110012L19Rik, 1110020A21Rik, 1110034G24Rik, 1190002N15Rik, 130004C03, 1500011K16Rik, 1500032P08Rik, 1700001J11Rik, 1700018L02Rik, 1700024P16Rik, 1700040L02Rik, 1810008I18Rik, 1810010H24Rik, 1810032O08Rik, 1810046K07Rik, 1810053B23Rik, 1810062G17Rik, 2010012O05Rik, 2010107E04Rik, 2210013O21Rik, 2310007J06Rik, 2310034O05Rik, 2310047D07Rik, 2310061I04Rik, 2510039O18Rik, 2610002J02Rik, 2610002M06Rik, 2900009J06Rik, 2900040J22Rik, 3000002C10Rik, 3010003L21Rik, 3110043O21Rik, 3110045C21Rik, 4631423B10Rik, 4732419C18Rik, 4833420G17Rik, 4921506L19Rik, 4921517O11Rik, 4930452B06Rik,

4930455C13Rik, 4930516B21Rik, 4930588G05Rik, 4931408D14Rik, 4932439E07Rik, 5330406M23Rik, 5430421B17, 5730411F24Rik, 6030422H21Rik, 6430562O15Rik, 6430573F11Rik, 6720427H10Rik, 8430408G22Rik, 8430419L09Rik, 9230104K21Rik, 9430078K24Rik, 9530006C21Rik, 9530025L08Rik, 9530046B11Rik, 9630009C16, 9630019E01Rik, 9930014A18Rik, A4gnt, A530079E22Rik, A930026I22Rik, A930036K24Rik, AA415014, AA415398, AA675035, Aacs, Aard, Abca3, Abca7, Abca8a, Abcb1a, Abcb1b, Abcc4, Abcc6, Abcd2, Abcg1, Abcg5, Abcg8, Abhd14b, Abtb2, Acaca, Acacb, Acadl, Acap3, Ackr1, Acly, Acmsd, Aco2, Acot13, Acot3, Acot7, Acot9, Acox1, Acp6, Acpp, Acsi3, Acsi4, Acsi5, Acss2, Acta1, Acta2, Actn1, Actr6, Acvr2b, Acy1, Acyp2, Adam8, Adam9, Adamts7, Adarb1, Adck4, Adcy6, Adcy9, Adgrl1, Adnp, Adora1, Adprm, Adrb2, Adrb3, Adssi1, Agap2, Agpat1, Agpat6, Agpat9, AI118078, AI132709, AI195470, AI413582, AI842136, Aifm1, Aim1, Aimp1, Ak2, Ak4, Akap12, Akap8, Akap8l, Akr1b3, Akr1b7, Akr1c18, Alas1, Alb, Aldh1a7, Aldh1b1, Aldh1l1, Aldh4a1, Aldoa, Aldoc, Alkbh6, Alox15, Als2cl, Amer2, Amigo2, Amotl2, Amy2a5, Angel2, Angptl4, Ank, Ank3, Ankrd16, Ankrd23, Ankrd27, Anks1, Antxr2, Anxa3, Anxa4, Anxa5, Anxa6, Ap1g2, Ap2a1, Apip, Apoo, App, Aprt, Aqp11, Arg1, Arhgap18, Arhgap21, Arhgdia, Arhgdib, Arhgef16, Arhgef26, Arhgef5, Arid1a, Arid5b, Arl15, Arl3, Arl4d, Arl6, Armc10, Arrdc1, Arrdc2, Arrdc4, As3mt, Asah1, Asf1b, Asl, Asns, Aspg, Asrgl1, Ass1, Atad3a, Atg14, Atg16l2, Atg2a, Atg9a, Atl2, Atn1, Atp10d, Atp13a1, Atp13a2, Atp1a1, Atp1b1, Atp4b, Atp5a1, Atp5c1, Atp5f1, Atp5g1, Atp5l, Atp5o, Atp5sl, Atp6ap2, Atp6v0a1, Atp6v0d1, Atp6v0e2, Atp6v1d, Atp6v1g1, AU015603, AU021025, AU040320, Avil, Avpi1, AW549877, AW555355, Awat1, Aym1, B230214O09Rik, B430203I24Rik, B9d1, B9d2, Bag2, Bag6, Baiap2l1, Banf1, Banp, BC021614, BC021891, BC023829, BC027231, BC038156, BC049762, BC089597, BC094435, Bcat2, Bckdhb, Bcl10, Bcl6b, Bdh2, Bdkrb1, Bex4, Blhfb9, Blhhe40, Bivm, Blvra, Bmi1, Bmp, Bnip1, Bnip3, Brca1, Brca2, Bri3, Bscl2, Bst2, Btg1, Btg2, C1qtnf1, C3, C730049O14Rik, C77027, C77080, C77097, C78692, C79242, Calm2, Calml4, Camk1d, Camsap3, Caprin2, Car2, Card10, Card6, Casp3, Casp6, Catsperd, Cbr1, Cbx7, Cc2d2a, Ccbl2, Ccdc152, Ccdc28b, Ccdc57, Ccdc61, Ccdc68, Ccdc84, Ccl21a, Ccl24, Ccl8, Ccnd3, Ccndbp1, Ccne1, Ccng2, Ccnk, Ccnl2, Ccp110, Ccrn4l, Ccz1, Cd14, Cd163, Cd209f, Cd55, Cd63, Cd7, Cd9, Cd99, Cdc25b, Cdc42bpg, Cdc42ep2, Cdc42ep5, Cdcp1, Cdh1, Cdip1, Cdk1, Cdk18, Cdk5rap1, Cdk8, Cdkn1c, Cdkn2c, Cdt1, Ceacam1, Ceacam9, Cebpb, Cela2a, Celsr1, Cenpa, Cenpq, Cep19, Cep85l, Cers6, Ces2g, Cetn2, Cfap126, Cfap20, Cfl1, Chac1, Chaf1b, Chchd4, Chchd6, Chchd7, Chd1, Chd1l, Chd2, Chd3, Chd7, Chic2, Chid1, Chmp2a, Chp1, Chpf, Chrd, Chrna4, Ciart, Cib3, Cic, Cir1, Cirbp, Cish, Ciz1, Cks2, Clasrp, Clk1, Clk4, Clmn, Cln6, Cln3, Clybl, Cmya5, Cnn3, Cnot3, Cnppd1, Cnst, Cntrl, Coa6, Cobll1, Col1a2, Col3a1, Col4a2, Col6a1, Col6a2, Col6a3, Commd1, Commd10, Commd4, Cope, Coprs, Coq10b, Coq6, Coq7, Coro7, Cox5a, Cox7a1, Cox7b, Cpeb2, Cpm, Cpne2, Cpne3, Cps1, Cpsf4, Cpsf7, Cr1l, Crcp, Crebrf, Crip1, Crtap, Cry1, Cry2, Crybb3, Crybg3, Cryl1, Crym, Cs, Csnk1d, Csrp1, Csrp2, Ctcflos, Ctdsp2, Ctgf, Cth, Ctif, Ctnnbip1, Ctps2, Ctrl, Ctse, Ctsk, Ctsl, Ctsw, Cuta, Cux2, Cxcl13, Cyb5r1, Cyb5r3, Cyc1, Cycs, Cyp17a1, Cyp20a1, Cyp26a1, Cyp2b10, Cyp2b13, Cyp2b23, Cyp2b9, Cyp2c38, Cyp2c39, Cyp2c55, Cyp2c66, Cyp2d11, Cyp2d22, Cyp2g1, Cyp2j9, Cyp2r1, Cyp2t4, Cyp39a1, Cyp3a57, Cyp4a10, Cyp4a14, Cyp4a31, Cyp4b1, Cyp51, Cystm1, Cyth1, D030056L22Rik, D030063E12, D12Ertd123e, D130037M23Rik, D130043K22Rik, D130062J21Rik, D18Ertd169e, D18Ertd232e, D1Ertd622e, D4Ertd298e, D530037H12Rik, D5Ertd215e, D630033O11Rik, D7Ertd187e, D7Ertd59e, D9Wsu90e, Dad1, Dap, Dap3, Dapk1, Dazap1, Dbi, Dbp, Dbt, Dcbl1, Dclk3, Dctn1, Dctpp1, Ddah2, Ddb1, Ddb2, Ddhd2, Ddit4, Ddost, Ddt, Ddx1, Ddx46, Defa24, Defa-rs12, Defb1, Dhcr7, Dhrs7, Dhrs8, Dhx16, Dicer1, Dio3os, Dixdc1, Dlat, Dlg3, Dli1, Dlx6os2, Dmpk, Dmxl2, Dnajb6, Dnajc13, Dnmbp, Dntt, Dock4, Dock6, Dot1l, Dqx1, Dsg1c, Dus1l, Dusp1, Dusp14, Dync1h1, Dync1li1, Dync2h1, Dync2li1, Dynll1, Dynlt3, E030016H06Rik, E130012A19Rik, E230001N04Rik, E2f5, E330018D03Rik, Eaf2, Ebp, Echdc1, Echdc2, Ecm1, Ecscr, Edem1, Edn1, Eef2, Eef2k, Eglf8, Egr1, Ehmt1, Eid1, Eif1, Eif1-ps3, Eif2a, Eif2b1, Eif3c, Eif3j1, Eif4e3, Eif4ebp1, Eif4ebp3, Elovl6, Emc2, Emc7, Emcn, Emilin1, Eml2, Emp2, Enc1,

Upregulated CR genes (gene symbols known)

Eno1, Eno3, Enpep, Enpp1, Entpd2, Epas1, Ebpb4.1, Ebpb41l4aos, Epcam, Epg5, Ephb8, Ephb4, Epn1, Epor, Eppk1, Eps15, Erbb3, Erdr1, Errfi1, Esam, Esrra, Esyt2, Ethe1, Etnk2, Etnpl, Evx2, F7, Fabp5, Fads1, Fads2, Fah, Fam103a1, Fam110c, Fam117a, Fam120a, Fam126a, Fam134b, Fam13a, Fam167b, Fam177a, Fam179a, Fam180a, Fam193b, Fam213a, Fam3c, Fam46a, Fam53b, Fam58b, Fam60a, Fam63b, Fam76b, Fam98c, Fars2, Fasn, Fastkd1, Fastkd3, Fat1, Fbxl12, Fbxl15, Fbxl4, Fbxl6, Fbxo18, Fbxo21, Fbxo31, Fbxo33, Fbxo47, Fbxo9, Fcgr2b, Fchsd2, Fdlt1, Fdps, Fdxr, Fech, Fes, Fgd6, Fh1, Fhit, Flh1, Fkbp14, Fkbp5, Flcn, Flot1, Flt4, Fmn1, Fmo2, Fmo3, Fmo4, Fmo5, Fn1, Fnip2, Foxa1, Foxo1, Foxo3, Frat2, Frmd4b, Fscn1, Fst, Fuom, Furin, Fzd4, Fzd5, G6pdx, Gab1, Gadd45g, Gal3st1, Gale, Galt, Gapdh, Gapdhs, Garem, Gba, Gbe1, Gca, Gdap2, Gemin6, Gfm1, Gfpt2, Gfra1, Ggt6, Ggt7, Gimap4, Gja4, Gkap1, Glb1, Glrx, Gls2, Glud1, Gm10029, Gm10052, Gm10144, Gm10845, Gm11827, Gm12657, Gm12866, Gm13194, Gm13375, Gm14325, Gm14403, Gm14420, Gm15408, Gm15453, Gm15666, Gm15698, Gm15754, Gm15772, Gm15800, Gm16381, Gm16519, Gm19313, Gm19648, Gm19929, Gm21541, Gm2889, Gm3579, Gm3715, Gm4825, Gm4910, Gm5039, Gm5131, Gm5177, Gm5215, Gm5321, Gm5560, Gm5591, Gm5600, Gm5643, Gm5779, Gm5905, Gm6981, Gm7634, Gm826, Gm8580, Gm9202, Gm9222, Gm9391, Gm9405, Gm9731, Gm9798, Gml, Gmnn, Gnat1, Gnat2, Gne, Gng10, Gng11, Gng3, Gnpda1, Gnptab, Gns, Golin4, Got1, Gpam, Gpcpd1, Gpd1l, Gpi1, Gpihbp1, Gpn3, Gpr108, Gpr157, Gpr182, Gps2, Gpx3, Gpx6, Gpx7, Grem2, Grina, Grm8, Grn, Grsf1, Gspt2, Gsta2, Gsta4, Gstk1, Gstm1, Gstm2, Gstm3, Gstm5, Gstm6, Gstm7, Gsto1, Gstt2, Gstt3, Gtdc1, Gtf2a2, Gtf3c3, Gtpbp2, Guca1a, Gucd1, Gucy1b3, Gucy2c, Gusb, Gyg, Gypc, Gys2, H2afv, H2afz, H2-M10.5, Hal, Hao2, Hapl1n1, Hapl1n4, Hat1, Haus7, Hax1, Hbs1l, Hcls1, Hdac5, Hdac7, Hdhd2, Herpud1, Herpud2, Hexb, Hey1, Hgd, Hhex, Hic2, Hint1, Hist1h4c, Hjurnp, Hlx, Hmgcr, Hmgcs1, Hmgn2, Hmgn5, Hmox1, Hnrnpa1, Hnrnpa1, Hnrnpa2b1, Hnrnpdl, Hnrnph1, Hnrnpl, Hnrnpll, Hnrnpm, Hook2, Hoxa5, Hoxc6, Hpcal1, Hsbp1l1, Hsf4, Hspa2, Hspa9, Hspb1, Iah1, Iars2, Icam1, Id1, Idh1, Idh2, Idh3b, Idh3g, Idi1, Ier3, Ifi27l2b, Ifi30, Ifitm1, Ifna2, Ifna7, Ifngr1, Ifnz, Ifrd1, Ift27, Ift43, Igf2, Igfbp1, Igfbp5, Igsf23, Igsf8, Il10rb, Il11, Il15, Il15ra, Il17ra, Il17rb, Il18, Il1f9, Il1r1, Il1rn, Il33, Il34, Il6ra, Imp3, Impa2, Inmt, Inpp4a, Inpp5j, Ins2, Ip6k2, Ipmk, Ireb2, Irs2, Irx3, Isca1, Isyna1, Itga1, Itga6, Itgb1bp1, Itgb2, Itih5, Itpka, Itpr1, Ivd, Ivns1abp, Izumo4, Jup, Kansl3, Kat2b, Kcnn2, Kcnq1, Kcns1, Kctd12, Kdelc1, Kdm3a, Kdm6b, Kdr, Khdc3, Khk, Khnyn, Kif4, Kin, Klc4, Klf1, Klf13, Klf15, Klf2, Klf6, Klhdc2, Klhl21, Klhl24, Klhl3, Klk15, Klk1b4, Klrg2, Krcc1, Krt23, Krtcap2, L2hgdh, Lacc1, Lactb2, Lamc1, Lancl1, Lap3, Las1l, Lbr, Lce1i, Lce3b, Lce3c, Lclat1, Ldb1, Ldhb, Lect1, Lect2, Lefty1, Leng8, Lepr, Lgals4, Lgalsl, Lgmn, Lhfp, Lig1, Lims2, Lipo1, Llgl1, Llgl2, Lmf2, Lmntd2, Lmo1, Lmo4, Lmo7, Lmod1, Lmod2, LOC100044087, LOC100044177, LOC100044756, LOC100044862, LOC100045522, LOC100045617, LOC100045887, LOC100046044, LOC100046080, LOC100046136, LOC100046406, LOC100046918, LOC100047012, LOC100047167, LOC100047261, LOC100047353, LOC100047579, LOC100047619, LOC100047794, LOC100047856, LOC100048187, LOC100048445, LOC100048480, LOC100048613, LOC100048622, LOC654842, LOC677528, LOC74457, Lonp1, Lonrf1, Lonrf3, Lor, Lpcat3, Lpin1, Lpin2, Lpin3, Lrit3, Lrp1, Lrpprc, Lrrc16a, Lrrc24, Lrrc28, Lrrc45, Lrrn3, Lsm10, Lsm4, Lsm8, Lsr, Lss, Ltbp4, Ly1, Lyve1, Mad2l1, Maf1, Mak16, Man2a1, Manea, Maoa, Map1lc3b, Map2k6, Map3k15, Map3k4, Map3k5, Map3k6, Mapk12, Mapk1ip1, Mapre2, Marco, Marf1, Mast3, Mat2b, Mbd6, Mcam, Mccc1, Mcl1, Mcoln1, Mcts1, Mdh2, Me1, Med10, Med12, Med20, Med23, Med25, Med30, Med31, Med4, Mef2d, Mertk, Mettl17, Mettl20, Mettl7a1, Mettl7a2, Mff, Mfge8, Mfsd12, Mfsd2a, Mfsd7c, Mgam, Mgp, Mgst3, Mical1, Mid1ip1, Miip, Mknk2, Mkrn2os, Mlx, Mlxip, Mmadhc, Mmp7, Mms19, Morc3, Mpp1, Mp17l, Mp17l2, Mr1, Mrgprb13, Mrpl20, Mrpl23, Mrpl40, Mrpl48, Mrpl53, Mrps17, Mrps27, Mrps28, Mrps35, Mrps36, Msh2, Msh5, Msn, Mst1, Mt1, Mtch2, Mthfd1, Mthfd1l, Mtmr11, Mtmr14, Mtmr7, Mtss1, Mtx2, Mug-ps1, Mum1, Mvd, Mvk, Myadm, Myc, Mycl, Myh10,

Upregulated CR genes (gene symbols known)

Myh14, Myl1, Myl6, Myl9, Myo10, Myo1b, Myo1d, Myo7a, Myo7b, Myom1, N4bp2l1, Naa10, Naa20, Naa38, Nab2, Nadsyn1, Naglu, Naip2, Nampt, Nans, Nbl1, Nbr1, Ncapd3, Nccrp1, Ncoa4, Ncoa6, Ndel1, Ndrg1, Ndrg3, Ndst2, Ndufa11, Ndufa5, Ndufa6, Ndufa8, Ndufaf4, Ndufaf5, Ndufaf6, Ndufb10, Ndufb6, Ndufc1, Ndufc2, Ndufs1, Ndufs2, Ndufs8, Ndufv1, Nedd1, Nedd4, Nek3, Nelfe, Net1, Neurl2, Neurl4, Nfkbia, Nfkbiz, Nfx1, Ngef, Nhirc2, Nhirc3, Nhp2l1, Nid2, Nipa1, Nipal1, Nkiras2, Nlrc5, Nme1, Nme4, Nmnat1, Nmral1, Nmrk1, Nmu, Nnmt, Nog, Nol3, Nop56, Npbwr1, Npc1, Npc2, Npm1, Nqo1, Nqo2, Nr0b2, Nr1d2, Nr4a1, Nrbp2, Nrg4, Nsdhl, Nsmaf, Nsmce2, Nt5c, Nt5c3b, Nucb2, Nudt16, Nudt19, Nudt2, Nudt21, Nudt22, Nudt4, Nup54, Nup62cl, Nupr1, Nx1f1, Oat, Obp2b, Ocln, Ogdh, Ogt, Olfr1175-ps, Olfr467, Omp, Orc6, Orm1, Orm2, Orm3, Osmr, Otud1, Ovgp1, P2ry1, P2ry14, P2ry2, Pafah1b3, Pafah2, Pah, Paics, Pan2, Pank3, Paox, Papl, Papss1, Papss2, Parl, Parp2, Pawr, Paxbp1, Pcbp2, Pcca, Pcgf1, Pcgf5, Pcid2, Pck1, Pcmd2, Pcn1, Pcnxl3, Pcolce, Pcsk4, Pcsk9, Pcyt2, Pcd5, Pde6c, Pde9a, Pdha1, Pdhb, Pdk1, Pdk2, Pdk4, Pdpn, Pdss1, Pdxk, Pdzk1ip1, Peli1, Pemt, Per1, Per2, Per3, Pex7, Pf4, Pfkfb1, Pfkm, Pfn1, Pgam1, Pgd, Pgk1, Pgr15l, Phf11c, Phf19, Phgdh, Phpt1, Pi4ka, Pigf, Pigp, Pigx, Pik3c2g, Pik3cb, Pim3, Pink1, Pitpnc1, Pkd1, Pkd2, Pklr, Pkm, Pla1a, Pla2g16, Plce1, Pld3, Plec, Plekhd1, Plekhd1os, Plekhg2, Plekhj1, Plet1, Plin3, Plin4, Plk3, Pls1, Plxnb1, Plxnb2, Plxnd1, Pm20d1, Pmpcb, Pmvk, Pnp, Pnpla2, Pnpla3, Pnpla5, Podn, Podxl, Pold4, Pole, Polg2, Poli, Polk, Polr2a, Polr2g, Polr2h, Polr3c, Pom121, Por, Postn, Ppa2, Ppap2a, Ppargc1a, Ppargc1b, Ppat, Ppfibp2, Ppib, Ppic, Ppif, Ppip5k2, Ppl, Ppnr, Ppp1r15a, Ppp1r1a, Ppp1r21, Ppp1r3g, Ppp2r3d, Ppp3cc, Ppt2, Pptc7, Praf2, Pramel5, Prdx2, Prdx3, Prelid2, Prg4, Prickle1, Prkab1, Prkag2, Prkcsh, Prod1, Prom1, Pros1, Prosc, Prpf8, Prps1, Prpsap1, Prr3, Prss8, Psat1, Psca, Psmb10, Psmb2, Psmd9, Psme2, Psme2b, Psph, Pstk, Pstkip2, Ptdss2, Ptgds, Ptger1, Ptgis, Ptgr1, Ptgn3, Ptprg, Pus1, Pygl, Qdpr, Qtrt1, Rab20, Rab23, Rab3a, Rab3b, Rab3d, Rab3ip, Rab43, Rab4a, Rab9, Rabep2, Rabggtb, Rad1, Ralgapa2, Ralgds, Ramp2, Ranbp10, Ranbp3l, Ranbp9, Rangrf, Rap1gap, Rapgef5, Raph1, Rarb, Rasa3, Rasgef1b, Rasgrp2, Rasip1, Rassf3, Rassf6, Rbl2, Rbm14, Rbm15, Rbm33, Rbm43, Rbms1, Rbmxl2, Rbp1, Rbpms, Rcbtb1, Rdh11, Rdh14, Rdh5, Rdh9, Reck, Reep4, Rere, Rev1, Rgag1, Rgs16, Rgs4, Rhbdf1, Rhcg, Rhebl1, Rhob, Rhobtb1, Rhod, Rhpn2, Rilp, Ripk4, Rmnd1, Rmnd5a, Rnase1, Rnase4, Rnaseh2b, Rnasek, Rnd1, Rnd2, Rnf14, Rnf145, Rnf167, Rnf19b, Rnf217, Rnf39, Rnf4, Rnf44, Rnh1, Ropn1l, Rorc, Rpa3, Rpap3, Rpe, Rpe65, Rpia, Rpl12, Rpl13, Rpl14, Rpl15, Rpl22l1, Rpl24, Rpl27, Rpl30, Rpl31-ps21, Rpl32, Rpl32, Rpl34, Rpl36, Rpl41, Rpl7a, Rpl9, Rplp0, Rpn2, Rpp38, Rprd1a, Rps12, Rps13, Rps15a, Rps21, Rps27a, Rps4x, Rps5, Rps7, Rras, Rrm1, Rrm2, Rrn3, Rsrc2, Rsrp1, Rtkn, Rundc3b, S100a13, S100a6, Safb, Safb2, Sag, Sall1, Samm50, Sap130, Sbk1, Sbpl, Scamp1, Scamp3, Scarb1, Scarf1, Scarf1a17, Sccpdh, Scd1, Scg3, Scn1b, Scpep1, Sdf2l1, Sdhaf4, Sdhb, Sdhc, Sdhd, Sdpr, Sds, Sdsl, Sec14l1, Sec23ip, Sec24b, Sec31a, Sec61a1, Sec61b, Selenbp1, Selk, Selo, Sema3f, Sema4a, Sep15, Sept11, Sept8, Serpinb6a, Serpine2, Serpinh1, Sertad1, Sesn1, Sesn3, Setd1b, Setd4, Setd6, Setdb2, Sftp, Sfxn1, Sgca, Sgk1, Sh2b3, Sh2d3c, Sh2d4a, Sh3bp1, Sh3d21, Shank3, Siah1a, Siah2, Sidt2, Sigirr, Sik1, Sirt1, Slc12a7, Slc15a2, Slc15a4, Slc16a10, Slc16a5, Slc18b1, Slc19a1, Slc19a2, Slc20a1, Slc22a16, Slc22a17, Slc22a18, Slc22a26, Slc22a27, Slc22a29, Slc23a1, Slc23a2, Slc25a1, Slc25a11, Slc25a12, Slc25a14, Slc25a22, Slc25a29, Slc25a32, Slc25a37, Slc25a51, Slc27a1, Slc29a1, Slc2a4, Slc2a5, Slc2a8, Slc30a1, Slc35b3, Slc35d2, Slc37a4, Slc38a2, Slc38a3, Slc3a1, Slc43a1, Slc44a3, Slc46a3, Slc50a1, Slc51b, Slc5a6, Slc6a6, Slc6a8, Slc7a2, Slc7a4, Slc7a7, Slc8b1, Slc9a1, Slco1a4, Slfn2, Slrp, Slpi, Slurp1, Smad9, Smap2, Smc5, Smdt1, Smg1, Smoc1, Smoc2, Smtn, Snai1, Snai2, Snhg1, Snhg11, Snora21, Snora34, Snora69, Snord15a, Snord16a, Snord1c, Snord22, Snord35a, Snord35b, Snord37, Snord47, Snord61, Snord82, Snord87, Snrk, Snrnp25, Snrnp70, Snrpa1, Snrpd1, Snrpd3, Snrpn, Snx12, Snx13, Snx14, Snx2, Soat2, Soc52, Sod3, Sorbs3, Sox17, Sox18, Sparc, Sparcl1, Spata13, Spata2l,

Upregulated CR genes (gene symbols known)

Spc24, Spcs3, Spdyb, Speer3, Spen1, Spen, Spg21, Spr, Spred1, Sprr1a, Sprtn, Spry4, Sptan1, Sptssa, Sqle, Sqrdl, Sra1, Srbd1, Srebf1, Srp54a, Srrm2, Srsf4, Srsf5, Ssfa2, Ssh2, Sspn, St3gal5, St3gal6, St8sia5, Stab1, Stag2, Stap2, Stard5, Stard7, Stard8, Steap2, Stk11ip, Stk25, Stmn4, Sts, Sulfl2, Sult1a1, Sult1d1, Sult1e1, Sult2a3, Sult3a1, Sumo1, Sumo2, Sun2, Supt3, Supt4a, Supt5, Susd2, Susd6, Suv420h2, Sv2a, Swap70, Swi5, Syde2, Syndig1, Syl1, Tacc2, Taf11, Taf9, Tagln, Taldo1, Tamm41, Tango2, Tapbp, Tars2, Tat, Tbc1d15, Tbc1d19, Tbc1d24, Tbc1d30, Tbc1d8, Tbca, Tbpl1, Tbxa2r, Tcaf2, Tceal8, Tcf19, Tcn2, Tcp11l2, Tctex1d2, Tdg, Tdp1, Tec, Tecr, Tef, Tekt4, Tfdp1, Tfdp2, Tfeb, Tff2, Tgds, Tgfbi, Tgfbr2, Tgif1, Tha1, Theg, Thrsp, Tial1, Tiam1, Tifa, Timm10, Timm17a, Timm8a1, Timm9, Timp3, Tinagl1, Tinf2, Tjp1, Tk1, Tkfc, Tkt, Tlcd1, Tlx2, Tm4sf4, Tm6sf2, Tmc4, Tmc5, Tmc7, Tmed2, Tmem106b, Tmem11, Tmem120b, Tmem126a, Tmem128, Tmem131, Tmem141, Tmem14c, Tmem159, Tmem176b, Tmem218, Tmem237, Tmem245, Tmem25, Tmem41a, Tmem5, Tmem50a, Tmem60, Tmem62, Tmem71, Tmem9, Tmem97, Tmtc2, Tmx2, Tnfrsf11b, Tnfrsf1a, Tnfrsf1b, Tnks1bp1, Tnncl, Tnni2, Tnrc6a, Tns1, Tns2, Tnxb, Tob1, Tom1, Tom1l1, Tomm40, Tox, Tpcn2, Tpd52l1, Tpi1, Tpm4, Tppp, Tpra1, Tpst1, Trak1, Trappc2l, Trdmt1, Trib3, Trim11, Trim28, Trim47, Trim80, Trip10, Trp53bp2, Trrap, Tsc2, Tsc22d3, Tsc22d4, Tspo, Tspyl4, Tsta3, Tstd3, Ttc17, Ttc23, Ttc38, Ttc39a, Tuba1a, Tuba1b, Tuba8, Tubb2a, Tubb4a, Tubb5, Tuft1, Tusc3, Txn2, Txndc16, Txnip, Uaca, Ubac1, Ubb, Ubc, Ube2h, Ube3b, Ube4b, Ubr2, Ubr4, Uchl5, Uck2, Ufc1, Ufsp2, Ugcg, Ugp2, Ugt1a10, Ugt1a2, Ulk1, Unc5b, Unc93a, Upp1, Uqcr10, Uqcrc1, Uqcrcfs1, Uqcrh, Urod, Usf2, Ushbp1, Usmg5, Usp2, Usp28, Uvrag, Vac14, Vamp2, Vamp3, Vangl1, Vasp, Vbp1, Vcam1, Vdac2, Vegfa, Vegfb, Vim, Vkorc1l1, Vldr, Vmn1r1, Vmn1r229, Vmn2r60, Vmn2r70, Vmn2r9, Vpreb3, Vps16, Vps51, Vrk3, Vsig10, Vwce, Vwf, Wbp1l, Wbp2, Wbp2nl, Wbscr27, Wdr1, Wdr55, Wdr81, Wee1, Wfdc15b, Wfdc17, Wipf3, Wnt5b, Wsb1, Xdh, Xlr, Xpa, Xrcc5, Xrcc6bp1, Ybx2, Ypel3, Ypel5, Ythdc1, Zbed4, Zbed5, Zbtb16, Zbtb48, Zc3h6, Zc3h7a, Zcchc14, Zfand4, Zfhx4, Zfp13, Zfp189, Zfp275, Zfp276, Zfp30, Zfp335, Zfp36, Zfp36l2, Zfp467, Zfp579, Zfp607, Zfp608, Zfp612, Zfp639, Zfp647, Zfp653, Zfp703, Zfp750, Zfp930, Zfp933, Zfp97, Zfpm1, Zfyve21, Zhx2, Zkscan17, Zmat1, Zmynd10, Zpr1, Zscan26, Zyx

Upregulated CR genes (only array IDs known)

10338069, 10338074, 10338078, 10338084, 10338085, 10338091, 10338097, 10338118, 10338130, 10338150, 10338163, 10338169, 10338177, 10338178, 10338184, 10338192, 10338205, 10338217, 10338224, 10338225, 10338231, 10338250, 10338255, 10338261, 10338262, 10338271, 10338286, 10338295, 10338300, 10338304, 10338315, 10338319, 10338329, 10338332, 10338339, 10338344, 10338345, 10338348, 10338368, 10338369, 10338384, 10338390, 10338403, 10338412, 10338434, 10338438, 10338448, 10338450, 10338461, 10338473, 10338474, 10338483, 10338490, 10338506, 10338514, 10338515, 10338518, 10338524, 10338535, 10338542, 10338580, 10338584, 10338598, 10338602, 10338604, 10338628, 10338634, 10338637, 10338644, 10338645, 10338646, 10338655, 10338674, 10338678, 10338681, 10338687, 10338699, 10338701, 10338704, 10338709, 10338716, 10338726, 10338742, 10338750, 10338772, 10338773, 10338779, 10338791, 10338799, 10338811, 10338838, 10338841, 10338847, 10338848, 10338852, 10338853, 10338860, 10338865, 10338881, 10338896, 10338905, 10338930, 10338941, 10338946, 10338970, 10338976, 10338989, 10338992, 10339002, 10339013, 10339035, 10339041, 10339063, 10339069, 10339081, 10339086, 10339092, 10339138, 10339145, 10339153, 10339157, 10339161, 10339167, 10339171, 10339174, 10339190, 10339192, 10339241, 10339242, 10339249, 10339270, 10339286, 10339305, 10339323, 10339341, 10339344, 10339348, 10339349, 10339352, 10339363, 10339365, 10339371, 10339394, 10339399, 10339407, 10339457, 10339462, 10339478, 10339479, 10339481, 10339483, 10339490, 10339491, 10339509, 10339510, 10339515, 10339516, 10339539, 10339550, 10339563, 10339581, 10339582,

Upregulated CR genes (only array IDs known)

10339588, 10339593, 10339595, 10339596, 10339604, 10339627, 10339628, 10339631, 10339645, 10339646, 10339648, 10339650, 10339654, 10339655, 10339663, 10339666, 10339670, 10339672, 10339686, 10339687, 10339699, 10339707, 10339722, 10339726, 10339740, 10339756, 10339772, 10339781, 10339783, 10339784, 10339789, 10339797, 10339803, 10339810, 10339813, 10339837, 10339848, 10339856, 10339859, 10339864, 10339874, 10339881, 10339886, 10339897, 10339907, 10339918, 10339931, 10339943, 10339944, 10339945, 10339953, 10339956, 10339971, 10339980, 10339995, 10340000, 10340002, 10340016, 10340019, 10340021, 10340043, 10340056, 10340057, 10340064, 10340069, 10340071, 10340076, 10340080, 10340083, 10340089, 10340110, 10340116, 10340121, 10340127, 10340140, 10340148, 10340163, 10340168, 10340172, 10340186, 10340190, 10340197, 10340206, 10340221, 10340245, 10340258, 10340261, 10340269, 10340270, 10340272, 10340275, 10340277, 10340281, 10340286, 10340301, 10340304, 10340306, 10340326, 10340330, 10340332, 10340338, 10340344, 10340349, 10340389, 10340399, 10340402, 10340407, 10340418, 10340420, 10340444, 10340446, 10340448, 10340459, 10340472, 10340480, 10340497, 10340498, 10340506, 10340511, 10340520, 10340551, 10340563, 10340566, 10340567, 10340574, 10340581, 10340586, 10340588, 10340589, 10340590, 10340607, 10340616, 10340618, 10340627, 10340641, 10340656, 10340670, 10340671, 10340681, 10340695, 10340699, 10340702, 10340712, 10340740, 10340751, 10340756, 10340757, 10340778, 10340791, 10340798, 10340799, 10340801, 10340802, 10340825, 10340834, 10340844, 10340851, 10340875, 10340885, 10340886, 10340898, 10340899, 10340901, 10340909, 10340925, 10340929, 10340932, 10340956, 10340967, 10340973, 10340976, 10340983, 10340991, 10340993, 10341006, 10341017, 10341031, 10341034, 10341039, 10341047, 10341055, 10341062, 10341063, 10341067, 10341068, 10341072, 10341085, 10341093, 10341094, 10341101, 10341106, 10341107, 10341123, 10341131, 10341132, 10341134, 10341140, 10341148, 10341157, 10341160, 10341164, 10341167, 10341174, 10341188, 10341205, 10341216, 10341217, 10341230, 10341232, 10341233, 10341234, 10341245, 10341247, 10341269, 10341296, 10341304, 10341311, 10341348, 10341360, 10341375, 10341383, 10341384, 10341397, 10341428, 10341432, 10341437, 10341444, 10341453, 10341465, 10341473, 10341485, 10341504, 10341518, 10341520, 10341521, 10341530, 10341542, 10341544, 10341549, 10341550, 10341551, 10341565, 10341568, 10341573, 10341580, 10341584, 10341589, 10341612, 10341661, 10341670, 10341679, 10341681, 10341684, 10341691, 10341697, 10341703, 10341710, 10341720, 10341724, 10341732, 10341733, 10341734, 10341744, 10341768, 10341771, 10341777, 10341780, 10341794, 10341808, 10341820, 10341824, 10341852, 10341892, 10341907, 10341914, 10341918, 10341989, 10341992, 10341996, 10341999, 10342005, 10342006, 10342010, 10342017, 10342019, 10342022, 10342031, 10342044, 10342045, 10342070, 10342114, 10342126, 10342140, 10342156, 10342168, 10342178, 10342201, 10342207, 10342211, 10342225, 10342248, 10342257, 10342258, 10342263, 10342276, 10342298, 10342320, 10342340, 10342344, 10342362, 10342363, 10342367, 10342376, 10342380, 10342413, 10342437, 10342446, 10342447, 10342450, 10342451, 10342456, 10342476, 10342479, 10342487, 10342488, 10342496, 10342497, 10342504, 10342508, 10342522, 10342524, 10342570, 10342580, 10342594, 10342606, 10342610, 10342618, 10342621, 10342626, 10342637, 10342649, 10342659, 10342662, 10342663, 10342665, 10342666, 10342668, 10342674, 10342688, 10342692, 10342698, 10342706, 10342708, 10342714, 10342749, 10342754, 10342759, 10342769, 10342789, 10342797, 10342815, 10342835, 10342838, 10342847, 10342851, 10342905, 10342911, 10342916, 10342920, 10342938, 10342939, 10342954, 10342960, 10342963, 10342971, 10342981, 10343008, 10343030, 10343039, 10343044, 10343064, 10343066, 10343072, 10343077, 10343089, 10343090, 10343093, 10343108, 10343124, 10343135, 10343141, 10343148, 10343182, 10343189, 10343197, 10343219,

Upregulated CR genes (only array IDs known)

10343223, 10343237, 10343243, 10343250, 10343251, 10343252, 10343253, 10343272, 10343276, 10343295, 10343296, 10343329, 10343356, 10343358, 10343364, 10343368, 10343377, 10343387, 10343404, 10343407, 10343412, 10343423, 10343427, 10343432, 10343435, 10343438, 10343448, 10343451, 10343452, 10343453, 10343457, 10343462, 10343467, 10343471, 10343474, 10343480, 10343494, 10343496, 10343518, 10343537, 10343549, 10343551, 10343581, 10343595, 10343608, 10343610, 10343612, 10343621, 10343629, 10343633, 10343634, 10343636, 10343637, 10343652, 10343704, 10343705, 10343731, 10343737, 10343754, 10343768, 10343801, 10343802, 10343803, 10343805, 10343833, 10343841, 10343853, 10343863, 10343876, 10343887, 10343890, 10343930, 10343939, 10343942, 10343948, 10343966, 10343974, 10343977, 10343986, 10343990, 10343995, 10344000, 10344008, 10344009, 10344011, 10344012, 10344024, 10344042, 10344074, 10344080, 10344082, 10344085, 10344091, 10344108, 10344113, 10344118, 10344122, 10344131, 10344135, 10344137, 10344152, 10344159, 10344178, 10344221, 10344226, 10344227, 10344230, 10344253, 10344262, 10344263, 10344285, 10344288, 10344302, 10344303, 10344305, 10344315, 10344329, 10344332, 10344353, 10344354, 10344367, 10344384, 10344386, 10344390, 10344395, 10344404, 10344407, 10344412, 10344416, 10344418, 10344429, 10344432, 10344441, 10344443, 10344481, 10344483, 10344488, 10344489, 10344495, 10344497, 10344503, 10344506, 10344512, 10344516, 10344535, 10344536, 10344552, 10344558, 10344584, 10344608, 10344895, 10345436, 10345529, 10346222, 10346874, 10348945, 10351045, 10352556, 10353189, 10355264, 10355454, 10356461, 10356762, 10357298, 10358454, 10359642, 10360145, 10360802, 10361152, 10362033, 10362091, 10362672, 10362902, 10362937, 10362939, 10363005, 10363561, 10364091, 10365230, 10366705, 10367118, 10368220, 10368670, 10368748, 10368883, 10373452, 10374183, 10374313, 10374352, 10374764, 10375242, 10375324, 10378568, 10383887, 10384493, 10386543, 10386548, 10386947, 10388782, 10389162, 10394682, 10395275, 10399461, 10400892, 10401837, 10405969, 10406852, 10407051, 10407346, 10408008, 10410776, 10410919, 10410927, 10412036, 10412828, 10414431, 10415444, 10416107, 10416696, 10416698, 10417700, 10417728, 10420237, 10421917, 10422161, 10422247, 10424347, 10424377, 10426093, 10426435, 10427389, 10427603, 10427742, 10428396, 10428441, 10428918, 10430871, 10431585, 10433177, 10434289, 10434745, 10436200, 10439080, 10440419, 10441359, 10442493, 10445323, 10445875, 10447510, 10447699, 10448230, 10454057, 10454113, 10455595, 10455599, 10456719, 10457667, 10457729, 10457916, 10457927, 10458245, 10459389, 10459768, 10459802, 10461148, 10461150, 10461152, 10461154, 10461156, 10461158, 10461160, 10465043, 10466843, 10468487, 10469127, 10473240, 10474239, 10476399, 10478746, 10479456, 10482432, 10483161, 10483761, 10484355, 10485355, 10489721, 10489870, 10490244, 10490271, 10490815, 10491191, 10491518, 10494007, 10494753, 10495659, 10496032, 10496167, 10496336, 10497327, 10498405, 10498797, 10500527, 10501922, 10502772, 10503212, 10503963, 10505917, 10506498, 10507177, 10507475, 10507851, 10508723, 10510221, 10510546, 10512463, 10512935, 10518568, 10522742, 10526085, 10527425, 10527494, 10527571, 10528165, 10528205, 10528474, 10535902, 10538932, 10539015, 10540072, 10540531, 10541129, 10541480, 10542034, 10543029, 10545152, 10547054, 10547924, 10549534, 10550161, 10550163, 10551282, 10551529, 10552100, 10552106, 10552108, 10554817, 10556113, 10557175, 10560111, 10562639, 10563110, 10564565, 10564663, 10565341, 10565598, 10568534, 10568731, 10570000, 10571310, 10571364, 10571528, 10574432, 10574434, 10574595, 10576657, 10578155, 10578545, 10579047, 10579952, 10580231, 10580984, 10581009, 10581569, 10582983, 10586166, 10586170, 10586172, 10586174, 10586433, 10586863, 10587323, 10587331, 10587339, 10589974, 10593101, 10594186, 10595046, 10595145, 10595148, 10597573, 10601567, 10602176, 10603469, 10603549, 10603803, 10605782,

Upregulated CR genes (only array IDs known)

10607398, 10607429, 100323_at, 100324_g_at, 100380_at, 100578_at, 100669_at, 100694_at, 100727_at, 100732_at, 100734_at, 100767_at, 100984_at, 101013_at, 101024_i_at, 101025_f_at, 101129_at, 101207_at, 101214_f_at, 101227_at, 101324_r_at, 101349_at, 101489_at, 101573_f_at, 101577_at, 101613_at, 101630_f_at, 101663_s_at, 101768_at, 101781_f_at, 101804_at, 101908_s_at, 102004_at, 102091_f_at, 102093_f_at, 102094_f_at, 102098_at, 102102_at, 102109_at, 102126_at, 102804_at, 102818_at, 102847_s_at, 103058_f_at, 103562_f_at, 103728_at, 103969_at, 103978_at, 104074_at, 104228_at, 104456_at, 104693_at, 1415733_a_at, 1415895_at, 1415896_x_at, 1416021_a_at, 1416226_at, 1416415_a_at, 1417366_s_at, 1417505_s_at, 1417580_s_at, 1417823_at, 1417838_at, 1418366_at, 1418536_at, 1418625_s_at, 1419022_a_at, 1419188_s_at, 1419444_at, 1419658_at, 1419836_at, 1419857_at, 1419880_x_at, 1419923_at, 1419958_at, 1420357_s_at, 1420570_x_at, 1421041_s_at, 1421063_s_at, 1421184_a_at, 1421682_a_at, 1422115_a_at, 1422123_s_at, 1422230_s_at, 1422327_s_at, 1422411_s_at, 1422948_s_at, 1423090_x_at, 1423666_s_at, 1423799_at, 1423816_at, 1424245_at, 1424269_a_at, 1424609_a_at, 1424775_at, 1425182_x_at, 1425222_x_at, 1425469_a_at, 1426060_at, 1426061_x_at, 1426359_at, 1426451_at, 1426710_at, 1426906_at, 1427111_s_at, 1427404_x_at, 1427743_at, 1427798_x_at, 1427820_at, 1428014_at, 1428116_a_at, 1428301_at, 1429079_a_at, 1429466_s_at, 1429708_at, 1429993_s_at, 1430019_a_at, 1430172_a_at, 1430375_a_at, 1430641_at, 1431008_at, 1431213_a_at, 1431214_at, 1432589_at, 1433107_at, 1433507_a_at, 1433685_a_at, 1433866_x_at, 1434150_a_at, 1434279_at, 1434362_at, 1434632_at, 1435017_at, 1435630_s_at, 1435716_x_at, 1435819_at, 1435872_at, 1435947_at, 1435948_at, 1436101_at, 1436320_at, 1436666_at, 1436956_at, 1437003_at, 1437584_at, 1437867_at, 1438009_at, 1438177_x_at, 1438276_at, 1438331_at, 1438345_at, 1438879_at, 1438917_x_at, 1438959_x_at, 1439121_at, 1439136_at, 1439404_x_at, 1439468_at, 1439537_at, 1439816_at, 1440016_at, 1440084_at, 1440304_at, 1440427_at, 1440506_at, 1440587_at, 1440710_at, 1440815_x_at, 1441242_at, 1441298_at, 1441324_at, 1441347_at, 1441415_at, 1441430_at, 1441539_at, 1441732_at, 1441793_at, 1442025_a_at, 1442026_at, 1442051_at, 1442182_at, 1442262_at, 1442268_a_at, 1442420_at, 1442514_a_at, 1442725_at, 1442771_at, 1443027_at, 1443095_at, 1443166_at, 1443206_at, 1443322_at, 1443377_at, 1443393_at, 1443430_at, 1443452_at, 1443719_x_at, 1444022_at, 1444113_at, 1444244_at, 1444258_at, 1444366_at, 1444522_at, 1444616_x_at, 1444751_at, 1444765_at, 1444909_at, 1445001_at, 1445031_at, 1445042_at, 1445062_at, 1445492_at, 1445534_at, 1445562_at, 1445591_at, 1445612_at, 1445622_at, 1445640_at, 1445641_at, 1445662_x_at, 1445676_at, 1445688_at, 1445728_at, 1445968_at, 1445987_at, 1445988_at, 1446040_at, 1446081_at, 1446090_at, 1446230_at, 1446293_at, 1446383_at, 1446413_at, 1446461_at, 1446567_at, 1446614_at, 1446615_at, 1446644_at, 1446833_at, 1446903_at, 1446950_at, 1447005_at, 1447092_at, 1447105_at, 1447176_at, 1447188_at, 1447209_at, 1447257_at, 1447322_at, 1447408_at, 1447443_at, 1447482_at, 1447819_x_at, 1447891_at, 1447901_x_at, 1447905_x_at, 1447939_a_at, 1447951_at, 1447999_x_at, 1448715_x_at, 1448775_at, 1448975_s_at, 1449467_at, 1449625_at, 1449745_at, 1449787_at, 1450619_x_at, 1450864_at, 1451077_at, 1451626_x_at, 1451635_at, 1452590_a_at, 1452731_x_at, 1453023_at, 1453299_a_at, 1453473_a_at, 1454686_at, 1455065_x_at, 1455869_at, 1455892_x_at, 1455930_at, 1456032_x_at, 1456393_at, 1456563_at, 1456590_x_at, 1456955_at, 1456973_at, 1457056_at, 1457189_at, 1457486_at, 1457549_at, 1457652_x_at, 1457657_at, 1457830_at, 1457851_at, 1458006_at, 1458007_at, 1458042_at, 1458127_at, 1458266_at, 1458279_at, 1458302_at, 1458382_a_at, 1458542_at, 1458543_at, 1458577_at, 1458751_at, 1458829_at, 1458835_at, 1458947_at, 1458974_at, 1459078_at, 1459196_at, 1459242_at, 1459333_at, 1459339_at, 1459425_at, 1459557_at, 1459626_at, 1459695_at, 1459747_at,

Upregulated CR genes (only array IDs known)

1459775_at, 1459959_at, 1460032_at, 1460072_at, 1460132_at, 1460351_at, 1460466_at, 1460682_s_at, 160081_at, 160098_s_at, 160272_at, 160292_at, 160420_r_at, 160568_at, 160628_at, 160799_at, 161053_at, 161076_at, 161095_i_at, 161127_i_at, 161213_r_at, 161219_r_at, 161252_r_at, 161258_at, 161282_r_at, 161299_r_at, 161300_r_at, 161343_r_at, 161352_r_at, 161363_r_at, 161369_r_at, 161393_at, 161443_r_at, 161462_r_at, 161487_f_at, 161499_f_at, 161503_f_at, 161523_r_at, 161572_r_at, 161578_r_at, 161600_r_at, 161624_r_at, 161657_f_at, 161671_at, 161674_i_at, 161742_r_at, 161782_r_at, 161843_at, 161897_f_at, 161905_r_at, 161944_at, 161972_r_at, 162017_at, 162020_at, 162025_r_at, 162044_f_at, 162060_r_at, 162084_i_at, 162149_i_at, 162210_r_at, 162240_r_at, 162317_r_at, 162318_r_at, 162338_r_at, 162341_r_at, 162386_at, 162396_at, 162434_i_at, 162476_r_at, 162501_at, 92272_at, 92330_r_at, 92408_at, 92577_f_at, 92625_at, 92628_at, 92636_f_at, 92775_at, 92800_i_at, 92855_at, 92900_at, 93022_i_at, 93030_at, 93293_at, 93336_at, 93488_at, 93516_at, 93525_f_at, 93730_at, 93817_at, 93818_g_at, 93883_at, 93904_f_at, 93907_f_at, 93921_at, 93924_f_at, 93987_f_at, 94090_at, 94106_at, 94240_i_at, 94479_at, 94539_f_at, 94685_at, 94716_f_at, 94735_s_at, 94845_at, 95291_r_at, 95523_at, 95643_at, 95766_f_at, 95795_at, 95796_g_at, 95911_at, 95955_at, 96041_at, 96071_at, 96082_at, 96146_at, 96215_f_at, 96255_at, 96290_f_at, 96291_f_at, 96292_r_at, 96300_f_at, 96307_s_at, 96322_at, 96334_f_at, 96339_at, 96358_at, 96485_at, 96530_at, 96606_at, 96607_at, 96611_at, 96718_at, 96741_at, 96906_at, 96912_s_at, 96977_at, 97017_f_at, 97129_at, 97181_f_at, 97252_at, 97293_at, 97401_at, 97536_at, 97598_at, 97647_at, 97751_f_at, 97909_at, 98119_at, 98254_f_at, 98327_at, 98342_at, 98353_at, 98480_s_at, 98564_f_at, 98577_f_at, 98849_at, 99153_at, 99422_at, 99498_at, 99503_at, 99512_at, 99571_at, 99575_at, 99580_s_at, 99593_at, 99868_at, AFFX-18SRNAMur/X00686_3_at, AFFX-BioB-3_st, AFFX-BioB-5_st, AFFX-BioB-M_st, AFFX-BioC-3_st, AFFX-BioC-5_st, AFFX-BioDn-3_st, AFFX-BioDn-5_st, AFFX-CreX-3_st, AFFX-CreX-5_st,

Downregulated CR genes (gene symbols known)

0610009B22Rik, 0610011F06Rik, 0610040J01Rik, 1110004F10Rik, 1110059E24Rik, 1110059G02Rik, 1110059G10Rik, 1110065P20Rik, 1300017J02Rik, 1500012F01Rik, 1600002H07Rik, 1600012H06Rik, 1700007L15Rik, 1700008J07Rik, 1700023H06Rik, 1700066M21Rik, 1810009A15Rik, 1810011O10Rik, 1810013L24Rik, 1810022K09Rik, 1810026B05Rik, 1810026J23Rik, 1810058I24Rik, 2010003K11Rik, 2010309G21Rik, 2010315B03Rik, 2010320M18Rik, 2210016L21Rik, 2210418O10Rik, 2310007O11Rik, 2310009A05Rik, 2310010J17Rik, 2310011J03Rik, 2310030G06Rik, 2310036O22Rik, 2310067E19Rik, 2410004B18Rik, 2410006H16Rik, 2410127L17Rik, 2610001J05Rik, 2610002J23Rik, 2610002M06Rik, 2610034B18Rik, 2610305D13Rik, 2610507B11Rik, 2610524H06Rik, 2700046G09Rik, 2810004N23Rik, 2810416G20Rik, 2810427A07Rik, 2810428I15Rik, 2810459M11Rik, 2810474O19Rik, 2900009J20Rik, 2900026A02Rik, 2900064B18Rik, 2900097C17Rik, 3010026O09Rik, 3110001I22Rik, 3110005L24Rik, 3110073H01Rik, 3110082I17Rik, 3830406C13Rik, 4632427E13Rik, 4633401B06Rik, 4833411C07Rik, 4833417J20Rik, 4833423F13Rik, 4930402H24Rik, 4930453N24Rik, 4930459C07Rik, 4930480G23Rik, 4930522L14Rik, 4930526I15Rik, 4930581F22Rik, 4931406C07Rik, 4933438K21Rik, 4933440N22Rik, 5031434C07Rik, 5033423O07Rik, 5430416N02Rik, 5530402H23Rik, 5730414N17Rik, 5830443J22Rik, 5830469G19Rik, 6030451C04Rik, 6330564D18Rik, 6820402I19Rik, 9030417H13Rik, 9030622O22Rik, 9130016M20Rik, 9130023H24Rik, 9130221J18Rik, 9130401M01Rik, 9130409I23Rik, 9230114K14Rik, 9430016H08Rik, 9930038B18Rik, A130012E19Rik, A130040M12Rik, A1cf, A230050P20Rik, A430035B10Rik, A430102J17Rik, A530040E14Rik, A630089N07Rik, A930004J17Rik, A930033H14Rik, A930035D04Rik, AA408650, AA409587, AA415038, AA792892, Aadat, Aaed1, Aamdc, Aars, Aars2, Aasdhppt, Aatf, Aatk, Abat, Abca1, Abca2, Abcb4, Abcc2, Abcc3, Abcc9, Abcd1, Abcd3, Abce1, Abcf3,

Downregulated CR genes (gene symbols known)

Abcg2, Abhd1, Abhd17b, Abhd2, Abhd3, Abhd6, Abi1, Abi3bp, Ablim3, Acaa1a, Acaa1b, Acad11, Acad12, Acad9, Acadm, Acat1, Acat2, Acbd4, Acbd5, Acnat1, Acnat2, Acot1, Acot12, Acot2, Acot4, Acot8, Acox2, Acp2, Acp5, Acsf2, Acsf1, Acsm3, Acsm5, Actb, Actg1, Actl6a, Actn2, Actr3, Acy3, Ada, Adam10, Adam11, Adam17, Adamdec1, Adap2, Adar, Adck1, Adck3, Adck5, Adgre1, Adgre4, Adgrg3, Adgrv1, Adh4, Adh6-ps1, Adipor2, Adprhl2, Adra1b, Adrm1, Adss, Adtrp, Aen, Aes, Afmid, Aggf1, Agl, Agmo, Agpat2, Agpat3, Agpat5, Agrn, Ahcyl1, Ahcyl2, Ahr, Ahsa1, Ahsa2, Al314180, Al428301, Al464131, Al606181, Al987944, Aif1, Aifm3, Aim1l, Aimp2, Ak3, Akap1, Akap9, Akirin1, Akr1c14, Akr1c20, Akr1d1, Akr1e1, Akt1s1, Alad, Alas2, Alcam, Aldh1a1, Aldh3a2, Aldh5a1, Aldh9a1, Alg11, Alg12, Alg13, Alg2, Alkbh2, Alkbh5, Alkbh7, Alpl, Als2, Alyref, Alyref2, Amacr, Amd1, Amdhd1, Amica1, Ampd2, Amy1, Anapc2, Anapc4, Ang, Ang6, Angptl3, Ankhd1, Ankrd13a, Ankrd28, Ankrd40, Ankrd46, Ankrd49, Ankrd50, Anks4b, Anp32a, Anp32b, Anp32e, Anpep, Aox1, Aox3, Ap2a2, Ap3s1, Apba3, Apcs, Apoa4, Apob, Apobec1, Apof, Apol7a, Apol9b, Apon, Aptx, Aqp4, Aqp8, Aqp9, Ar, Are1, Arf2, Arf4, Arf6, Arfgap1, Argef1, Arglu1, Arhgap35, Arhgap5, Arhgef19, Arl1, Arl4a, Arl5a, Arl5b, Arl6ip1, Arl8b, Arpp19, Arrdc3, Ars, Arsg, Asap3, Asb13, Asb2, Ascc3, Asgr2, Asic5, Asna1, Asnsd1, Aspa, Aspdh, Astx6, Atad2b, Atad3aos, Atf2, Atf5, Atf6, Atf6b, Atg13, Atg5, Atg7, Athl1, Atm, Atoh8, Atp11c, Atp13a3, Atp6v1a, Atp6v1h, Atp7a, Atp8b1, Atp9a, Atpaf1, Atrip, Atxn1, Atxn2, Atxn7l3b, AU014972, AU041133, Avpr1a, AW111846, AW112010, AW209491, AW413774, AW987390, Awat2, Axl, Azin1, B130006D01Rik, B230114P17Rik, B230308N11Rik, B2m, B3galnt1, B3galt1, B3gat3, B4galnt1, B630005N14Rik, Baat, Bace1, Bach1, Bag1, Bag3, Bak1, Baz2a, Bbox1, Bbs5, BC003331, BC004004, BC005537, BC005561, BC005624, BC010981, BC024137, BC025446, BC030308, BC030336, BC038331, BC062258, Bcap29, Bcar1, Bche, Bcl3, Bcl6, Bd1, Bet1, Bglap3, Bhmt, Bhmt2, Bid, Bmp1, Bmp4, Bmp5, Bmp7, Bpgm, Bpnt1, Brd2, Bri3bp, Bsdc1, Bsg, Btaf1, Btc, Btd, Bub3, Bysl, C1d, C1galt1, C1galt1c1, C1qa, C1qb, C1qc, C1s1, C2, C230091D08Rik, C330006A16Rik, C330007P06Rik, C330018D20Rik, C330021F23Rik, C430003N24Rik, C4b, C4bp, C6, C630043F03Rik, C730027H18Rik, C730029A08Rik, C730036E19Rik, C730037M02Rik, C8a, C8b, C8g, C9, Caap1, Cab39, Cables2, Cacybp, Cadm1, Cadm4, Cald1, Caln1, Calr, Camk2n2, Camkk2, Cand1, Canx, Cap1, Capn1, Capn2, Capn7, Caprin1, Capra1, Capza2, Car14, Car3, Car8, Carm1, Casd1, Casp1, Casp4, Casp8, Casp8ap2, Cbl1, Cc2d1a, Ccdc101, Ccdc107, Ccdc115, Ccdc117, Ccdc122, Ccdc126, Ccdc159, Ccdc162, Ccdc181, Ccdc32, Ccdc47, Ccdc58, Ccdc59, Ccdc85b, Ccdc86, Ccdc90b, Ccl5, Ccl9, Ccm2, Ccnc, Ccne2, Ccnf, Ccng1, Ccni, Ccpq1, Ccr5, Ccs, Cd1d1, Cd207, Cd24a, Cd274, Cd2ap, Cd2bp2, Cd300lb, Cd300ld, Cd302, Cd36, Cd3eap, Cd48, Cd52, Cd53, Cd59b, Cd5l, Cd68, Cd74, Cd81, Cd83, Cda, Cdc123, Cdc27, Cdc34, Cdh2, Cdhr5, Cdpt, Cdkn1b, Cdkn2aipnl, Cdo1, Ceacam3, Cebpa, Cebpe, Cebpzos, Cela1, Cela3a, Celf2, Cenpb, Cenpv, Cept1, Ces1d, Ces1e, Ces1f, Ces1g, Ces2a, Ces3a, Cetn3, Cfap36, Cfap97, Cfd, Cfdp1, Cfh, Cfhr1, Cflar, Cfp, Cgref1, Chac2, Chaf1a, Chchd3, Chd4, Chdh, Chic1, Chka, Chkb, Chm, Chmp1a, Chmp2b, Chmp4b, Chmp6, Chn2, Chordc1, Chpf2, Chpt1, Chrna2, Chsy1, Chtop, Chuk, Cidea, Cideb, Cidec, Cited2, Cks1b, Clcn2, Clcn3, Cldn1, Cldn14, Cldn2, Cldn3, Cldnd1, Clec14a, Clec2d, Clec2h, Clec4f, Clec4n, Clec7a, Clint1, Clip2, Cln5, Clock, Clpx, Clstn3, Cltb, Cltc, Cmah, Cmas, Cmbl, Cmip, Cml2, Cml5, Cmpk2, Cmtm4, Cmtm6, Cmtr1, Cnbp, Cnot11, Cnot6, Cnot7, Cnpy3, Coa3, Coa7, Cog8, Coil, Col15a1, Col18a1, Col27a1, Col5a3, Colec10, Commd3, Commd6, Commd7, Commd9, Comt, Copa, Copb1, Copb2, Cops2, Cops6, Copz2, Coq3, Corin, Cot1, Cox15, Cox18, Cox19, Cox6b2, Cp, Cpd, Cpeb4, Cpn1, Cpn2, Cpxo, Cpsf2, Cpt1a, Cpt2, Cptp, Crat, Crbn, Creb3l3, Crebf, Creg1, Creld1, Creld2, Crem, Crip2, Cris1, Crnk1, Crot, Crp, Crygn, Cryz, Csad, Csde1, Cse1l, Csf1r, Csgalnact2, Csnk1a1, Cspp1, Csprs, Csrp3, Cst3, Cstf2, Cstf2t, Ctbp1, Ctcf, Ctcfl, Ctdp1, Ctdsp1, Ctlal2b, Ctnnb1, Ctnnb1, Ctps, Ctss, Ctsz, Cul2, Cul3, Cul4a, Cul4b, Cul5, Cwf19l2, Cxcl1, Cxcl10, Cxcl12, Cxcl9, Cyb561, Cyb561a3, Cyb5a, Cyb5b, Cyb5d2, Cyb5rl, Cyba,

Downregulated CR genes (gene symbols known)

Cybb, Cyp2a12, Cyp2a22, Cyp2c29, Cyp2c37, Cyp2c50, Cyp2c54, Cyp2c70, Cyp2d10, Cyp2d12, Cyp2d13, Cyp2d26, Cyp2d37-ps, Cyp2d40, Cyp2d9, Cyp2f2, Cyp2j5, Cyp2j6, Cyp2u1, Cyp3a13, Cyp3a25, Cyp46a1, Cyp4a12a, Cyp4a12b, Cyp4v3, Cyp7a1, Cyp7b1, Cyp8b1, Cyr61, D10Ertd709e, D130020L05Rik, D17Wsu92e, D1Ertd448e, D2Ertd612e, D430033H22Rik, D4Ertd76e, D5Ertd505e, D5Ertd579e, D5Ertd798e, D5Wsu152e, D630024D03Rik, D630039A03Rik, D830012I24Rik, D830014E11Rik, D930001B02, D930016D06Rik, D930017J03Rik, D9Ertd306e, Daam1, Dag1, Daglb, Dancr, Dapk2, Dars, Daxx, Dbnl, Dbr1, Dcaf10, Dcaf11, Dcaf15, Dcaf5, Dcaf8, Dcakd, Dct, Dcun1d1, Dcun1d4, Dcxr, Ddah1, Ddc, Ddx10, Ddx17, Ddx28, Ddx3x, Ddx3y, Ddx42, Ddx52, Ddx6, Ddx60, Decr1, Decr2, Def8, Dennd4c, Depdc7, Derl1, Derl2, Derl3, Dexi, Dhdh, Dhhs13, Dhrs3, Dhrs4, Dhrs7b, Dhx15, Dhx29, Dhx30, Dhx36, Dhx58, Diablo, Diap2, Dio1, Dip2c, Dis3l, Dkc1, Dkk4, Dleu2, Dmbt1, Dmgdh, Dnaic1, Dnaja1, Dnajb1, Dnajb11, Dnajb14, Dnajb2, Dnajb4, Dnajb9, Dnajc1, Dnajc14, Dnajc2, Dnajc3, Dnajc30, Dnajc5, Dnase1, Dnase2a, Dnase2b, Dnlz, Dnph1, Dnttip2, Dok2, Dolk, Dpagt1, Dpm1, Dpp7, Dpp9, Dpt, Dpy19l1, Dpy19l4, Dpys, Dr1, Dram2, Drg2, Dsg2, Dsp, Dtx3l, Dtx4, Dus2, Dusp23, Dusp3, Dusp6, Dut, Dyncl12, Dynlrb1, E130110O22Rik, E130311K13Rik, E2f1, E2f8, E430010N07Rik, Eaf1, Ears2, Ebag9, Ech1, Eci1, Eci2, Edem3, Ednrb, Eea1, Eef1b2, Eef1e1, Efemp1, Efhd2, Efna1, Efr3a, Egfr, Egln1, Egln3, Ehd1, Ehhadh, Ei24, Eif1a, Eif1ax, Eif1b, Eif2ak2, Eif2b3, Eif2b4, Eif2b5, Eif2s2, Eif2s3x, Eif3e, Eif3k, Eif3m, Eif4a2, Eif4b, Eif4e, Eif4ebp2, Eif4g1, Eif5, Eif5b, El2, Elmod2, Elovl3, Elovl5, Eme2, Eml3, Endod1, Enpp2, Enpp3, Enpp4, Enpp5, Entpd5, Eny2, Epc2, Eph2, Ephx1, Ephx2, Epm2aip1, Eps8l2, Ept1, Ercc6, Ergic1, Ergic2, Erich1, Ermp1, Ero1lb, Esco1, Esd, Esf1, Esrp2, Etfdh, Etnk1, Eva1a, Evi2a, Evi5, Exoc2, Exoc3, Exoc4, Exoc5, Extl1, F8, F9, Fabp2, Fabp7, Fads6, Faf2, Fahd1, Faim, Fam102a, Fam107b, Fam118b, Fam126b, Fam129b, Fam131c, Fam136a, Fam13b, Fam149a, Fam149b, Fam151b, Fam168a, Fam174a, Fam198a, Fam19a5, Fam206a, Fam208b, Fam20a, Fam210a, Fam210b, Fam213b, Fam214a, Fam219b, Fam220a, Fam228b, Fam25c, Fam26f, Fam32a, Fam35a, Fam46d, Fam47e, Fam63a, Fam73b, Fam83f, Fam83h, Fam89a, Fam96a, Fam98a, Fam98b, Farsb, Fas, Fastkd5, Fbxo11, Fbxo30, Fbxw11, Fbxw4, Fcgr1, Fcgr4, Fcgrt, Fcho2, Fchsdl, Fcmr, Fcna, Fen1, Fermt2, Fetub, Fez2, Fgf1, Fgf21, Fgfr1op, Fgfr1op2, Fgl1, Ficd, Fitm1, Fkbp3, Fkbp4, Fkbp8, Fkbp9, Fmn2, Fmr1nb, Fn3krp, Fnbp1l, Fndc3a, Fndc3b, Fnip1, Fntb, Folr2, Foxa2, Foxa3, Foxn2, Foxp1, Foxq1, Fpgs, Fpgt, Fpr2, Fsd1l, Ftsj2, Ftsj3, Fubp1, Fubp3, Fuca1, Fus, Fxn, Fyco1, Fzd7, Fzd8, G0s2, G3bp1, G3bp2, G630090E17Rik, G6pc, Gaa, Gabarapl1, Gabpa, Gabpb1, Gadd45a, Galk1, Galnt1, Galnt2, Gas2, Gas5, Gas6, Gatad2a, Gatc, Gbf1, Gbp10, Gbp2, Gbp2b, Gbp3, Gbp6, Gbp7, Gbp8, Gcat, Gcgr, Gck, Gclc, Gclm, Gcsh, Gdap10, Gdf15, Gemin2, Gfer, Gfm2, Gfod2, Ggact, Ggct, Ggcx, Ggh, Ggnbp2, Ghr, Gid8, Gigyf2, Gimap9, Gjb1, Gjb2, Glcd, Glo1, Glt28d2, Gltpd2, Glyat, Glyctk, Gm10104, Gm10804, Gm11437, Gm11772, Gm11837, Gm12250, Gm12474, Gm13212, Gm15417, Gm16551, Gm17586, Gm19331, Gm19696, Gm19966, Gm20300, Gm20319, Gm2568, Gm26853, Gm2701, Gm2788, Gm4285, Gm4354, Gm4794, Gm4951, Gm4956, Gm5150, Gm5486, Gm5590, Gm561, Gm5617, Gm5802, Gm5817, Gm6484, Gm6658, Gm6682, Gm7241, Gm8979, Gm8995, Gm9706, Gmfb, Gmppb, Gmps, Gna11, Gna12, Gna14, Gnai1, Gngt2, Gnpnat1, Gnrh1, Golga4, Golga7, Golph3, Gorasp1, Gorasp2, Gosr2, Gp49a, Gpatch11, Gpbp1, Gpbp1l, Gpc1, Gpc4, Gpd1, Gpd2, Gphn, Gpn2, Gpr137, Gpr146, Gpr155, Gpr52, Gprin3, Gps1, Gpx4, Grhpr, Grpel2, Gse1, Gsn, Gspt1, Gss, Gsta3, Gstm4, Gstp2, Gt(ROSA)26Sor, Gtf2f1, Gtf2f2, Gtf2h1, Gtf2h2, Gtf2i, Gtf2ird1, Gtf3c4, Gtf3c5, Gtpbp1, Gtpbp10, Gtpbp4, Guk1, Gyk, H19, H1f0, H2-Aa, H2-Ab1, H2afx, H2-DMa, H2-DMb1, H2-Eb1, H2-M2, H2-Oa, H2-Q1, H2-Q4, H2-T24, H6pd, Hacd1, Hacd3, Hacl1, Hadha, Hadhb, Hao1, Hars, Haus8, Hba-a1, Hbb-b2, Hc, Hccs, Hcfc1r1, Hdac1, Hdac11, Hdac2, Hdhd3, Heca, Hectd1, Helz2, Herc6, Hes1, Hes6, Hexim1, Hfe, Hgf, Hgfac, Hgs, Hhip, Hiatl1, Hif1a, Hint2, Hipk2, Hist1h1c, Hist1h2ad, Hist1h2ae, Hist1h2af, Hist1h2ah,

Downregulated CR genes (gene symbols known)

Hist1h2ak, Hist1h2an, Hist1h2ap, Hist1h3a, Hist1h3d, Hist1h3e, Hist1h4i, Hist1h4j, Hist1h4k, Hist1h4n, Hist2h2aa1, Hist2h2ab, Hist2h2ac, Hist2h3c1, Hivep1, Hlf, Hmgb3, Hmgcs2, Hn1, Hn1l, Hnf1b, Hnf4a, Hnf4aos, Hnrnpa3, Hnrnpab, Hnrnpc, Hnrnpd, Hnrnpf, Hnrnph2, Hnrnpk, Hnrnpr, Hnrnpu, Hnrnpul1, Hook1, Hopx, Hp, Hpgd, Hps1, Hps5, Hr, Hras, Hsd17b11, Hsd17b12, Hsd17b13, Hsd17b2, Hsd17b4, Hsd3b2, Hsd3b3, Hsd3b5, Hsd3b7, Hsdl1, Hsdl2, Hsp90aa1, Hsp90ab1, Hsp90b1, Hspa13, Hspa1b, Hspa4, Hspa5, Hspa8, Hspb6, Hspb8, Hspbpb1, Hspbh1, Hyal1, Hyou1, Iba57, Ice1, Id2, Id3, Ide, Ier2, Ifi202b, Ifi27l2a, Ifi35, Ifi44, Ifi47, Ifih1, Ifit1, Ifit2, Ifit3, Ifitm2, Ifngr2, Ift20, Ift22, Igf1it, Igfals, Igfbp2, Igfbp3, Igflr1, Igha, Igkv1-54, Igkv-V3660, Igkv1-133, Igkv4-91, Igsf11, Igsf5, Igtp, ligp1, ligp1b, Ik, Ikbke, Ikbkg, Ikfz5, Il13ra1, Il4i1, Il6st, Illdr2, Immp1l, Immt, Impa1, Impact, Impad1, Inafm2, Inca1, Inhba, Inhbc, Inhbe, Ino80b, Inpp5a, Inpp5f, Insc, Insig1, Insig2, Ints12, Ints2, Ints4, Ints7, Ipo13, Ipo4, Ipo8, Iqce, Iqgap2, Irak1, Irak2, Irf1, Irf5, Irf6, Irf7, Irf9, Irgm1, Irgm2, Irs1, Isg15, Isg20, Isg20l2, Islr, Isoc2b, Itfg1, Itga8, Itgb7, Itih1, Itm2b, Itpa, Itpk1, Jade1, Jagn1, Jak1, Jak3, Jchain, Jmjd8, Josd2, Jtb, Jun, Junb, Jund, Kank2, Kat2a, Kat5, Kat8, Katnbl1, Kbtbd7, Kcnk5, Kcnn1, Kcnt2, Kctd15, Kdm1b, Kdm5d, Kdsr, Keap1, Keg1, Kif5b, Kifc3, Klf10, Klf9, Klhdc7a, Klhl2, Klhl5, Klhl9, Klkb1, Klra2, Klra3, Kpna1, Kpna2, Kpna3, Kpna4, Kpnb1, Krr1, Krt10, Krt8, Kti12, L3mbtl2, Lactb, Lad1, Lag3, Lama3, Lamb3, Lamp2, Lamtor5, Laptm4a, Laptm4b, Larp1, Larp4, Larp4b, Larp7, Lars, Lasp1, Lbp, Lcat, Lcn10, Lcn2, Lcor, Lcp1, Ldha, Ldhd, Ldlr, Leap2, Leprotl1, Lgals1, Lgals3bp, Lgals8, Lgals9, Lhpp, Lifr, Lig4, Lima1, Lin7a, Lin7c, Lipa, Lipc, Lipg, Litaf, Lman2, Lman2l, Lmf1, Lmna, Lmo2, Lnx2, LOC100040592, LOC100044221, LOC100044439, LOC100044576, LOC100045567, LOC100045981, LOC100046232, LOC100047046, LOC100047173, LOC100047200, LOC100047427, LOC100047651, LOC100047674, LOC100047707, LOC100047963, LOC100048105, LOC100048733, LOC382693, LOC641240, LOC669168, LOC670044, LOC75771, Lonp2, Loxl1, Lpar6, Lpgat1, Lppos, Lrat, Lrfn3, Lrg1, Lrif1, Lrp11, Lrp5, Lrrc14b, Lrrc24, Lrrc3, Lrrc58, Lrrc59, Lrrc8d, Lsm12, Lsm2, Lsm5, Lst1, Lt, Ltbr, Ltv1, Luc7l3, Lum, Ly6a, Ly6c2, Ly6d, Ly6e, Ly6g, Ly86, Lypla1, Lyrm2, Lyrm4, Lysmd3, Lyz2, Lztfl1, Mab21l2, Macrod1, Mad2l2, Mafb, Magix, Magoh, Magt1, Mal2, Malsu1, Mamdc2, Man1a, Man2b2, Manf, Map1lc3a, Map1s, Map2k4, Map3k11, Map3k2, Map3k7, Mapk1, Mapk15, Mapk1ip1l, Mapkapk5, Mapre1, March6, March8, Mars, Mat2a, Matr3, Mbd2, Mbd3, Mbl1, Mcat, Mcfd2, McM10, McM2, Mcs2, Mdfic, Med19, Med21, Med28, Megf9, Meig1, Meiob, Metap2, Mettl1, Mettl10, Mettl23, Mettl3, Mettl5, Mettl8, Mfsd3, Mgll, Mgmt, Mgst1, Mia2, Mia3, Micu1, Midn, Mief2, Mier3, Mif4gd, Mir15a, Mir22hg, Mis12, Mlec, Mlf2, Mlh3, Mlxip1, Mmd, Mme, Mmgt1, Mmp15, Mmp19, Mn1, Mob4, Mocos, Mocs2, Mogat1, Mon1a, Morf4l2, Morn2, Mospd3, Moxd1, Mpc1, Mpdz, Mpeg1, Mphosph10, Mplkip, Mpp6, Mpst, Mpv17, Mpzl2, Mreg, Mrpl10, Mrpl36, Mrpl50, Mrpl57, Mrps18a, Mrps22, Mrs2, Ms4a4d, Ms4a6c, Msantd4, Msh6, Msl1, Msl2, Msmo1, Msrb1, Mt2, Mtap, Mtch1, Mterf2, Mtfr1, Mtif2, Mtif3, Mtm1, Mtmr4, Mtmr9, mt-Nd6, Mtnr1a, Mtpn, Mtrf1l, Mtpp, Mtus1, Mug1, Mug2, Mup10, Mup2, Mup20, Mup21, Mup3, Mup4, Mup5, Mup6, Mus81, Mustn1, Mx1, Mx2, Mxd4, Mydgf, Myh1, Myh9, Myl2, Myo5b, N6amt2, Naa15, Naa16, Naa50, Naa60, Nacc2, Nadk2, Nae1, Nap1l1, Nat6, Nat8, Nat9, Nbeal1, Nbn, Ncald, Ncbp1, Ncbp2, Ncf4, Nckap1l, Ncl, Ncmap, Ncor1, Ndnl2, Ndrg2, Ndrl1, Ndrl2, Ndrl3, Neat1, Neb, Necab1, Nedd4l, Nek4, Nek6, Nek7, Nelfcd, Nenf, Nespas, Nfatc3, Nfe2l1, Nfe2l2, Nfia, Nfib, Nfic, Nfil3, Nfix, Ngfr, Nhirc1, Nhp2, Nicn1, Nif3l1, Nifk, Ninj1, Nip7, Nipa2, Nipsnap3b, Nit2, Nkiras1, Nlk, Nln, Nlrp12, Nlrp6, Nlrx1, Nmd3, Nmi, Nol8, Nolc1, Nop58, Nop9, Notum, Nox4, Noxo1, Npepl1, Nploc4, Npm3, Npr2, Nr1d1, Nr1h3, Nr1i3, Nr2c2ap, Nr2f2, Nr2f6, Nr3c1, Nr5a2, Nras, Nrbf2, Nrbp1, Nrep, Nrip1, Nrp1, Nsd1, Nsf1c, Nsmf, Nt5dc1, Nt5dc2, Ntan1, Nthl1, Ntmt1, Ntn4, Ntpcr, Nuak2, Nub1, Nubp2, Nudcd2, Nudt1, Nudt12, Nudt16l1, Nudt18, Nudt7, Nudt8, Nufip2, Nup153, Nup35, Nup93, Nus1, Nxpe2, Nxt2, Oard1, Oasl1, Oasl2, Oaz2, Obp2a, Ocel1, Ociad1, Odc1, Odf3b, Ogfr, Olfr1126, Olfr1381, Olfr206, Olfr30, Olfr378, Olfr538, Olfr60, Olfr607, Olfr702, Olfr786, Olfr974, Olig1, Oma1, Omd,

Downregulated CR genes (gene symbols known)

Onecut1, Onecut2, Opa3, Oplah, Oraov1, Orc3, Orc4, Ormdl1, Osbp, Osbpl3, Osbpl5, Osbpl8, Osgin1, Ostc, Otud6b, Oulin, Ovca2, Oxr1, Oxsr1, Pa2g4, Pabpc1, Pafah1b1, Paip1, Paip2, Palld, Papd5, Papola, Paqr3, Paqr7, Paqr9, Pard6b, Parg, Parn, Parp12, Parp16, Parp3, Parp9, Parva, Pbdc1, Pbld2, Pbld2, Pbrm1, Pcbd2, Pcbp1, Pcbp4, Pcdh1, Pcdhga8, Pcgf6, Pcm1, Pcnp, Pcp4l1, Pcsk6, Pctp, Pcyt1a, Pdcd10, Pdcd2, Pdcd4, Pdcl, Pde12, Pde6d, Pdia3, Pdia4, Pdia6, Pdik1l, Pdilt, Pdlim5, Pdpr, Pdzk1, Pelo, Perp, Pes1, Pex11a, Pex12, Pex13, Pex14, Pex19, Pex2, Pex26, Pfdn1, Pfn2, Pgam5, Pgap1, Pgap2, Pggt1b, Pglyrp1, Pglyrp2, Pgm3, Pgpep1, Phf20l1, Phf23, Phkb, Phlda1, Phldb2, Phospho1, Pias2, Piccalm, Pid1, Pigk, Pigl, Pigq, Pigr, Pigt, Pigyl, Pik3ap1, Pik3r1, Pik3r2, Pik3r3, Pilrb1, Pim1, Pisd, Pitpna, Pitpn, Pja2, Pkdcc, Pknox1, Pkp4, Pla2g12a, Pla2g12b, Pla2g15, Pla2g4a, Pla2g6, Plaa, Plac8, Plcg2, Plcx2d, Pld4, Plekhb1, Plekhf1, Plin2, Plin5, Pllp, Plrg1, Pls3, Plscr2, Pltp, Pmm1, Pnkd, Pnkp, Pnlcd1, Pnn, Pno1, Pnpla7, Pnpo, Pnrc2, Poc1b, Pofut1, Polb, Pold2, Polr1e, Polr2i, Polr2l, Polr3k, Pon1, Ppan, Ppap2b, Ppapdc1b, Ppara, Ppcs, Ppdf, Pphln1, Ppid, Ppig, Ppm1m, Ppp1ca, Ppp1cb, Ppp1r11, Ppp1r12a, Ppp1r14a, Ppp1r37, Ppp1r3b, Ppp1r3c, Ppp1r9a, Ppp2ca, Ppp2r1b, Ppp2r3a, Ppp2r4, Ppp2r5a, Ppp2r5c, Ppp2r5e, Ppp3ca, Ppp4c, Ppp4r1, Ppp4r2, Ppp6c, Pprc1, Pqb1, Pramef8, Prdx6, Prelp, Prepl, Prkaa2, Prkaca, Prkar2a, Prkd3, Prkrir, Prlr, Prmt1, Proca1, Prorsd1, Prox1, Proz, Prpf38b, Prpf40a, Prr13, Prr16, Prr33, Prrg2, Prss36, Prtn3, Psap, Psen2, Psma3, Psma4, Psma5, Psmb8, Psmb9, Psmc2, Psmc4, Psmc5, Psmc6, Psmd1, Psmd11, Psmd12, Psmd14, Psmd4, Psmd6, Psme3, Pspc1, Ptbp3, Ptdss1, Pten, Pter, Ptk2b, Ptma, Ptms, Ptp4a2, Ptpn1, Ptpn12, Ptpn6, Ptprb, Ptprc, Ptprf, Ptrh2, Pttg1ip, Pum1, Pura, Purb, Pus7, Pvrl2, Pvrl3, Pwwp2b, Pxmp4, Qk, Qpct, Qrich1, Qrs1, Qsox1, R3hdm2, Rab1, Rab10, Rab11a, Rab14, Rab18, Rab21, Rab24, Rab29, Rab30, Rab33b, Rab5a, Rab6a, Rab7, Rabepk, Rabl3, Rad17, Rad21, Rad23b, Rad51b, Raet1d, Rai14, Rala, Ralb, Ran, Ranbp1, Ranbp2, Ranbp6, Rangap1, Rap1a, Rap1b, Rapgef1, Rarres1, Rars, Rasa1, Rassf8, Rbbp4, Rbbp9, Rbck1, Rbm19, Rbm25, Rbm39, Rbm45, Rbm7, Rbm8a, Rbmxl1, Rcan1, Rcbtb2, Rcc2, Rchy1, Rcor1, Rdh16, Rdh7, Rdx, Recql5, Reep3, Reep6, Reg1, Rela, Rel1, Retsat, Rexo2, Rfc1, Rfesd, Rfk, Rfwd2, Rfxank, Rgp1, Rgs12, Rgs3, Rgs5, Rhbdd3, Rhbdl1, Rhbg, Rheb, Rhog, Rhou, RIBC1, Rif1, Rin3, RioK2, Rnase2a, Rnf11, Rnf121, Rnf123, Rnf125, Rnf135, Rnf138, Rnf152, Rnf168, Rnf186, Rnf215, Rnf31, Rnf43, Rnf5, Rnf6, Rnf7, Rnft1, Rngtt, Rnpsc3, Rnpep, Rock1, Rock2, Rom1, Rpl22, Rpl35a, Rpl7I1, Rpp14, Rpp40, Rpr1, Rpr12, Rps25, Rps27, Rps4l, Rraga, Rras2, Rrbp1, Rrp12, Rrp36, Rrp8, Rrs1, Rsad2, Rsl24d1, Rspry1, Rtfdc1, Rtn4, Rtn4ip1, Rtn4rl1, Rtp3, Rtp4, Rufy1, Rwd4a, Rxra, S100a10, S100a8, S100a9, S1pr1, S1pr5, Saa1, Saa2, Saa3, Saa4, Samd9l, Sap30, Sap30l, Sar1a, Saraf, Saraf, Sat1, Sc5d, Scaf11, Scamp2, Scamp5, Scara5, Scly, Scnm1, Sco1, Sco2, Scoc, Scp2, Scyl3, Sdc1, Sdc2, Sdc4, Sdcbp, Sdhaf1, Sdr42e1, Sdr9c7, Sec14l2, Sec16b, Sec22b, Sec23b, Sec24d, Sec62, Sec63, Secisbp2, Secisbp2l, Selenbp2, Selm, Selplg, Sema4g, Sema5b, Senp5, Senp7, Sephs1, Sept7, Sept9, Sepw1, Serbp1, Serinc1, Serinc2, Serinc5, Serpina10, Serpina11, Serpina12, Serpina1e, Serpina3a, Serpina3f, Serpina3g, Serpina3k, Serpina3n, Serpina4-ps1, Serpina7, Serpina9, Serpinb1a, Serpinb9, Serpine1, Serpinf2, Serping1, Setbp1, Sf3b1, Sf3b6, Sfr1, Sgk2, Sgms2, Sgol2b, Sgpl1, Sgpp1, Sh3bgrl, Sh3bgrl2, Sh3bp2, Shc1, Shisa5, Shoc2, Shtn1, Siae, Sigmar1, Sipa1l1, Skil, Skiv2l, Skp1a, Slc10a2, Slc10a3, Slc10a5, Slc11a1, Slc12a2, Slc13a2, Slc13a3, Slc15a3, Slc16a1, Slc16a12, Slc16a13, Slc16a2, Slc16a6, Slc16a7, Slc17a1, Slc17a2, Slc17a3, Slc17a4, Slc17a5, Slc17a9, Slc1a2, Slc1a4, Slc20a2, Slc22a1, Slc22a23, Slc22a30, Slc22a5, Slc22a7, Slc25a10, Slc25a13, Slc25a16, Slc25a20, Slc25a23, Slc25a25, Slc25a30, Slc25a33, Slc25a45, Slc26a1, Slc27a2, Slc27a4, Slc28a1, Slc30a10, Slc30a5, Slc30a9, Slc33a1, Slc35a1, Slc35a3, Slc35b4, Slc35d1, Slc35e3, Slc39a1, Slc39a14, Slc39a2, Slc39a4, Slc39a7, Slc39a8, Slc40a1, Slc41a2, Slc45a3, Slc4a4, Slc4a7, Slc6a13, Slc6a16, Slc6a9, Slco1a1, Slco1b2, Smad3, Smad6, Smagp, Smarca2, Smarca4, Smarca5, Smarcal1, Smarcc1, Smarcc2, Smc4, Smco4,

Downregulated CR genes (gene symbols known)

Smek2, Smg5, Smg7, Smim10l1, Smim14, Smim15, Smim4, Smim8, Smn1, Smo, Smurf2, Smyd2, Smyd4, Snap23, Snap47, Snapin, Snca, Snhg12, Snhg4, Snhg6, Snhg8, Snord118, Snrnp27, Snrnp35, Snrnp40, Snrbp2, Snrpe, Sntb1, Snx17, Snx19, Snx24, Snx27, Snx5, Soc3, Sorbs1, Sord, Sos2, Sox6os, Sp1, Sp3, Sp5, Spast, Spata2, Spcs2, Spg20, Spin1, Spon2, Spop, Sppl2a, Spryd7, Spsb3, Spsb4, Sqstm1, Srd5a1, Srgn, Srp68, Srp9, Srprb, Srr, Srsf1, Srsf10, Srsf2, Srsf3, Srsf6, Srsf7, Srxn1, Ss18, Ssb, Ssbp3, Ssr1, St3gal1, St3gal3, St3gal4, St5, St6gal1, St6galnac6, Stag1, Stap1, Stard10, Stard13, Stat1, Stat2, Stat5a, Stat6, Stbd1, Stip1, Stk16, Stk3, Stk4, Stom, Stra6l, Stradb, Strn3, Stt3a, Stt3b, Stx4a, Stxbp6, Styx, Sub1, Sucnr1, Sugct, Sult1c2, Sult2a8, Sult5a1, Sumf1, Sumf2, Sumo3, Suox, Susd4, Suz12, Swsap1, Syap1, Syce2, Syncp3, Syncrip, Synrg, Synv1, Szrd1, Tab2, Tada2b, Taf10, Taf1a, Taf6, Taf6l, Tap1, Tap2, Tapt1, Tardbp, Tars, Tatdn2, Tax1bp1, Tbc1d8b, Tbcb, Tbcel, Tbk1, Tbl1x, Tbl1xr1, Tbrg1, Tcaim, Tcea1, Tceb1, Tcerg1, Tcf4, Tefm, Tenm3, Tes3-ps, Tesk1, Tex12, Tex2, Tex264, Tex30, Tfam, Tfec, Tff3, Tfpi2, Tfr2, Tgfbr1, Tgm2, Tgs1, Thap11, Them7, Thoc2, Thpo, Thrap3, Thrb, Thtpa, Thy1, Tiam2, Tigd2, Timd2, Timd4, Timm8b, Timmdc1, Tipin, Tirap, Tjp3, Tk2, Tlcd2, Tle1, Tle2, Tle6, Tlk1, Tlr12, Tlr5, Tm2d1, Tm7sf2, Tm9sf2, Tm9sf3, Tmbim1, Tmcc1, Tmco1, Tmed7, Tmed9, Tmem101, Tmem102, Tmem120a, Tmem123, Tmem125, Tmem127, Tmem134, Tmem14a, Tmem150a, Tmem167, Tmem176a, Tmem177, Tmem180, Tmem184a, Tmem184b, Tmem19, Tmem230, Tmem243, Tmem248, Tmem254a, Tmem263, Tmem30a, Tmem30b, Tmem33, Tmem37, Tmem42, Tmem56, Tmem64, Tmem86a, Tmem98, Tmie, Tmprss6, Tmsb4x, Tmx1, Tnfaip2, Tnfaip8l1, Tnnt1, Tnnt3, Tnpo3, Tomm20, Tomm70a, Top1mt, Top2b, Tor1aip1, Tor1aip2, Tor1b, Tor2a, Tor3a, Tpgs1, Tpgs2, Tpmt, Tpp2, Tprgl, Traf3, Trappc10, Trappc13, Treh, Treml4, Trex1, Trhde, Triap1, Trim12c, Trim13, Trim2, Trim21, Trim30a, Trim30d, Trim32, Trim39, Trip6, Trmt10a, Trmt11, Trmt6, Trp53inp1, Trp53inp2, Trp53rka, Trpm7, Trpv2, Trub2, Tsacc, Tsc22d1, Tsku, Tspan12, Tspan14, Tspan31, Tspan33, Tspan4, Tspyl1, Tsr2, Tssc4, Tstd2, Ttc13, Ttc14, Ttc27, Ttc30b, Ttc32, Ttc39c, Ttc7b, Tti1, Ttpa, Ttyh2, Tuba1c, Tuba4a, Tubb4b, Tubg1, Tug1, Tusc2, Tvp23b, Twf1, Twf2, Twistnb, Twsg1, Txndc17, Txndc5, Txndc9, Txnl1, Txnl4b, Txndl1, Tyrobp, Tysnd1, Tyw5, U2af2, Uba3, Uba5, Uba6, Ubd, Ube2a, Ube2d1, Ube2d2a, Ube2d3, Ube2e1, Ube2e2, Ube2e3, Ube2f, Ube2g1, Ube2g2, Ube2l3, Ube2l6, Ube2o, Ube2q2, Ube2r2, Ube2v2, Ube3a, Ubl4, Ubl7, Ublcp1, Ubp1, Ubr3, Ubxn1, Ubxn2a, Ubxn4, Uck1, Ufd1l, Ufl1, Ugdh, Ugt2b1, Ugt2b34, Ugt2b35, Ugt2b37, Ugt2b5, Ugt3a1, Ugt3a2, Uhmkl1, Ulk2, Unc119, Ung, Unkl, Upf1, Upf2, Uqcc2, Uros, Uso1, Usp1, Usp14, Usp18, Usp33, Usp34, Usp39, Usp7, Usp9x, Uspl1, Utp18, Utp6, Uxt, Vamp7, Vapb, Vaultrc5, Vcl, Vcp, Vcpip1, Vma21, Vmac, Vmn1r174, Vmn2r110, Vmn2r121, Vmn2r37, Vmn2r61, Vmn2r96, Vmo1, Vnn1, Vnn3, Vprbp, Vps35, Vps37a, Vps37b, Vps41, Vps72, Vti1b, Vwa8, Wac, Wbp1, Wbp5, Wbscr16, Wdr12, Wdr18, Wdr43, Wdr45b, Wdr74, Wdr77, Wdr92, Wdtc1, Wfdc12, Wfdc2, Wfdc21, Wnk1, Wnt2, Wrap73, Wrnip1, Wtap, Wwc1, Wwp1, Xbp1, Xiap, Xist, Xkr9, Xlr4a, Xpnpep3, Xpo1, Xpo6, Xpot, Xrn2, Xylb, Xylt2, Yae1d1, Yaf2, Yap1, Yipf4, Yipf5, Yipf6, Ykt6, Yme1l1, Ythdc2, Ythdf1, Ythdf2, Ythdf3, Ywhae, Ywhag, Ywhaz, Zbed6, Zbp1, Zbtb11os1, Zbtb18, Zbtb20, Zbtb22, Zbtb33, Zbtb41, Zbtb6, Zc3h12d, Zc3h18, Zc3hav1, Zcchc10, Zcchc2, Zcchc6, Zdhhc16, Zfand1, Zfand2a, Zfhx3, Zfp1, Zfp110, Zfp114, Zfp131, Zfp141, Zfp146, Zfp148, Zfp213, Zfp229, Zfp260, Zfp263, Zfp281, Zfp292, Zfp322a, Zfp324, Zfp326, Zfp362, Zfp367, Zfp36l1, Zfp386, Zfp397, Zfp41, Zfp422, Zfp442, Zfp445, Zfp456, Zfp472, Zfp51, Zfp518a, Zfp551, Zfp563, Zfp58, Zfp617, Zfp62, Zfp655, Zfp664, Zfp672, Zfp687, Zfp729b, Zfp74, Zfp748, Zfp758, Zfp759, Zfp770, Zfp790, Zfp799, Zfp809, Zfp825, Zfp866, Zfp871, Zfp931, Zfp932, Zfp935, Zfp943, Zfp946, Zfp955b, Zfp958, Zfp1, Zfr, Zfyve16, Zfyve19, Zfyve26, Zkscan7, Zmynd8, Znfx1, Znhit1, Znhit2, Znhit3, Znhit6, Znrd1, Znrf2, Zranb2, Zscan29, Zswim1, Zswim7, Zwint

Downregulated CR genes (only array IDs known)

10338065, 10338073, 10338087, 10338114, 10338146, 10338149, 10338153, 10338160, 10338168, 10338179, 10338185, 10338186, 10338190, 10338200, 10338201, 10338222, 10338243, 10338246, 10338256, 10338263, 10338264, 10338274, 10338289, 10338297, 10338307, 10338318, 10338321, 10338334, 10338343, 10338365, 10338377, 10338386, 10338394, 10338406, 10338407, 10338409, 10338417, 10338466, 10338497, 10338502, 10338526, 10338527, 10338556, 10338579, 10338586, 10338591, 10338599, 10338603, 10338607, 10338613, 10338617, 10338620, 10338625, 10338640, 10338647, 10338653, 10338661, 10338670, 10338694, 10338696, 10338708, 10338717, 10338732, 10338749, 10338755, 10338770, 10338774, 10338776, 10338784, 10338824, 10338829, 10338877, 10338879, 10338884, 10338891, 10338894, 10338906, 10338910, 10338914, 10338917, 10338931, 10338932, 10338942, 10338945, 10338951, 10338952, 10338968, 10338969, 10339007, 10339008, 10339011, 10339026, 10339032, 10339034, 10339037, 10339042, 10339052, 10339058, 10339060, 10339065, 10339078, 10339083, 10339085, 10339087, 10339096, 10339098, 10339107, 10339120, 10339129, 10339133, 10339155, 10339162, 10339206, 10339212, 10339214, 10339260, 10339274, 10339282, 10339284, 10339288, 10339289, 10339310, 10339312, 10339321, 10339335, 10339338, 10339340, 10339356, 10339368, 10339369, 10339395, 10339396, 10339424, 10339429, 10339450, 10339452, 10339453, 10339482, 10339494, 10339544, 10339559, 10339607, 10339610, 10339611, 10339614, 10339635, 10339676, 10339696, 10339703, 10339706, 10339729, 10339733, 10339743, 10339750, 10339771, 10339777, 10339795, 10339801, 10339808, 10339827, 10339834, 10339842, 10339851, 10339883, 10339891, 10339900, 10339902, 10339922, 10339924, 10339930, 10339939, 10339940, 10339946, 10339950, 10339951, 10339978, 10339991, 10339992, 10339994, 10340048, 10340065, 10340066, 10340075, 10340090, 10340091, 10340098, 10340099, 10340122, 10340130, 10340141, 10340142, 10340147, 10340153, 10340169, 10340173, 10340218, 10340220, 10340233, 10340241, 10340252, 10340255, 10340263, 10340282, 10340289, 10340302, 10340339, 10340351, 10340353, 10340354, 10340383, 10340390, 10340396, 10340401, 10340413, 10340426, 10340432, 10340447, 10340467, 10340474, 10340475, 10340478, 10340496, 10340508, 10340517, 10340519, 10340526, 10340527, 10340540, 10340542, 10340547, 10340550, 10340560, 10340569, 10340591, 10340623, 10340644, 10340655, 10340659, 10340667, 10340674, 10340682, 10340691, 10340701, 10340708, 10340713, 10340722, 10340731, 10340739, 10340749, 10340759, 10340763, 10340776, 10340787, 10340813, 10340819, 10340833, 10340835, 10340837, 10340857, 10340868, 10340878, 10340884, 10340887, 10340888, 10340892, 10340894, 10340895, 10340933, 10340944, 10340954, 10340966, 10340968, 10340987, 10340989, 10340999, 10341008, 10341030, 10341052, 10341054, 10341071, 10341080, 10341084, 10341088, 10341089, 10341097, 10341109, 10341118, 10341122, 10341146, 10341151, 10341154, 10341178, 10341179, 10341182, 10341183, 10341184, 10341186, 10341191, 10341192, 10341202, 10341214, 10341228, 10341237, 10341242, 10341250, 10341256, 10341273, 10341276, 10341280, 10341302, 10341305, 10341306, 10341314, 10341344, 10341354, 10341359, 10341361, 10341363, 10341368, 10341371, 10341394, 10341412, 10341413, 10341429, 10341435, 10341451, 10341454, 10341467, 10341472, 10341479, 10341493, 10341498, 10341511, 10341545, 10341566, 10341578, 10341586, 10341590, 10341595, 10341643, 10341650, 10341651, 10341660, 10341669, 10341690, 10341702, 10341704, 10341711, 10341712, 10341716, 10341748, 10341761, 10341767, 10341781, 10341785, 10341796, 10341802, 10341815, 10341816, 10341825, 10341830, 10341846, 10341854, 10341857, 10341865, 10341871, 10341886, 10341913, 10341921, 10341923, 10341924, 10341928, 10341932, 10341938, 10341950, 10341959, 10341960, 10341964, 10341975, 10341978, 10342004, 10342016, 10342021, 10342024, 10342033, 10342037, 10342043, 10342050, 10342051, 10342052, 10342054, 10342055, 10342081, 10342101, 10342104, 10342122,

Downregulated CR genes (only array IDs known)

10342161, 10342166, 10342174, 10342179, 10342200, 10342206, 10342221, 10342242, 10342260, 10342269, 10342292, 10342297, 10342315, 10342317, 10342330, 10342364, 10342379, 10342407, 10342427, 10342434, 10342440, 10342449, 10342463, 10342464, 10342470, 10342490, 10342494, 10342514, 10342528, 10342535, 10342536, 10342540, 10342558, 10342559, 10342560, 10342561, 10342583, 10342590, 10342596, 10342619, 10342634, 10342644, 10342652, 10342701, 10342711, 10342720, 10342727, 10342745, 10342748, 10342755, 10342763, 10342764, 10342765, 10342779, 10342782, 10342798, 10342807, 10342826, 10342846, 10342849, 10342853, 10342881, 10342885, 10342889, 10342898, 10342900, 10342910, 10342919, 10342925, 10342928, 10342933, 10342937, 10342943, 10342957, 10342958, 10342977, 10342978, 10342987, 10342989, 10342997, 10342999, 10343007, 10343023, 10343062, 10343074, 10343080, 10343085, 10343088, 10343105, 10343117, 10343118, 10343127, 10343128, 10343131, 10343137, 10343138, 10343139, 10343142, 10343159, 10343174, 10343178, 10343184, 10343188, 10343193, 10343196, 10343205, 10343221, 10343224, 10343233, 10343240, 10343277, 10343306, 10343331, 10343335, 10343382, 10343384, 10343385, 10343392, 10343401, 10343415, 10343425, 10343443, 10343450, 10343455, 10343473, 10343475, 10343490, 10343508, 10343512, 10343514, 10343515, 10343516, 10343536, 10343553, 10343558, 10343565, 10343574, 10343579, 10343586, 10343609, 10343625, 10343630, 10343641, 10343654, 10343666, 10343668, 10343674, 10343682, 10343687, 10343695, 10343717, 10343719, 10343721, 10343732, 10343742, 10343758, 10343762, 10343766, 10343789, 10343798, 10343800, 10343814, 10343817, 10343822, 10343823, 10343845, 10343852, 10343871, 10343877, 10343882, 10343895, 10343896, 10343900, 10343901, 10343919, 10343928, 10343936, 10343943, 10343963, 10343969, 10343979, 10343980, 10343987, 10343997, 10344002, 10344018, 10344021, 10344048, 10344049, 10344063, 10344064, 10344068, 10344070, 10344075, 10344077, 10344081, 10344096, 10344097, 10344098, 10344114, 10344123, 10344156, 10344157, 10344164, 10344173, 10344182, 10344190, 10344191, 10344197, 10344210, 10344220, 10344223, 10344232, 10344236, 10344250, 10344264, 10344267, 10344273, 10344275, 10344278, 10344292, 10344296, 10344297, 10344319, 10344344, 10344346, 10344372, 10344413, 10344420, 10344431, 10344463, 10344469, 10344475, 10344478, 10344508, 10344513, 10344522, 10344532, 10344533, 10344540, 10344543, 10344557, 10344566, 10344580, 10344592, 10344595, 10344609, 10344610, 10345203, 10347915, 10347925, 10351504, 10353303, 10356262, 10356274, 10358432, 10360391, 10362422, 10362424, 10362426, 10365830, 10375121, 10375123, 10375515, 10376326, 10383168, 10383192, 10383198, 10383200, 10383206, 10383208, 10385513, 10385518, 10385526, 10398039, 10399354, 10400189, 10403015, 10403018, 10403021, 10403028, 10403034, 10403038, 10403043, 10403048, 10403054, 10403060, 10403063, 10403079, 10403957, 10404053, 10405916, 10408092, 10408557, 10408647, 10409202, 10410386, 10410388, 10411622, 10412657, 10414527, 10415081, 10418185, 10418198, 10418210, 10423002, 10424383, 10425049, 10426650, 10427624, 10428940, 10429568, 10430174, 10432398, 10438415, 10450694, 10451287, 10453736, 10455957, 10455970, 10457020, 10458547, 10458890, 10460253, 10461055, 10461636, 10464445, 10467390, 10468877, 10468885, 10476939, 10477004, 10482762, 10485357, 10485580, 10489343, 10491820, 10493494, 10496592, 10500327, 10504755, 10505237, 10505240, 10505246, 10505249, 10505252, 10505254, 10505258, 10505261, 10505266, 10505268, 10513420, 10513428, 10513437, 10513455, 10513467, 10513472, 10513497, 10513504, 10513512, 10514532, 10514558, 10520121, 10531928, 10531994, 10533246, 10535900, 10537062, 10537227, 10538880, 10538903, 10539773, 10541112, 10542981, 10544106, 10545184, 10545187, 10545196, 10545202, 10545208, 10545237, 10545247, 10545450, 10549917, 10550320, 10551250, 10554162, 10556018, 10558740, 10559467, 10565432, 10566254, 10566258, 10566333, 10566346, 10566571,

Downregulated CR genes (only array IDs known)

10566574, 10566585, 10570201, 10576901, 10578405, 10581011, 10582862, 10582882, 10582916, 10585680, 10585972, 10585976, 10585980, 10585982, 10586076, 10587873, 10588786, 10594513, 10598603, 10599956, 10600122, 10601581, 10604996, 10605007, 10607865, 10608136, 10608646, 10608695, 100101_at, 100322_at, 100328_s_at, 100332_s_at, 100348_at, 100358_s_at, 100378_at, 100379_f_at, 100405_at, 100437_g_at, 100566_at, 100577_at, 100708_at, 100718_at, 100722_r_at, 100769_at, 101121_at, 101141_at, 101167_at, 101168_at, 101179_at, 101301_at, 101306_f_at, 101331_f_at, 101336_at, 101566_f_at, 101589_at, 101633_at, 101640_f_at, 101656_f_at, 101670_at, 101675_at, 101687_r_at, 101732_at, 101753_s_at, 101773_r_at, 101780_at, 101784_f_at, 101787_f_at, 101792_at, 101845_s_at, 101869_s_at, 101876_s_at, 101883_s_at, 101886_f_at, 101897_g_at, 101971_at, 102096_f_at, 102125_f_at, 102141_f_at, 102156_f_at, 102161_f_at, 102230_at, 102234_at, 102348_at, 102570_at, 102649_s_at, 102728_f_at, 102821_s_at, 102870_at, 102906_at, 103010_at, 103012_at, 103033_at, 103074_f_at, 103240_f_at, 103267_i_at, 103609_at, 103703_f_at, 103789_at, 103847_at, 103865_at, 103873_i_at, 104263_at, 104502_f_at, 104588_at, 104641_f_at, 1415716_a_at, 1415781_a_at, 1415782_at, 1415788_at, 1415789_a_at, 1415790_at, 1415852_at, 1416128_at, 1416176_at, 1416184_s_at, 1416228_at, 1416336_s_at, 1416461_at, 1416705_at, 1416706_at, 1416790_a_at, 1416811_s_at, 1416884_at, 1416946_a_at, 1416947_s_at, 1417009_at, 1417057_a_at, 1417296_at, 1417714_x_at, 1417810_a_at, 1417886_at, 1417907_at, 1417998_at, 1418072_at, 1418188_a_at, 1418189_s_at, 1418273_a_at, 1418274_at, 1418328_at, 1418335_a_at, 1418520_at, 1419075_s_at, 1419194_s_at, 1419443_at, 1419977_s_at, 1420136_a_at, 1420310_at, 1420809_a_at, 1420810_at, 1420867_at, 1420868_s_at, 1420873_at, 1420874_at, 1421534_at, 1421609_a_at, 1422140_at, 1422141_s_at, 1422155_at, 1422414_a_at, 1422660_at, 1422993_s_at, 1422997_s_at, 1423103_at, 1423262_a_at, 1423263_at, 1423307_s_at, 1423308_at, 1423309_at, 1423397_at, 1423455_at, 1423522_at, 1423617_at, 1423681_at, 1423740_a_at, 1423798_a_at, 1423857_at, 1424008_a_at, 1424300_at, 1424518_at, 1424752_x_at, 1424853_s_at, 1425048_a_at, 1425545_x_at, 1425665_a_at, 1425918_at, 1426154_s_at, 1426266_s_at, 1426327_s_at, 1426394_at, 1426395_s_at, 1426419_at, 1426558_x_at, 1426678_at, 1426679_at, 1426726_at, 1426964_at, 1427174_at, 1427455_x_at, 1427660_x_at, 1427913_at, 1427932_s_at, 1428047_s_at, 1428219_at, 1428658_at, 1428679_s_at, 1428738_a_at, 1429040_at, 1429169_at, 1429184_at, 1429252_at, 1429272_a_at, 1429468_at, 1429504_at, 1429554_at, 1429712_at, 1429784_at, 1430135_at, 1430293_a_at, 1431158_at, 1431591_s_at, 1433549_x_at, 1433592_at, 1433622_at, 1433898_at, 1433903_at, 1434054_at, 1434110_x_at, 1434161_at, 1434280_at, 1434328_at, 1434367_s_at, 1434503_s_at, 1434568_at, 1434861_at, 1435137_s_at, 1435150_at, 1435267_at, 1435384_at, 1435439_at, 1435544_at, 1435574_at, 1435715_x_at, 1435792_at, 1435985_at, 1436247_at, 1436338_at, 1436389_at, 1436412_at, 1436464_at, 1436540_at, 1436933_at, 1436944_x_at, 1436946_s_at, 1436971_x_at, 1437026_at, 1437073_x_at, 1437162_at, 1437350_at, 1437354_at, 1437546_at, 1437547_s_at, 1437600_at, 1437694_at, 1437821_at, 1437987_at, 1438076_at, 1438104_at, 1438221_at, 1438250_s_at, 1438265_at, 1438313_at, 1438398_at, 1438403_s_at, 1438412_at, 1438415_s_at, 1438516_at, 1438543_at, 1438596_at, 1438657_x_at, 1438689_at, 1438714_at, 1438750_at, 1438786_a_at, 1438848_at, 1438906_at, 1438908_at, 1438929_at, 1438937_x_at, 1438997_at, 1439169_at, 1439175_at, 1439224_at, 1439290_at, 1439298_at, 1439305_at, 1439307_at, 1439416_x_at, 1439426_x_at, 1439444_x_at, 1439460_a_at, 1439463_x_at, 1439553_s_at, 1439600_at, 1439616_at, 1439623_at, 1439643_at, 1439831_at, 1439832_at, 1439964_at, 1440020_at, 1440038_at, 1440078_at, 1440151_s_at, 1440219_at, 1440264_at, 1440272_at, 1440284_at, 1440314_at, 1440325_at, 1440351_at, 1440365_at, 1440373_at, 1440388_at, 1440490_at, 1440511_at, 1440533_at,

Downregulated CR genes (only array IDs known)

1440551_at, 1440563_at, 1440624_at, 1440665_at, 1440790_x_at, 1440841_at, 1440896_at, 1441052_at, 1441072_at, 1441108_at, 1441141_at, 1441328_at, 1441370_at, 1441388_at, 1441413_at, 1441498_at, 1441505_at, 1441547_at, 1441565_at, 1441657_at, 1441722_at, 1441765_at, 1442024_at, 1442049_at, 1442097_at, 1442109_at, 1442311_at, 1442331_at, 1442381_at, 1442407_at, 1442470_at, 1442509_at, 1442537_at, 1442569_at, 1442615_at, 1442626_at, 1442641_at, 1442671_at, 1442679_at, 1442710_at, 1442775_at, 1442837_at, 1442867_at, 1442905_at, 1442916_at, 1443003_at, 1443053_at, 1443056_at, 1443075_at, 1443138_at, 1443147_at, 1443196_at, 1443254_at, 1443338_at, 1443489_at, 1443505_at, 1443533_at, 1443620_at, 1443628_at, 1443649_at, 1443673_x_at, 1443835_x_at, 1443872_at, 1443949_at, 1444001_at, 1444032_at, 1444057_at, 1444072_at, 1444140_at, 1444143_at, 1444194_at, 1444265_at, 1444326_at, 1444333_at, 1444377_at, 1444498_at, 1444506_at, 1444531_at, 1444562_at, 1444611_at, 1444735_at, 1444746_at, 1444869_at, 1445065_at, 1445188_at, 1445235_at, 1445267_at, 1445349_at, 1445377_at, 1445379_at, 1445402_at, 1445966_at, 1445980_at, 1446007_at, 1446143_at, 1446167_at, 1446201_at, 1446234_at, 1446278_at, 1446330_at, 1446410_at, 1446504_at, 1446835_at, 1446850_at, 1446868_at, 1447023_at, 1447040_at, 1447064_at, 1447122_at, 1447143_at, 1447156_at, 1447164_at, 1447167_at, 1447227_at, 1447270_at, 1447285_at, 1447329_at, 1447437_at, 1447458_at, 1447502_at, 1447527_at, 1447644_at, 1447651_x_at, 1447977_x_at, 1448037_at, 1448232_x_at, 1448235_s_at, 1448347_a_at, 1448462_at, 1448463_at, 1448548_at, 1448854_s_at, 1449009_at, 1449322_at, 1449347_a_at, 1449480_at, 1449556_at, 1449855_s_at, 1449910_at, 1449972_s_at, 1450007_at, 1450638_at, 1451052_at, 1451082_at, 1451184_at, 1451477_at, 1451600_s_at, 1451683_x_at, 1452052_s_at, 1452300_at, 1452402_at, 1452417_x_at, 1452426_x_at, 1452433_at, 1452463_x_at, 1452485_at, 1452596_at, 1452935_at, 1453078_at, 1453733_a_at, 1454826_at, 1455054_a_at, 1455077_a_at, 1455418_at, 1455503_at, 1455538_at, 1455680_at, 1455773_at, 1455904_at, 1455961_at, 1455998_at, 1456078_x_at, 1456159_at, 1456257_at, 1456272_at, 1456386_at, 1456398_at, 1456433_at, 1456437_x_at, 1456494_a_at, 1456514_at, 1456686_at, 1456782_at, 1456851_at, 1456864_at, 1456904_at, 1456960_at, 1456979_at, 1456991_at, 1457020_at, 1457070_at, 1457178_at, 1457263_at, 1457302_at, 1457367_at, 1457398_at, 1457454_at, 1457468_at, 1457483_at, 1457489_at, 1457553_at, 1457588_at, 1457666_s_at, 1457731_at, 1457765_at, 1457832_at, 1457842_at, 1457908_at, 1457988_at, 1457991_at, 1458134_at, 1458144_at, 1458205_at, 1458282_at, 1458290_at, 1458324_x_at, 1458356_at, 1458589_at, 1458618_at, 1458637_x_at, 1458686_at, 1458832_at, 1458847_at, 1458943_at, 1458994_at, 1459008_at, 1459061_at, 1459187_at, 1459238_at, 1459325_at, 1459360_at, 1459371_at, 1459434_at, 1459488_at, 1459556_at, 1459734_at, 1459813_at, 1459948_at, 1460052_at, 1460086_at, 1460138_at, 1460151_at, 1460221_at, 1460232_s_at, 1460416_s_at, 1460590_s_at, 160119_at, 160159_at, 160193_at, 160682_at, 160906_i_at, 161057_at, 161077_f_at, 161173_f_at, 161227_r_at, 161232_r_at, 161362_at, 161406_at, 161410_r_at, 161411_i_at, 161448_f_at, 161531_r_at, 161590_r_at, 161617_f_at, 161724_r_at, 161815_f_at, 161930_r_at, 162045_r_at, 162083_f_at, 162195_i_at, 162309_at, 162368_r_at, 162457_f_at, 162496_r_at, 92198_s_at, 92217_s_at, 92470_f_at, 92611_at, 92632_at, 92665_f_at, 92812_f_at, 92847_s_at, 93023_f_at, 93059_at, 93077_s_at, 93086_at, 93087_r_at, 93321_at, 93646_at, 93714_f_at, 93833_s_at, 93865_s_at, 93874_s_at, 93881_i_at, 93882_f_at, 93908_f_at, 93968_at, 94000_at, 94007_at, 94028_f_at, 94042_f_at, 94224_s_at, 94259_at, 94424_at, 94477_at, 94489_at, 94541_at, 94756_at, 94805_f_at, 94838_r_at, 94850_at, 95018_r_at, 95081_at, 95124_i_at, 95398_at, 95409_at, 95415_f_at, 95634_at, 95635_g_at, 95636_at, 95654_at, 95657_f_at, 95773_at, 95852_at, 95858_at, 95869_at, 95892_at, 95901_f_at, 95919_at, 95926_at, 95936_at, 95937_at, 95988_r_at, 96017_at, 96135_at, 96220_at, 96416_f_at, 96482_at, 96483_at, 96486_at,

Downregulated CR genes (only array IDs known)

96489_at, 96522_at, 96536_at, 96547_at, 96550_at, 96555_at, 96653_at, 96959_at, 96964_at, 96967_at, 96970_at, 97135_at, 97173_f_at, 97176_at, 97205_at, 97287_at, 97288_at, 97448_at, 97462_at, 97464_at, 97479_at, 97523_i_at, 97524_f_at, 97541_f_at, 97566_f_at, 97567_f_at, 97575_f_at, 97578_at, 97666_r_at, 97702_at, 97706_at, 97708_at, 98065_at, 98096_f_at, 98438_f_at, 98446_s_at, 98472_at, 98516_at, 98600_at, 98761_i_at, 99143_at, 99144_s_at, 99681_at, 99702_at, 99821_at, 99866_at, 99872_s_at, AFFX-18SRNAMur/X00686_5_at, AFFX-18SRNAMur/X00686_M_at, AFFX-BioB-3_at, AFFX-BioB-5_at, AFFX-BioB-M_at, AFFX-BioC-3_at, AFFX-BioC-5_at, AFFX-BioDn-3_at, AFFX-BioDn-5_at, AFFX-CreX-3_at, AFFX-CreX-5_at, AFFX-DapX-3_at, AFFX-DapX-M_at, AFFX-r2-Bs-dap-3_at, AFFX-r2-Bs-dap-5_at, AFFX-r2-Bs-dap-M_at, AFFX-r2-Ec-bioB-3_at, AFFX-r2-Ec-bioB-5_at, AFFX-r2-Ec-bioB-M_at, AFFX-r2-Ec-bioC-3_at, AFFX-r2-Ec-bioC-5_at, AFFX-r2-Ec-bioD-3_at, AFFX-r2-Ec-bioD-5_at, AFFX-YEL002c/WBP1_at

D: Pathways mentioned by CR in overrepresentation analysis and the number of overlapping genes with plant bioactives

Pathways	regu- la- tion	genes up- regulated by CR	Overlap					genes down- regulated by CR	Overlap					
			R	P	Q	Qi	Qi		R	P	Q	Qi	Qi	
(deoxy)ribose phosphate degradation - MouseCyc	↑	4 of 10	1 of 4	2 of 4	1 of 4	1 of 4	1 of 4	2 of 10	1 of 2	1 of 2	1 of 2	0 of 2	0 of 2	
2-amino-3-carboxymuconate semialdehyde degradation to glutaryl-CoA - MouseCyc	↑	2 of 2	0 of 2	0 of 1	0 of 1	0 of 2	0 of 2	0 of 2						
2-methylbutyrate biosynthesis - MouseCyc	↓	1 of 5	0 of 1	0 of 1	0 of 1	1 of 1	1 of 1	4 of 5	0 of 4	3 of 3	0 of 3	1 of 4	1 of 4	
3, -UTR-mediated translational regulation - Reactome	↑	24 of 100	0 of 12	1 of 10	0 of 10	2 of 23	1 of 23	15 of 100	2 of 14	4 of 13	0 of 13	1 of 15	0 of 15	
4-hydroxyproline degradation I - MouseCyc	↑	2 of 4	1 of 2	1 of 2	1 of 2	0 of 2	1 of 2	0 of 4						
ABC transporters - Mus musculus (mouse) - KEGG	↑	14 of 46	5 of 14	0 of 14	0 of 14	0 of 14	0 of 14	11 of 46	3 of 11	6 of 11	0 of 11	1 of 11	0 of 11	
ABCA transporters in lipid homeostasis - Reactome	↑	6 of 13	4 of 6	0 of 6	0 of 6	0 of 6	0 of 6	3 of 13	1 of 3	0 of 3	0 of 3	0 of 3	0 of 3	
ABC-family proteins mediated transport - Reactome	↑	12 of 31	5 of 12	0 of 12	0 of 12	0 of 12	0 of 12	6 of 31	1 of 6	3 of 6	0 of 6	0 of 6	0 of 6	
acetyl-CoA biosynthesis (from pyruvate) - MouseCyc	↑	3 of 5	0 of 3	0 of 3	0 of 3	0 of 3	0 of 3	0 of 5						
Activated TLR4 signalling - Reactome	↓	13 of 94	4 of 12	2 of 11	2 of 11	1 of 12	0 of 12	28 of 94	2 of 26	16 of 28	4 of 28	1 of 28	0 of 28	
Activation of Gene Expression by SREBP (SREBF) - Reactome	↑	3 of 7	0 of 3	0 of 3	0 of 3	1 of 3	1 of 3	0 of 7						
acyl-CoA hydrolysis - MouseCyc	↓	4 of 11	1 of 4	1 of 4	0 of 4	1 of 4	2 of 4	5 of 11	0 of 5	1 of 5	1 of 5	0 of 5	0 of 5	
Adaptive Immune System - Reactome	↓	56 of 441	15 of 52	4 of 52	6 of 52	3 of 55	2 of 55	98 of 441	15 of 93	33 of 93	3 of 93	13 of 97	8 of 97	
adenosine nucleotides de novo biosynthesis - MouseCyc	↑	8 of 22	1 of 7	0 of 7	0 of 8	0 of 8	0 of 8	2 of 22	0 of 2	2 of 2	0 of 2	0 of 2	0 of 2	
Adipocytokine signaling pathway - Mus musculus (mouse) - KEGG	↑	18 of 71	4 of 16	1 of 18	1 of 18	4 of 18	2 of 18	16 of 71	3 of 15	7 of 16	4 of 16	2 of 16	0 of 16	
aerobic respiration -- electron donor II - MouseCyc	↑	24 of 80	0 of 22	0 of 22	0 of 22	1 of 24	0 of 24	6 of 80	1 of 5	3 of 5	0 of 5	1 of 5	1 of 5	
Alanine and aspartate metabolism - Wikipathways	↑	4 of 12	1 of 4	1 of 4	1 of 4	1 of 4	1 of 4	4 of 12	0 of 4	3 of 4	0 of 4	1 of 4	1 of 4	

128

Alanine, aspartate and glutamate metabolism - Mus musculus (mouse) - KEGG	↑	14 of 33	1 of 11	3 of 14	3 of 14	0 of 14	1 of 14	8 of 33	1 of 8	5 of 8	0 of 8	1 of 8	1 of 8
alpha-linolenic (omega3) and linoleic (omega6) acid metabolism - Reactome	↑↓	4 of 12	1 of 4	1 of 4	0 of 4	2 of 4	0 of 4	8 of 12	2 of 8	1 of 8	0 of 8	2 of 8	1 of 8
alpha-linolenic acid (ALA) metabolism - Reactome	↑↓	4 of 12	1 of 4	1 of 4	0 of 4	2 of 4	0 of 4	8 of 12	2 of 8	1 of 8	0 of 8	2 of 8	1 of 8
Amino Acid metabolism - Wikipathways	↑	4 of 6	0 of 4	0 of 6									
Amino acid synthesis and interconversion (transamination) - Reactome	↑	8 of 17	1 of 7	2 of 8	1 of 8	0 of 8	1 of 8	1 of 17	0 of 1	0 of 1	0 of 1	0 of 1	0 of 1
Amino sugar and nucleotide sugar metabolism - Mus musculus (mouse) - KEGG	↑	14 of 48	3 of 14	1 of 14	2 of 14	0 of 14	0 of 14	12 of 48	5 of 12	4 of 12	2 of 12	2 of 12	0 of 12
Androgen biosynthesis - Reactome	↓	1 of 13	1 of 1	0 of 1	0 of 1	1 of 1	1 of 1	4 of 13	3 of 4	3 of 4	2 of 4	2 of 4	1 of 4
Androgen Receptor Signaling Pathway - Wikipathways	↓	15 of 109	4 of 12	2 of 13	0 of 13	1 of 13	1 of 13	37 of 109	6 of 35	13 of 35	2 of 35	6 of 37	4 of 37
Antigen processing and presentation - Mus musculus (mouse) - KEGG	↓	13 of 76	5 of 10	1 of 11	2 of 11	0 of 11	0 of 11	26 of 76	3 of 22	11 of 22	3 of 23	6 of 23	4 of 25
Antigen processing: Ubiquitination & Proteasome degradation - Reactome	↓	12 of 128	3 of 11	0 of 10	2 of 10	1 of 11	0 of 11	34 of 128	8 of 31	14 of 31	1 of 34	3 of 34	2 of 34
Antigen processing-Cross presentation - Reactome	↓	6 of 33	2 of 5	0 of 5	0 of 5	0 of 6	0 of 6	13 of 33	0 of 11	3 of 11	0 of 10	4 of 10	3 of 12
Antiviral mechanism by IFN-stimulated genes - Reactome	↓	2 of 25	1 of 2	0 of 2	0 of 2	0 of 2	0 of 2	9 of 25	1 of 8	5 of 8	0 of 8	4 of 9	4 of 9
Apoptotic cleavage of cellular proteins - Reactome	↑	9 of 34	3 of 8	0 of 8	1 of 8	1 of 9	1 of 9	9 of 34	1 of 9	3 of 9	1 of 9	2 of 9	0 of 9
Arachidonic acid metabolism - Mus musculus (mouse) - KEGG	↑↓	25 of 71	7 of 23	4 of 22	3 of 22	5 of 25	6 of 25	19 of 71	4 of 17	7 of 17	1 of 18	6 of 18	5 of 19
Arachidonic acid metabolism - Reactome	↑↓	19 of 71	2 of 17	2 of 16	1 of 16	4 of 18	4 of 18	19 of 71	4 of 16	8 of 16	1 of 18	7 of 18	4 of 19
Arginine and proline metabolism - Mus musculus (mouse) - KEGG	↑	16 of 55	3 of 13	1 of 16	1 of 16	0 of 16	2 of 16	9 of 55	0 of 7	2 of 9	0 of 9	0 of 9	0 of 9
arginine biosynthesis IV - MouseCyc	↑	4 of 5	0 of 4	0 of 5									
arginine degradation I (arginase pathway) - MouseCyc	↑	3 of 4	0 of 3	0 of 3	0 of 3	0 of 3	1 of 3	0 of 4					
arsenate detoxification I (glutaredoxin) - MouseCyc	↑	5 of 6	2 of 5	0 of 5	0 of 5	0 of 5	0 of 5	1 of 6	1 of 1	0 of 1	0 of 1	0 of 1	0 of 1
Ascorbate and aldarate metabolism - Mus musculus (mouse) - KEGG	↑	6 of 22	3 of 5	0 of 3	1 of 3	0 of 3	0 of 3	8 of 22	3 of 7	3 of 8	1 of 8	3 of 8	2 of 8

beta-Alanine metabolism - <i>Mus musculus</i> (mouse) - KEGG	↓	3 of 30	1 of 3	0 of 3	0 of 3	1 of 3	1 of 3	8 of 30	0 of 7	6 of 8	0 of 8	1 of 8	1 of 8
Beta-oxidation of pristanoyl-CoA - Reactome	↓	0 of 8						7 of 8	0 of 7	1 of 7	0 of 7	4 of 7	2 of 7
Beta-oxidation of very long chain fatty acids - Reactome	↓	1 of 7	1 of 1	0 of 1	0 of 1	0 of 1	0 of 1	6 of 7	2 of 6	1 of 6	0 of 6	1 of 6	0 of 6
Bile acid and bile salt metabolism - Reactome	↓	9 of 36	3 of 9	1 of 9	1 of 9	0 of 9	0 of 9	19 of 36	3 of 19	7 of 19	1 of 19	8 of 19	4 of 19
bile acid biosynthesis, neutral pathway - MouseCyc	↓	2 of 6	1 of 2	1 of 2	1 of 2	0 of 2	0 of 2	4 of 6	1 of 4	0 of 4	1 of 4	0 of 4	0 of 4
Bile secretion - <i>Mus musculus</i> (mouse) - KEGG	↑↓	24 of 71	7 of 23	2 of 23	1 of 23	2 of 24	1 of 24	19 of 71	5 of 19	12 of 19	3 of 19	5 of 19	5 of 19
Binding and Uptake of Ligands by Scavenger Receptors - Reactome	↑	11 of 35	4 of 11	1 of 11	2 of 11	0 of 11	0 of 11	11 of 35	2 of 11	2 of 11	1 of 11	1 of 11	0 of 11
Biological oxidations - Reactome	↑↓	62 of 158	22 of 54	5 of 51	9 of 51	12 of 59	11 of 59	50 of 158	10 of 44	16 of 47	7 of 47	22 of 49	15 of 49
Biosynthesis of the N-glycan precursor (dolichol lipid-linked oligosaccharide, LLO) and transfer to a nascent protein - Reactome	↓	2 of 29	0 of 2	0 of 2	1 of 2	0 of 2	0 of 2	11 of 29	4 of 11	2 of 11	0 of 11	0 of 11	1 of 11
Biosynthesis of unsaturated fatty acids - <i>Mus musculus</i> (mouse) - KEGG	↑↓	10 of 25	2 of 10	3 of 10	0 of 10	4 of 10	3 of 10	12 of 25	1 of 4	3 of 4	0 of 4	0 of 4	0 of 4
Biotin transport and metabolism - Reactome	↑	5 of 10	1 of 5	1 of 5	0 of 5	0 of 5	0 of 5	2 of 10	1 of 2	1 of 2	0 of 2	0 of 2	0 of 2
biotin-carboxyl carrier protein assembly - MouseCyc	↑	2 of 3	0 of 2	1 of 2	0 of 2	2 of 2	2 of 2	0 of 3					
bupropion degradation - MouseCyc	↑↓	14 of 36	3 of 11	1 of 12	3 of 12	2 of 14	3 of 14	16 of 36	2 of 14	9 of 16	1 of 16	10 of 16	8 of 16
Butanoate metabolism - <i>Mus musculus</i> (mouse) - KEGG	↓	7 of 29	1 of 7	0 of 7	0 of 7	1 of 7	1 of 7	11 of 29	0 of 11	9 of 10	0 of 10	4 of 11	4 of 11
Caffeine metabolism - <i>Mus musculus</i> (mouse)	↑	5 of 8	1 of 5	0 of 4	0 of 4	1 of 5	1 of 5	1 of 8	0 of 1	1 of 1	0 of 1	0 of 1	0 of 1
Cap-dependent Translation Initiation - Reactome	↑	26 of 108	0 of 14	1 of 12	0 of 12	2 of 15	1 of 25	20 of 108	2 of 19	7 of 18	0 of 18	1 of 20	0 of 20
Caspase-mediated cleavage of cytoskeletal proteins - Reactome	↑	7 of 12	3 of 6	0 of 7	1 of 7	0 of 7	1 of 7	4 of 12	0 of 4	1 of 4	0 of 4	0 of 4	0 of 4
catecholamine biosynthesis - MouseCyc	↓	0 of 6						2 of 6	1 of 2	1 of 2	0 of 2	0 of 2	0 of 2
CDT1 association with the CDC6:ORC:origin complex - Reactome	↑	6 of 14	2 of 5	0 of 4	0 of 4	0 of 5	0 of 5	2 of 14	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2
Chemical carcinogenesis - <i>Mus musculus</i> (mouse)	↑↓	39 of 87	11 of 33	1 of 30	3 of 30	7 of 35	7 of 35	20 of 87	3 of 18	11 of 19	2 of 19	12 of 19	11 of 19

Cholesterol biosynthesis - Reactome	↑	19 of 22	14 of 19	6 of 19	2 of 19	1 of 19	1 of 19	6 of 22	1 of 6	2 of 6	1 of 6	3 of 6	1 of 6		
Cholesterol biosynthesis - Wikipathways	↑	15 of 15	12 of 15	6 of 15	2 of 15	1 of 15	1 of 15	5 of 15	1 of 5	2 of 5	1 of 5	3 of 5	1 of 5		
cholesterol biosynthesis I - MouseCyc	↑	11 of 12	8 of 11	3 of 11	1 of 11	1 of 11	1 of 11	4 of 12	0 of 4	1 of 4	1 of 4	3 of 4	0 of 4		
cholesterol biosynthesis II (via 24,25-dihydrolanosterol) - MouseCyc	↑	8 of 8	5 of 8	2 of 8	1 of 8	1 of 8	1 of 8	3 of 8	0 of 3	1 of 3	1 of 3	3 of 3	0 of 3		
cholesterol biosynthesis III (via desmosterol) - MouseCyc	↑	10 of 11	7 of 10	3 of 10	1 of 10	1 of 10	1 of 10	4 of 11	0 of 4	1 of 4	1 of 4	3 of 4	0 of 4		
Circadian Clock - Reactome	↑	6 of 13	3 of 6	1 of 6	2 of 6	0 of 6	0 of 6	2 of 13	1 of 2	2 of 2	1 of 2	0 of 2	0 of 2		
Circadian clock tutorial Cgrove - Wikipathways	↑	4 of 6	3 of 4	1 of 4	2 of 4	0 of 4	0 of 4	2 of 6	1 of 2	2 of 2	1 of 2	0 of 2	0 of 2		
Circadian rhythm - Mus musculus (mouse) - KEGG	↑	11 of 30	7 of 11	2 of 11	2 of 11	0 of 11	0 of 11	6 of 30	1 of 5	3 of 6	2 of 6	0 of 6	0 of 6		
Citrate cycle (TCA cycle) - Mus musculus (mouse) - KEGG	↑	18 of 32	4 of 18	0 of 18	0 of 18	1 of 18	1 of 18	1 of 32	0 of 1	1 of 1	0 of 1	0 of 1	0 of 1		
Citric acid cycle (TCA cycle) - Reactome	↑	12 of 20	2 of 12	0 of 12	0 of 12	0 of 12	0 of 12	0 of 12							
citrulline biosynthesis - MouseCyc	↑	4 of 9	0 of 4	0 of 4	0 of 4	0 of 4	1 of 4	0 of 9							
citrulline-nitric oxide cycle - MouseCyc	↑	2 of 7	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2	0 of 7							
Class I MHC mediated antigen processing & presentation - Reactome	↓	22 of 173	6 of 20	0 of 19	2 of 19	1 of 21	0 of 21	51 of 173	9 of 46	18 of 48	1 of 48	7 of 50	5 of 50		
Classical antibody-mediated complement activation - Reactome	↓	0 of 8						5 of 8	0 of 5	2 of 5	0 of 5	3 of 5	2 of 5		
Clathrin derived vesicle budding - Reactome	↓	6 of 60	2 of 6	1 of 6	0 of 6	0 of 6	0 of 6	19 of 60	2 of 16	7 of 19	1 of 19	3 of 19	1 of 19		
Cleavage of Growing Transcript in the Termination Region - Reactome	↓	4 of 43	0 of 6	0 of 6	1 of 6	0 of 6	0 of 6	16 of 43	1 of 16	7 of 15	0 of 15	3 of 16	2 of 16		
colanic acid building blocks biosynthesis - MouseCyc	↑↓	6 of 14	1 of 6	1 of 6	1 of 6	0 of 6	0 of 6	5 of 14	2 of 5	0 of 5	1 of 5	1 of 5	0 of 5		
Complement Activation, Classical Pathway - Wikipathways	↓	2 of 17	1 of 2	0 of 2	0 of 2	0 of 2	0 of 2	11 of 17	0 of 10	5 of 11	0 of 11	7 of 11	4 of 11		
Complement and coagulation cascades - Mus musculus (mouse) - KEGG	↓	12 of 77	6 of 11	1 of 11	0 of 11	0 of 12	0 of 12	24 of 77	1 of 23	8 of 22	0 of 22	12 of 24	8 of 24		
Complement and Coagulation Cascades - Wikipathways	↓	10 of 61	5 of 10	1 of 10	0 of 10	0 of 10	0 of 10	20 of 61	1 of 19	6 of 20	0 of 20	10 of 20	8 of 20		

Complement cascade - Reactome	↓	4 of 36	3 of 4	0 of 4	0 of 4	0 of 4	0 of 4	17 of 36	1 of 16	5 of 16	0 of 16	9 of 17	5 of 17
Cysteine and methionine metabolism - Mus musculus (mouse) - KEGG	↑	9 of 35	3 of 9	2 of 9	3 of 9	0 of 9	2 of 9	11 of 35	4 of 11	3 of 11	1 of 11	1 of 11	2 of 11
cysteine biosynthesis II - MouseCyc	↑	2 of 2	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2					
cysteine biosynthesis/homocysteine degradation - MouseCyc	↑	2 of 2	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2					
Cysteine formation from homocysteine - Reactome	↑	2 of 2	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2					
Cytochrome P450 - arranged by substrate type - Reactome	↑↓	31 of 78	9 of 24	3 of 23	6 of 23	7 of 30	7 of 30	27 of 78	5 of 22	11 of 25	5 of 25	13 of 27	10 of 27
Cytochrome P450 - Wikipathways	↑	12 of 40	4 of 11	2 of 12	3 of 12	1 of 12	1 of 12	10 of 40	4 of 9	1 of 10	2 of 10	3 of 10	1 of 10
Cytokine Signaling in Immune system - Reactome	↓	20 of 190	8 of 19	4 of 20	5 of 20	3 of 20	3 of 20	50 of 190	7 of 48	22 of 49	3 of 49	12 of 50	8 of 50
Cytoplasmic Ribosomal Proteins - Wikipathways	↑	21 of 78	0 of 9	1 of 9	0 of 9	0 of 21	0 of 21	5 of 78	0 of 4	0 of 3	0 of 3	1 of 5	0 of 5
Cytosolic sensors of pathogen-associated DNA - Reactome	↓	9 of 55	3 of 7	1 of 7	1 of 7	1 of 8	0 of 8	17 of 55	1 of 17	8 of 16	1 of 16	3 of 17	3 of 17
Cytosolic sulfonation of small molecules - reactome	↑	7 of 16	3 of 6	0 of 6	1 of 6	2 of 7	2 of 7	5 of 16	1 of 5	1 of 5	2 of 5	0 of 5	0 of 5
Cytosolic tRNA aminoacylation - Reactome	↓	0 of 23						10 of 23	1 of 8	6 of 10	0 of 10	1 of 10	1 of 10
Deadenylation of mRNA - Reactome	↓	3 of 21	0 of 3	0 of 3	0 of 3	1 of 3	0 of 3	10 of 21	3 of 9	5 of 10	0 of 10	0 of 10	0 of 10
Deadenylation-dependent mRNA decay - Reactome	↓	5 of 45	0 of 4	0 of 5	0 of 5	1 of 5	0 of 5	15 of 45	3 of 13	7 of 15	1 of 15	0 of 15	0 of 15
Degradation of cysteine and homocysteine - Reactome	↓	3 of 8	0 of 3	0 of 3	0 of 3	0 of 3	0 of 3	5 of 8	3 of 5	2 of 5	0 of 5	2 of 5	1 of 5
Destabilization of mRNA by AUF1 (hnRNP D0) - Reactome	↓	3 of 7	0 of 1	2 of 3	2 of 3	0 of 2	0 of 2	7 of 7	2 of 4	4 of 7	3 of 7	0 of 6	2 of 6
Diurnally Regulated Genes with Circadian Orthologs - Wikipathways	↑↓	21 of 48	9 of 18	3 of 21	5 of 21	1 of 21	1 of 21	18 of 48	4 of 17	11 of 18	4 of 18	7 of 18	3 of 18
Drug metabolism - cytochrome P450 - Mus musculus (mouse) - KEGG	↑↓	39 of 88	12 of 35	1 of 32	3 of 32	7 of 36	7 of 36	25 of 88	6 of 23	12 of 24	2 of 24	14 of 24	13 of 24
Drug metabolism - other enzymes - Mus musculus (mouse) - KEGG	↑↓	16 of 51	7 of 14	2 of 12	3 of 12	2 of 13	2 of 13	19 of 51	6 of 18	11 of 19	2 of 19	6 of 19	6 of 19
EGFR1 Signaling Pathway - Wikipathways	↓	25 of 175	5 of 25	2 of 25	5 of 25	0 of 25	0 of 25	45 of 175	6 of 45	14 of 45	4 of 45	7 of 45	3 of 45

Electron Transport Chain - Wikipathways	↑	28 of 93	0 of 23	0 of 26	0 of 26	1 of 28	0 of 28	4 of 93	0 of 2	3 of 3	0 of 3	0 of 3	0 of 3
Endogenous sterols - Reactome	↑↓	4 of 14	3 of 4	1 of 4	1 of 4	1 of 4	1 of 4	4 of 14	1 of 4	0 of 4	1 of 4	1 of 4	0 of 4
epoxysqualene biosynthesis - Mousecyc	↑	2 of 2	1 of 2	1 of 2	1 of 2	0 of 2	0 of 2	1 of 2	0 of 1	0 of 1	0 of 1	1 of 1	0 of 1
Estrogen metabolism - Wikipathways	↑	8 of 11	3 of 8	0 of 5	1 of 5	2 of 6	2 of 6	1 of 11	0 of 1	0 of 1	0 of 1	1 of 1	0 of 1
Eukaryotic Translation Elongation - Reactome	↑	23 of 82	0 of 11	1 of 10	0 of 10	3 of 22	1 of 22	6 of 82	0 of 5	0 of 4	0 of 4	2 of 6	0 of 6
Eukaryotic Translation Initiation - Reactome	↑	26 of 108	0 of 14	1 of 12	0 of 12	2 of 25	1 of 25	20 of 108	2 of 19	7 of 18	0 of 18	1 of 20	0 of 20
Eukaryotic Translation Termination - Reactome	↑	23 of 79	0 of 11	1 of 10	0 of 10	2 of 22	1 of 22	5 of 79	0 of 4	0 of 3	0 of 3	1 of 5	0 of 5
Exercise-induced Circadian Regulation - Wikipathways	↑↓	22 of 49	9 of 19	4 of 22	5 of 22	1 of 22	1 of 11	20 of 49	4 of 18	12 of 19	5 of 19	7 of 19	3 of 17
Facilitative Na ⁺ -independent glucose transporters - reactome	↑	4 of 11	1 of 4	0 of 4	0 of 4	1 of 4	0 of 4	0 of 11					
Fat digestion and absorption - Mus musculus (mouse)	↓	5 of 40	3 of 5	0 of 5	0 of 5	0 of 5	0 of 5	12 of 40	2 of 12	5 of 11	2 of 11	4 of 12	1 of 12
fatty acid activation - MouseCyc	↑	3 of 6	0 of 3	0 of 6									
Fatty Acid Beta Oxidation - Wikipathways	↑↓	7 of 34	2 of 7	1 of 7	0 of 7	0 of 7	0 of 7	16 of 34	2 of 16	8 of 16	1 of 16	3 of 16	2 of 16
Fatty acid biosynthesis - Mus musculus (mouse) - KEGG	↑	3 of 6	0 of 3	1 of 3	0 of 3	0 of 3	0 of 3	1 of 6	0 of 1	0 of 1	0 of 1	0 of 1	0 of 1
Fatty Acid Biosynthesis - Wikipathways	↑	11 of 22	1 of 10	3 of 11	0 of 11	1 of 11	1 of 11	5 of 22	0 of 5	2 of 5	1 of 5	2 of 5	2 of 5
Fatty acid degradation - Mus musculus (mouse) - KEGG	↑↓	11 of 45	4 of 10	1 of 10	1 of 11	4 of 11	3 of 11	20 of 45	4 of 18	9 of 18	0 of 18	5 of 20	5 of 20
Fatty acid elongation - Mus musculus (mouse) - KEGG	↑↓	7 of 24	1 of 7	1 of 7	0 of 7	2 of 7	3 of 7	10 of 24	2 of 10	4 of 10	1 of 10	3 of 10	2 of 10
fatty acid β-oxidation I - MouseCyc	↓	5 of 27	0 of 5	0 of 5	0 of 5	1 of 5	1 of 5	12 of 27	0 of 12	7 of 12	0 of 12	3 of 12	1 of 12
fatty acid β-oxidation III (unsaturated, odd number) - MouseCyc	↓	2 of 3	0 of 2	0 of 2	0 of 2	1 of 2	1 of 2	3 of 3	0 of 3	2 of 3	0 of 3	0 of 3	0 of 3
fatty acid β-oxidation IV (unsaturated, even number) - MouseCyc	↓	2 of 8	0 of 2	0 of 2	0 of 2	1 of 2	1 of 2	7 of 8	0 of 7	4 of 7	1 of 7	1 of 7	1 of 7
Fatty acid, triacylglycerol, and ketone body metabolism - Reactome	↑↓	23 of 85	2 of 20	1 of 23	1 of 23	2 of 23	3 of 23	30 of 85	9 of 26	11 of 30	3 of 30	5 of 30	3 of 30

Fatty acids - Reactome	↑↓	6 of 16	2 of 4	1 of 5	1 of 5	3 of 6	2 of 6	6 of 16	2 of 5	2 of 5	0 of 5	3 of 6	2 of 6
Fatty Acyl-CoA Biosynthesis - Reactome	↑	9 of 17	0 of 8	0 of 9	0 of 9	2 of 9	2 of 9	4 of 17	2 of 4	2 of 4	0 of 4	2 of 4	1 of 4
formaldehyde oxidation I - MouseCyc	↑	2 of 5	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2	1 of 5	0 of 1	0 of 1	0 of 1	0 of 1	0 of 1
Formation of a pool of free 40S subunits - Reactome	↑	24 of 89	0 of 12	1 of 10	0 of 10	2 of 23	1 of 23	8 of 89	0 of 7	0 of 6	0 of 6	1 of 8	0 of 8
Formation of ATP by chemiosmotic coupling - Reactome	↑	6 of 13	0 of 4	0 of 6	0 of 6	0 of 6	0 of 6	0 of 13					
Formation of the Editosome - Reactome	↓	0 of 2						2 of 2	1 of 2	2 of 2	0 of 2	0 of 2	0 of 2
Fructose and mannose metabolism - Mus musculus (mouse) - KEGG	↑	12 of 39	3 of 12	1 of 12	0 of 12	0 of 12	0 of 12	5 of 39	1 of 5	1 of 5	0 of 5	1 of 5	1 of 5
G1 to S cell cycle control - Wikipathways	↑	16 of 61	4 of 16	3 of 16	1 of 16	0 of 16	0 of 16	10 of 61	1 of 10	2 of 10	1 of 10	0 of 10	0 of 10
Galactose metabolism - Mus musculus (mouse) - KEGG	↑	10 of 32	3 of 10	1 of 10	1 of 10	1 of 10	1 of 10	5 of 32	3 of 5	2 of 5	3 of 5	1 of 5	1 of 5
Gamma-carboxylation, transport, and amino-terminal cleavage of proteins - Reactome	↑	4 of 10	1 of 4	0 of 4	0 of 4	0 of 4	0 of 4	4 of 10	0 of 4	2 of 4	0 of 4	1 of 4	1 of 4
GDP-mannose biosynthesis I - MouseCyc	↓	1 of 6	0 of 1	0 of 1	0 of 1	0 of 1	0 of 1	2 of 6	1 of 2	0 of 2	0 of 2	0 of 2	0 of 2
Gene Expression - Reactome	↓	107 of 646	8 of 83	3 of 87	3 of 87	4 of 104	3 of 104	167 of 646	24 of 148	59 of 158	9 of 158	13 of 163	6 of 163
Gluconeogenesis - Reactome	↑	18 of 33	3 of 16	3 of 18	1 of 18	1 of 18	2 of 18	4 of 33	0 of 4	2 of 4	0 of 4	1 of 4	0 of 4
gluconeogenesis I - MouseCyc	↑	12 of 24	1 of 9	1 of 12	0 of 12	1 of 12	1 of 12	0 of 24					
Glucose metabolism - Reactome	↑	27 of 67	5 of 24	3 of 26	1 of 26	2 of 27	2 of 27	10 of 27	1 of 10	7 of 10	1 of 10	1 of 10	0 of 10
Glucuronidation - Reactome	↑	4 of 5	3 of 4	0 of 2	1 of 2	0 of 2	0 of 2	2 of 5	1 of 2	0 of 2	0 of 2	1 of 2	0 of 2
glutathione biosynthesis - MouseCyc	↓	0 of 3						3 of 3	2 of 3	2 of 3	0 of 3	2 of 3	1 of 3
Glutathione conjugation - Reactome	↑	11 of 26	4 of 11	0 of 11	0 of 11	1 of 11	1 of 11	8 of 26	2 of 7	2 of 7	0 of 7	4 of 7	3 of 7
Glutathione metabolism - Mus musculus (mouse) - KEGG	↑	27 of 54	8 of 26	3 of 26	0 of 26	3 of 27	2 of 27	14 of 54	2 of 11	3 of 13	0 of 13	5 of 13	3 of 13
Glutathione metabolism - Wikipathways	↑	7 of 19	4 of 7	1 of 6	0 of 6	2 of 7	1 of 7	7 of 19	2 of 6	3 of 7	0 of 7	3 of 7	1 of 7

134

Glutathione synthesis and recycling - Reactome	↓	0 of 6							4 of 6	2 of 4	2 of 4	0 of 4	2 of 4	1 of 4
glutathione-mediated detoxification - MouseCyc	↑	15 of 26	4 of 14	0 of 14	0 of 14	3 of 15	2 of 15		6 of 26	0 of 4	0 of 5	0 of 5	3 of 5	2 of 5
glycerol degradation IV - MouseCyc	↓	0 of 3						3 of 6	2 of 3	1 of 3	0 of 3	0 of 3	0 of 3	0 of 3
Glycerolipid metabolism - Mus musculus (mouse) - KEGG	↑	15 of 55	6 of 14	0 of 15	1 of 15	1 of 15	1 of 15	11 of 55	3 of 10	5 of 11	3 of 11	2 of 11	2 of 11	2 of 11
Glycerophospholipid biosynthesis - Reactome	↑↓	20 of 85	6 of 18	1 of 20	2 of 20	0 of 20	2 of 20	25 of 85	7 of 24	6 of 24	2 of 24	5 of 25	2 of 25	2 of 25
Glycerophospholipid metabolism - Mus musculus (mouse) - KEGG	↓	18 of 90	6 of 16	0 of 18	1 of 18	0 of 18	1 of 18	27 of 90	8 of 26	7 of 26	2 of 26	6 of 27	4 of 27	4 of 27
Glycine, serine and threonine metabolism - Mus musculus (mouse) - KEGG	↑↓	15 of 38	4 of 13	1 of 15	2 of 15	0 of 15	0 of 15	11 of 38	5 of 11	6 of 10	1 of 10	2 of 11	3 of 11	3 of 11
glycogen biosynthesis II (from UDP-D-Glucose) - MouseCyc	↑	4 of 5	1 of 4	0 of 5										
Glycogen synthesis - Reactome	↑	4 of 9	1 of 4	0 of 4	0 of 4	0 of 4	0 of 4	1 of 9	1 of 1	1 of 1	1 of 1	0 of 1	0 of 1	0 of 1
Glycolysis - Reactome	↑	14 of 30	2 of 12	2 of 14	0 of 14	2 of 14	1 of 14	3 of 30	0 of 3	2 of 3	0 of 3	0 of 3	0 of 3	0 of 3
Glycolysis / Gluconeogenesis - Mus musculus (mouse) - KEGG	↑	23 of 64	5 of 21	2 of 23	1 of 23	3 of 23	2 of 23	7 of 64	2 of 6	5 of 7	2 of 7	2 of 7	2 of 7	2 of 7
Glycolysis and Gluconeogenesis - Wikipathways	↑	26 of 49	5 of 24	2 of 26	2 of 26	4 of 26	3 of 26	5 of 49	2 of 5	3 of 5	2 of 5	2 of 5	2 of 5	1 of 5
glycolysis I - MouseCyc	↑	14 of 26	2 of 12	1 of 14	0 of 14	2 of 14	1 of 14	0 of 26						
glycolysis III - MouseCyc	↑	14 of 25	2 of 12	1 of 14	0 of 14	2 of 14	1 of 14	1 of 25	1 of 1	1 of 1	1 of 1	0 of 1	0 of 1	0 of 1
glycolysis V (Pyrococcus) - MouseCyc	↑	9 of 17	2 of 9	0 of 9	0 of 9	2 of 9	1 of 9	0 of 17						
Growth hormone receptor signaling - Reactome	↑↓	3 of 15	2 of 3	2 of 3	2 of 3	3 of 3	3 of 3	9 of 15	3 of 9	5 of 9	2 of 9	3 of 9	0 of 9	0 of 9
GTP hydrolysis and joining of the 60S ribosomal subunit - Reactome	↑	24 of 101	0 of 12	1 of 10	0 of 10	2 of 23	1 of 23	16 of 101	2 of 15	5 of 14	0 of 14	1 of 16	0 of 16	0 of 16
Heme biosynthesis - Reactome	↓	3 of 11	1 of 3	1 of 3	1 of 3	0 of 3	0 of 3	6 of 11	1 of 6	4 of 6	1 of 6	2 of 6	2 of 6	2 of 6
Heme Biosynthesis - Wikipathways	↓	3 of 9	1 of 3	1 of 3	1 of 3	0 of 3	0 of 3	5 of 9	1 of 5	3 of 5	1 of 5	2 of 5	2 of 5	2 of 5
heme biosynthesis II - MouseCyc	↓	3 of 9	1 of 3	1 of 3	1 of 3	0 of 3	0 of 3	5 of 9	1 of 5	3 of 5	1 of 5	2 of 5	2 of 5	2 of 5

IL-5 Signaling Pathway - Wikipathways	↓	7 of 67	2 of 6	0 of 7	0 of 7	0 of 7	0 of 7	21 of 67	1 of 21	9 of 21	2 of 21	4 of 21	2 of 21
IL-6 signaling Pathway - Wikipathways	↓	15 of 98	6 of 15	2 of 15	4 of 15	0 of 15	1 of 15	31 of 98	6 of 31	12 of 31	3 of 31	6 of 31	3 of 31
IL-9 Signaling Pathway - Wikipathways	↓	2 of 24	0 of 2	0 of 2	0 of 2	1 of 2	1 of 2	11 of 24	2 of 11	5 of 11	1 of 11	2 of 11	1 of 11
Immune System - Reactome	↓	94 of 863	30 of 87	9 of 89	13 of 89	8 of 92	5 of 92	202 of 863	27 of 191	77 of 192	12 of 192	42 of 199	27 of 199
Initial triggering of complement - Reactome	↓	2 of 17	1 of 2	0 of 2	0 of 2	0 of 2	0 of 2	8 of 17	1 of 7	2 of 8	0 of 8	5 of 8	4 of 8
Innate Immune System - Reactome	↓	52 of 462	13 of 47	5 of 48	4 of 48	5 of 50	3 of 50	104 of 462	10 of 99	43 of 100	8 of 100	25 of 102	15 of 102
Insulin signaling pathway - Mus musculus (mouse) - KEGG	↑	29 of 142	5 of 27	2 of 28	3 of 28	5 of 29	4 of 29	27 of 142	7 of 25	11 of 27	6 of 27	5 of 27	4 of 27
Interferon Signaling - Reactome	↓	5 of 61	2 of 5	0 of 5	0 of 5	0 of 5	0 of 5	16 of 61	2 of 15	7 of 15	0 of 15	7 of 16	6 of 16
Interleukin-7 signaling - Reactome	↓	0 of 9						6 of 9	0 of 6	2 of 6	0 of 6	0 of 6	0 of 6
ISG15 antiviral mechanism - Reactome	↓	2 of 25	1 of 2	0 of 2	0 of 2	0 of 2	0 of 2	9 of 25	1 of 8	5 of 8	0 of 8	4 of 9	4 of 9
Kennedy pathway - Wikipathways	↓	4 of 14	2 of 4	0 of 4	0 of 4	0 of 4	0 of 4	9 of 14	2 of 9	4 of 9	2 of 9	2 of 9	0 of 9
ketogenesis - MouseCyc	↓	2 of 8	1 of 2	0 of 2	0 of 2	0 of 2	0 of 2	5 of 8	0 of 5	3 of 4	0 of 4	2 of 5	2 of 5
Ketone body metabolism - Reactome	↓	0 of 5						3 of 5	0 of 3	2 of 3	0 of 3	1 of 3	1 of 3
L13a-mediated translational silencing of Ceruloplasmin expression - Reactome	↑	24 of 100	0 of 12	1 of 10	0 of 10	2 of 23	1 of 23	15 of 100	2 of 14	4 of 13	0 of 13	1 of 15	0 of 15
Leptin Insulin Overlap - Wikipathways	↑	4 of 18	1 of 3	1 of 4	1 of 4	3 of 4	2 of 4	4 of 18	1 of 4	1 of 4	2 of 4	1 of 4	0 of 4
Leukocyte transendothelial migration - Mus musculus (mouse) - KEGG	↓	15 of 121	6 of 14	0 of 15	1 of 15	0 of 15	0 of 15	32 of 121	1 of 31	8 of 32	3 of 32	6 of 32	1 of 32
Linoleic acid (LA) metabolism - Reactome	↑	3 of 6	0 of 3	1 of 3	0 of 3	2 of 3	0 of 3	3 of 6	1 of 3	0 of 3	0 of 3	1 of 3	0 of 3
Linoleic acid metabolism - Mus musculus (mouse) - KEGG	↑	13 of 48	1 of 12	1 of 12	1 of 12	1 of 13	1 of 13	14 of 48	1 of 13	9 of 14	1 of 14	6 of 14	5 of 14
Lipid digestion, mobilization, and transport - Reactome	↓	7 of 44	4 of 7	0 of 7	0 of 7	0 of 7	0 of 7	16 of 44	1 of 16	3 of 16	2 of 16	7 of 16	4 of 16
Lipoprotein metabolism - Reactome	↓	4 of 28	2 of 4	0 of 4	0 of 4	0 of 4	0 of 4	11 of 28	1 of 11	3 of 11	2 of 11	5 of 11	3 of 11

L-serine degradation - MouseCyc	↑	2 of 2	1 of 2	0 of 2	1 of 2	1 of 2	1 of 2	0 of 2					
Macrophage markers - Wikipathways	↓	4 of 10	3 of 4	0 of 4	3 of 4	0 of 4	0 of 4	6 of 10	1 of 6	4 of 6	0 of 6	1 of 6	0 of 6
Meiotic Recombination - Reactome	↓	3 of 52	0 of 2	1 of 2	0 of 2	0 of 2	0 of 2	17 of 52	6 of 10	0 of 2	0 of 2	1 of 13	1 of 13
Membrane Trafficking - Reactome	↓	20 of 146	4 of 18	1 of 17	0 of 17	0 of 19	0 of 19	41 of 146	4 of 36	11 of 41	1 of 41	5 of 41	3 of 41
Metabolism - Reactome	↑↓	369 of 1346	86 of 342	27 of 352	24 of 352	27 of 366	27 of 366	304 of 1346	66 of 282	107 of 298	31 of 298	86 of 302	53 of 302
Metabolism of amino acids and derivatives	↑	50 of 142	5 of 46	2 of 49	2 of 49	1 of 50	4 of 50	30 of 142	8 of 29	12 of 40	1 of 40	6 of 40	4 of 40
Metabolism of carbohydrates - Reactome	↑	62 of 243	10 of 59	5 of 59	3 of 59	4 of 62	3 of 62	42 of 243	8 of 41	14 of 41	6 of 41	8 of 42	7 of 42
Metabolism of lipids and lipoproteins - Reactome	↑↓	114 of 483	38 of 107	13 of 111	7 of 111	10 of 113	10 of 113	135 of 483	31 of 124	47 of 133	13 of 133	39 of 135	21 of 135
Metabolism of mRNA - Reactome	↑	33 of 158	1 of 17	3 of 20	2 of 20	3 of 31	1 of 31	34 of 158	6 of 28	13 of 32	5 of 32	2 of 33	3 of 33
Metabolism of nucleotides - Reactome	↑	21 of 80	5 of 20	3 of 21	2 of 21	1 of 21	1 of 21	20 of 80	4 of 20	5 of 20	1 of 20	4 of 20	2 of 20
Metabolism of porphyrins - Reactome	↓	5 of 15	3 of 5	1 of 5	1 of 5	0 of 5	0 of 5	6 of 15	1 of 6	4 of 6	1 of 6	2 of 6	2 of 6
Metabolism of proteins - Reactome	↑	81 of 458	10 of 63	4 of 65	4 of 65	5 of 80	2 of 80	97 of 458	19 of 91	30 of 94	4 of 94	15 of 97	7 of 97
Metabolism of vitamins and cofactors	↑	21 of 73	5 of 21	2 of 21	1 of 21	0 of 21	0 of 21	18 of 73	4 of 14	8 of 16	1 of 16	6 of 16	4 of 16
Metabolism of water-soluble vitamins and cofactors - Reactome	↑	21 of 73	5 of 21	2 of 21	1 of 21	0 of 21	0 of 21	18 of 73	4 of 14	8 of 16	1 of 16	6 of 16	4 of 16
Metabolism of xenobiotics by cytochrome P450 - Mus musculus (mouse) - KEGG	↑↓	36 of 88	10 of 31	1 of 28	3 of 28	7 of 33	7 of 33	25 of 88	5 of 23	11 of 24	3 of 24	14 of 24	13 of 24
Metal ion SLC transporters - Reactome	↓	2 of 25	0 of 2	1 of 2	0 of 2	0 of 2	0 of 2	11 of 25	1 of 11	3 of 11	1 of 11	0 of 11	1 of 11
Metapathway biotransformation - Wikipathways	↑	39 of 137	17 of 36	4 of 35	4 of 35	7 of 36	5 of 36	29 of 137	7 of 27	7 of 29	3 of 29	8 of 29	6 of 29
Methylation - Wikipathways	↑	4 of 9	1 of 4	1 of 4	1 of 4	2 of 4	1 of 4	3 of 9	0 of 3	1 of 3	0 of 3	0 of 3	0 of 3
mevalonate pathway I - MouseCyc	↑↓	6 of 11	5 of 6	2 of 6	1 of 6	0 of 6	0 of 6	6 of 11	1 of 6	3 of 5	0 of 5	2 of 5	3 of 6
Mitochondrial Fatty Acid Beta-Oxidation - Reactome	↓	2 of 15	0 of 2	5 of 15	0 of 5	3 of 5	0 of 5	1 of 5	1 of 5				

mitochondrial fatty acid beta-oxidation of unsaturated fatty acids - Reactome	↓	0 of 3					3 of 3	0 of 3	2 of 3	0 of 3	0 of 3	0 of 3
Mitochondrial LC-Fatty Acid Beta-Oxidation - Wikipathways	↓	4 of 16	0 of 4	0 of 4	0 of 4	1 of 4	1 of 4	9 of 16	1 of 9	4 of 9	0 of 9	0 of 9
Mitochondrial Protein Import - Reactome	↑	13 of 43	2 of 13	0 of 13	0 of 13	0 of 13	0 of 13	9 of 43	1 of 9	4 of 9	0 of 9	4 of 9
mRNA 3,-end processing - Reactome	↓	4 of 34	0 of 2	0 of 4	1 of 4	0 of 4	0 of 4	15 of 34	1 of 15	7 of 14	0 of 14	2 of 15
mRNA Editing - Reactome	↓	1 of 4	0 of 1	3 of 4	1 of 3	2 of 3	0 of 3	0 of 3				
mRNA Editing: C to U Conversion - Reactome	↓	0 of 2						2 of 2	1 of 2	2 of 2	0 of 2	0 of 2
mRNA Processing - Reactome	↓	25 of 157	3 of 22	0 of 23	1 of 23	0 of 25	0 of 25	51 of 157	5 of 50	19 of 49	0 of 49	6 of 50
mRNA processing - Wikipathways	↓	71 of 447	8 of 61	3 of 66	2 of 66	3 of 71	0 of 71	112 of 447	8 of 108	50 of 110	5 of 150	14 of 112
mRNA Splicing - Major Pathway - Reactome	↓	19 of 107	3 of 16	0 of 17	1 of 17	0 of 19	0 of 19	38 of 107	4 of 37	13 of 36	0 of 36	5 of 37
mRNA Splicing - Minor Pathway - Reactome	↓	7 of 42	2 of 5	0 of 5	0 of 5	0 of 7	0 of 7	15 of 42	3 of 14	5 of 14	0 of 14	2 of 14
mRNA Splicing - Reactome	↓	19 of 107	3 of 16	0 of 17	1 of 17	0 of 19	0 of 19	38 of 107	4 of 37	13 of 36	0 of 36	5 of 37
mRNA surveillance pathway - Mus musculus (mouse) - KEGG	↓	10 of 92	1 of 7	0 of 10	0 of 10	0 of 10	0 of 10	28 of 92	3 of 27	13 of 27	0 of 27	3 of 28
MyD88-independent cascade - Reactome	↓	12 of 88	4 of 11	2 of 10	2 of 10	1 of 11	0 of 11	28 of 88	2 of 26	16 of 28	4 of 28	1 of 28
NAD biosynthesis II (from tryptophan) - MouseCyc	↑	5 of 10	1 of 5	0 of 5	0 of 5	1 of 5	1 of 5	1 of 10	0 of 1	0 of 1	0 of 1	0 of 1
Nicotinate and nicotinamide metabolism - Mus musculus (mouse) - Kegg	↑	10 of 31	4 of 10	2 of 10	1 of 10	1 of 10	0 of 10	5 of 31	2 of 5	4 of 5	0 of 5	1 of 5
nicotine degradation II - MouseCyc	↑↓	20 of 50	6 of 17	1 of 17	3 of 17	4 of 19	5 of 19	23 of 50	5 of 20	12 of 23	1 of 23	13 of 23
nicotine degradation III - MouseCyc	↑↓	15 of 43	4 of 12	1 of 12	3 of 12	2 of 14	3 of 14	22 of 43	5 of 19	12 of 22	1 of 22	13 of 22
Nonsense Mediated Decay Enhanced by the Exon Junction Complex - Reactome	↑	24 of 94	0 of 11	1 of 11	0 of 11	2 of 23	1 of 23	13 of 94	2 of 12	4 of 11	0 of 11	2 of 13
Nonsense Mediated Decay Independent of the Exon Junction Complex - Reactome	↑	23 of 84	0 of 11	1 of 10	0 of 10	2 of 22	1 of 22	10 of 84	2 of 12	3 of 8	0 of 8	1 of 10
Nonsense-Mediated Decay - Reactome	↑	24 of 94	0 of 11	1 of 11	0 of 11	2 of 23	1 of 23	13 of 94	2 of 12	4 of 11	0 of 11	2 of 13

Nuclear Receptor transcription pathway - Reactome	↓	9 of 49	2 of 9	0 of 9	0 of 9	0 of 9	1 of 9	17 of 49	2 of 14	7 of 17	3 of 17	0 of 17	0 of 17
Nuclear Receptors - Wikipathways	↓	7 of 38	1 of 7	0 of 7	0 of 7	0 of 7	1 of 7	14 of 38	2 of 13	5 of 14	2 of 14	0 of 14	0 of 14
Nuclear receptors in lipid metabolism and toxicity	↑↓	13 of 30	5 of 12	1 of 13	3 of 13	0 of 13	1 of 13	13 of 30	1 of 12	6 of 13	3 of 13	1 of 13	0 of 13
oleate biosynthesis II (animals) - MouseCyc	↑	3 of 7	0 of 3	2 of 3	0 of 3	2 of 3	0 of 3	2 of 7	0 of 2	0 of 2	1 of 2	2 of 2	2 of 2
One carbon metabolism and related pathways - Wikipathways	↓	10 of 49	4 of 10	1 of 10	0 of 10	0 of 10	0 of 10	20 of 49	7 of 18	8 of 20	1 of 20	6 of 20	2 of 20
Oxidative phosphorylation - <i>Mus musculus</i> (mouse) - KEGG	↑	39 of 124	0 of 33	0 of 36	0 of 36	1 of 39	0 of 39	7 of 124	1 of 6	3 of 6	0 of 6	1 of 6	0 of 6
Oxidative phosphorylation - Wikipathways	↑	18 of 52	1 of 14	0 of 17	0 of 17	1 of 18	0 of 18	1 of 52	0 of 0	0 of 0	0 of 0	0 of 0	0 of 0
Oxidative Stress - Wikipathways	↑↓	9 of 27	5 of 9	2 of 9	1 of 9	0 of 9	0 of 9	11 of 27	3 of 11	3 of 11	2 of 11	4 of 11	2 of 11
Pentose and glucuronate interconversions - <i>Mus musculus</i> (mouse) - KEGG	↑	11 of 29	7 of 10	0 of 7	1 of 7	0 of 8	0 of 8	10 of 29	4 of 9	4 of 10	2 of 10	3 of 10	2 of 10
pentose phosphate pathway - MouseCyc	↑	6 of 11	0 of 6	1 of 5	0 of 5	0 of 6	0 of 6	1 of 11	0 of 1	0 of 1	0 of 1	0 of 1	0 of 1
Pentose phosphate pathway - <i>Mus musculus</i> (mouse) - KEGG	↑	13 of 30	1 of 13	1 of 12	0 of 12	0 of 13	0 of 13	1 of 30	0 of 1	0 of 1	0 of 1	0 of 1	0 of 1
Pentose Phosphate Pathway - Wikipathways	↑	6 of 7	0 of 6	1 of 5	0 of 5	0 of 6	0 of 6	0 of 7					
Pentose phosphate pathway (hexose monophosphate shunt) - Reactome	↑	6 of 8	0 of 6	1 of 5	0 of 5	0 of 6	0 of 6	0 of 8					
pentose phosphate pathway (non-oxidative branch) - MouseCyc	↑	4 of 6	0 of 4	0 of 3	0 of 3	0 of 4	0 of 4	0 of 6					
pentose phosphate pathway (oxidative branch) - MOuseCyc	↑	2 of 5	0 of 2	1 of 2	0 of 2	0 of 2	0 of 2	1 of 5	0 of 1	0 of 1	0 of 1	0 of 1	0 of 1
Peptide chain elongation - Reactome	↑	23 of 79	1 of 11	1 of 10	0 of 10	3 of 22	1 of 22	5 of 79	0 of 4	0 of 3	0 of 3	1 of 5	0 of 5
Peroxisomal lipid metabolism - Reactome	↓	2 of 22	2 of 2	0 of 2	0 of 2	0 of 2	0 of 2	21 of 22	2 of 12	2 of 12	1 of 12	4 of 12	2 of 12
Peroxisome - <i>Mus musculus</i> (mouse) - KEGG	↑↓	21 of 80	7 of 19	1 of 21	0 of 21	1 of 21	1 of 21	32 of 80	7 of 32	9 of 32	2 of 32	7 of 32	4 of 32
Phagosome - <i>Mus musculus</i> (mouse)	↑	37 of 169	9 of 32	2 of 34	2 of 34	0 of 37	0 of 37	39 of 169	2 of 37	14 of 35	1 of 35	11 of 39	4 of 39
Phase 1 - Functionalization of compounds - Reactome	↑↓	37 of 96	11 of 30	4 of 29	6 of 29	8 of 36	8 of 36	31 of 96	6 of 26	12 of 29	5 of 25	16 of 31	11 of 31

Phase II conjugation - Reactome	↑	26 of 63	11 of 25	1 of 23	3 of 23	4 of 24	3 of 24	19 of 63	4 of 18	4 of 18	2 of 18	6 of 18	4 of 18
Phenylalanine and tyrosine catabolism - Reactome	↑	5 of 9	1 of 4	0 of 4	0 of 4	0 of 4	1 of 4	0 of 9					
Phenylalanine, tyrosine and tryptophan biosynthesis - Mus musculus (mouse) - KEGG	↑	4 of 7	2 of 4	2 of 4	2 of 4	0 of 4	2 of 4	1 of 7	0 of 1	0 of 1	0 of 1	0 of 1	0 of 1
phosphatidylcholine biosynthesis I - MouseCyc	↓	0 of 6						5 of 6	1 of 5	3 of 5	1 of 5	1 of 5	0 of 5
phosphatidylethanolamine biosynthesis II - MouseCyc	↓	3 of 8	1 of 2	0 of 3	0 of 3	0 of 3	0 of 3	5 of 8	2 of 4	2 of 5	1 of 5	1 of 5	0 of 5
Phospholipid metabolism - Reactome	↑	27 of 131	6 of 24	1 of 27	2 of 27	0 of 27	2 of 27	30 of 131	8 of 29	7 of 29	2 of 29	5 of 30	2 of 30
Platelet degranulation - Reactome	↑	24 of 85	12 of 22	1 of 21	1 of 21	1 of 24	0 of 24	20 of 85	2 of 19	5 of 19	0 of 19	7 of 20	5 of 20
Porphyrin and chlorophyll metabolism - Mus musculus (mouse) - KEGG	↓	9 of 37	6 of 6	1 of 6	1 of 6	0 of 6	0 of 6	14 of 37	5 of 14	8 of 14	2 of 14	4 of 14	4 of 14
Post-Elongation Processing of Intron-Containing pre-mRNA - Reactome	↓	4 of 34	0 of 4	0 of 4	1 of 4	0 of 4	0 of 4	15 of 34	1 of 15	7 of 14	0 of 14	2 of 15	2 of 15
Post-Elongation Processing of the Transcript - Reactome	↓	4 of 43	0 of 6	0 of 6	1 of 6	0 of 6	0 of 6	16 of 43	1 of 16	7 of 15	0 of 15	3 of 16	2 of 16
Post-translational protein modification - Reactome	↓	16 of 164	3 of 14	0 of 15	1 of 15	0 of 16	0 of 16	41 of 164	9 of 39	11 of 41	0 of 41	5 of 41	5 of 41
PPAR signaling pathway - Mus musculus (mouse) - KEGG	↑↓	20 of 78	8 of 17	4 of 19	3 of 19	6 of 19	4 of 19	32 of 78	5 of 30	13 of 31	4 of 31	7 of 32	5 of 32
PPAR signaling pathway - Wikipathways	↑↓	20 of 78	8 of 17	4 of 19	3 of 19	6 of 19	4 of 19	32 of 78	5 of 30	13 of 31	4 of 31	7 of 32	5 of 32
Primary bile acid biosynthesis - Mus musculus (mouse) - KEGG	↓	2 of 16	1 of 2	1 of 2	1 of 2	0 of 2	0 of 2	12 of 16	2 of 12	2 of 12	1 of 12	5 of 12	2 of 12
Processing of Capped Intron-Containing Pre-mRNA - Reactome	↓	22 of 138	3 of 19	0 of 20	1 of 20	0 of 22	0 of 22	45 of 138	4 of 44	15 of 43	0 of 43	6 of 44	2 of 44
Proline catabolism - Reactome	↑	2 of 2	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2					
Propanoate metabolism - Mus musculus (mouse) - KEGG	↑↓	9 of 31	3 of 9	2 of 9	0 of 9	1 of 9	1 of 9	9 of 31	0 of 8	6 of 8	0 of 8	3 of 9	3 of 9
Proteasome - Mus musculus (mouse) - KEGG	↓	3 of 44	1 of 3	0 of 3	0 of 3	0 of 3	0 of 3	17 of 44	3 of 16	6 of 17	0 of 17	5 of 17	5 of 17
Proteasome Degradation - Wikipathways	↓	9 of 57	2 of 9	1 of 9	0 of 9	0 of 8	0 of 8	19 of 57	3 of 18	6 of 19	1 of 19	5 of 19	5 of 19
Protein processing in endoplasmic reticulum - Mus musculus (mouse) - KEGG	↓	20 of 165	0 of 16	1 of 19	2 of 19	0 of 19	1 of 19	67 of 165	17 of 63	24 of 67	6 of 67	14 of 66	10 of 66

14

purine and pyrimidine metabolism - MouseCyc	↑	9 of 25	3 of 9	2 of 8	1 of 8	0 of 9	0 of 9	8 of 25	1 of 8	5 of 8	1 of 8	1 of 8	1 of 8
Purine metabolism - <i>Mus musculus</i> (mouse) - KEGG	↑	39 of 170	9 of 37	1 of 38	0 of 38	2 of 39	0 of 39	21 of 170	6 of 20	5 of 20	1 of 20	1 of 20	2 of 20
Purine metabolism - Wikipathways	↑	37 of 165	9 of 35	1 of 36	0 of 36	2 of 37	0 of 37	20 of 165	5 of 19	5 of 19	1 of 19	1 of 19	2 of 19
Pyrimidine salvage reactions - Reactome	↑	5 of 9	2 of 5	2 of 5	2 of 5	1 of 50	1 of 5	3 of 9	1 of 3	1 of 3	1 of 3	0 of 3	0 of 3
Pyruvate metabolism - <i>Mus musculus</i> (mouse) - KEGG	↑	16 of 42	6 of 15	2 of 16	0 of 16	1 of 16	0 of 16	10 of 42	1 of 10	6 of 10	0 of 10	2 of 11	2 of 11
Pyruvate metabolism - Reactome	↑	7 of 20	1 of 7	0 of 7	1 of 7	0 of 7	1 of 7	6 of 20	1 of 6	1 of 6	2 of 6	4 of 6	1 of 6
Pyruvate metabolism and Citric Acid (TCA) cycle - Reactome	↑	20 of 42	3 of 20	0 of 20	1 of 20	0 of 20	1 of 20	6 of 42	1 of 6	1 of 6	2 of 6	4 of 6	1 of 6
Regulation of Complement cascade - Reactome	↓	3 of 23	2 of 3	0 of 3	0 of 3	0 of 3	0 of 3	11 of 23	0 of 10	3 of 10	0 of 10	6 of 11	2 of 11
Regulation of pyruvate dehydrogenase (PDH) complex - Reactome	↑	6 of 12	1 of 6	0 of 6	1 of 6	0 of 6	1 of 6	2 of 12	1 of 2	1 of 2	2 of 2	0 of 2	0 of 2
Removal of aminoterminal propeptides from gamma-carboxylated proteins - Reactome	↑	4 of 9	1 of 4	0 of 4	0 of 4	0 of 4	0 of 4	3 of 9	0 of 3	1 of 3	0 of 3	1 of 3	0 of 3
Respiratory electron transport - Reactome	↑	23 of 68	0 of 21	0 of 21	0 of 21	1 of 23	0 of 23	2 of 68	0 of 1	1 of 1	0 of 1	1 of 1	0 of 1
Respiratory electron transport, ATP synthesis by chemiosmotic coupling, and heat production by uncoupling proteins - Reactome	↑	29 of 84	0 of 25	0 of 27	0 of 27	1 of 29	0 of 29	2 of 84	0 of 1	1 of 1	0 of 1	1 of 1	0 of 1
Response to elevated platelet cytosolic Ca2+	↑	24 of 90	12 of 22	1 of 21	1 of 21	1 of 24	0 of 24	21 of 90	2 of 20	6 of 20	0 of 20	7 of 21	5 of 21
Retinoid metabolism and transport - Reactome	↓	9 of 46	3 of 9	0 of 9	0 of 9	0 of 9	0 of 9	13 of 46	4 of 13	5 of 13	3 of 13	6 of 13	3 of 13
Retinol metabolism - <i>Mus musculus</i> (mouse) - KEGG	↑↓	29 of 77	9 of 25	2 of 22	5 of 22	6 of 26	6 of 26	31 of 77	9 of 29	16 of 29	4 of 29	15 of 31	12 of 31
RIG-I/MDA5 mediated induction of IFN-alpha/beta pathways - Reactome	↓	7 of 45	3 of 7	1 of 7	1 of 7	1 of 7	0 of 7	16 of 45	1 of 16	8 of 16	1 of 16	3 of 16	2 of 16
RNA Polymerase I Promoter Opening - Reactome	↓	1 of 21	0 of 0	0 of 0	0 of 0	0 of 0	0 of 0	6 of 21	1 of 3	1 of 1	0 of 1	0 of 4	0 of 4
RNA Polymerase II Transcription - Reactome	↓	14 of 101	1 of 12	0 of 13	1 of 13	0 of 14	0 of 14	30 of 101	5 of 28	13 of 28	0 of 28	3 of 29	2 of 29
RNA Polymerase II Transcription Termination - Reactome	↓	4 of 43	0 of 6	0 of 6	1 of 6	0 of 6	0 of 6	16 of 43	1 of 16	7 of 15	0 of 15	3 of 16	2 of 16
RNA transport - <i>Mus musculus</i> (mouse) - KEGG	↓	12 of 161	1 of 11	0 of 7	0 of 7	0 of 12	0 of 12	46 of 161	4 of 45	20 of 45	0 of 45	4 of 46	1 of 46

salvage pathways of pyrimidine ribonucleotides - MouseCyc	↑	7 of 15	3 of 7	4 of 7	1 of 7	1 of 7	1 of 7	3 of 15	1 of 3	1 of 3	1 of 3	0 of 3	0 of 3
Scavenging by Class B Receptors - Reactome	↓	2 of 6	1 of 2	1 of 2	1 of 2	0 of 2	0 of 2	4 of 6	0 of 4	1 of 4	1 of 4	1 of 4	0 of 4
Selenium - Wikipathways	↑	8 of 22	2 of 8	2 of 8	0 of 8	2 of 8	1 of 8	3 of 22	0 of 2	0 of 3	0 of 3	1 of 3	0 of 3
Selenium metabolism-Selenoproteins - Wikipathways	↓	8 of 47	3 of 7	1 of 7	0 of 7	0 of 8	0 of 8	18 of 47	4 of 16	4 of 17	3 of 17	3 of 18	3 of 18
Senescence and Autophagy - Wikipathways	↓	15 of 98	4 of 15	1 of 15	3 of 15	2 of 15	0 of 15	30 of 98	3 of 29	6 of 30	1 of 30	5 of 30	2 of 30
Serine biosynthesis - Reactome	↑	3 of 3	0 of 3	0 of 3	0 of 3	0 of 3	0 of 3	0 of 3					
Signaling by Interleukins - Reactome	↓	11 of 104	4 of 11	2 of 11	3 of 11	0 of 11	0 of 11	32 of 104	3 of 31	12 of 32	0 of 32	5 of 32	0 of 32
Spliceosome - Mus musculus (mouse) - KEGG	↓	20 of 129	3 of 14	1 of 17	2 of 17	0 of 19	0 of 19	43 of 129	3 of 38	16 of 42	3 of 42	4 of 42	3 of 42
SREBF and miR33 in cholesterol and lipid homeostasis - Wikipathways	↑	7 of 13	3 of 7	0 of 7	1 of 7	0 of 7	0 of 7	6 of 13	0 of 5	4 of 5	1 of 5	0 of 5	0 of 5
SRP-dependent cotranslational protein targeting to membrane - Reactome	↑	28 of 103	0 of 13	1 of 14	0 of 14	2 of 27	1 of 27	10 of 103	2 of 8	1 of 8	0 of 8	2 of 10	0 of 10
Starch and sucrose metabolism - Mus musculus (mouse) - KEGG	↑	15 of 46	5 of 13	0 of 11	2 of 11	1 of 12	1 of 12	13 of 46	5 of 13	6 of 13	3 of 13	5 of 13	5 of 13
Statin Pathway - Wikipathways	↓	5 of 19	2 of 5	1 of 5	1 of 5	0 of 5	0 of 5	9 of 19	1 of 9	3 of 9	2 of 9	3 of 9	2 of 9
Steroid biosynthesis - Mus musculus (mouse) - KEGG	↑	11 of 17	7 of 11	2 of 11	0 of 11	1 of 11	1 of 11	4 of 17	1 of 4	1 of 4	1 of 4	2 of 4	1 of 4
Steroid Biosynthesis - Wikipathways	↓	2 of 13	2 of 2	0 of 2	0 of 2	1 of 2	1 of 2	5 of 13	2 of 5	3 of 5	1 of 5	3 of 5	1 of 5
Steroid hormone biosynthesis - Mus musculus (mouse) - KEGG	↑↓	14 of 52	7 of 12	1 of 11	3 of 11	3 of 11	2 of 11	17 of 52	9 of 16	9 of 17	3 of 17	8 of 17	5 of 17
sucrose degradation V (mammalian) - MouseCyc	↑	4 of 7	1 of 4	0 of 4	0 of 4	0 of 4	0 of 4	0 of 7					
Sulfide oxidation to sulfate - Reactome	↓	2 of 5	0 of 2	0 of 2	0 of 2	0 of 2	0 of 2	3 of 5	1 of 3	1 of 3	0 of 3	1 of 3	1 of 3
Sulfur amino acid metabolism - Reactome	↓	7 of 25	1 of 7	1 of 7	1 of 7	0 of 7	1 of 7	10 of 25	6 of 10	5 of 10	0 of 10	3 of 10	1 of 10
superpathway of acetyl-CoA biosynthesis - MouseCyc	↑	4 of 6	0 of 4	0 of 4	0 of 4	1 of 4	1 of 4	0 of 6					
superpathway of cholesterol biosynthesis - MouseCyc	↑↓	18 of 25	14 of 18	6 of 18	2 of 18	1 of 18	1 of 18	10 of 25	1 of 10	4 of 9	1 of 9	5 of 10	3 of 10

superpathway of citrulline metabolism - MouseCyc	↑	7 of 17	0 of 5	0 of 7	0 of 7	0 of 7	1 of 7	0 of 17					
Synthesis and degradation of ketone bodies - Mus musculus (mouse)	↓	2 of 10	1 of 2	0 of 2	0 of 2	0 of 2	0 of 2	5 of 10	0 of 5	3 of 4	0 of 4	2 of 5	2 of 5
Synthesis and Degradation of Ketone Bodies - Wikipathways	↓	0 of 5						3 of 5	0 of 3	2 of 3	0 of 3	1 of 3	1 of 3
Synthesis and interconversion of nucleotide di- and triphosphates - Reactome	↑	7 of 18	2 of 7	1 of 7	0 of 7	0 of 7	0 of 7	4 of 18	1 of 4	1 of 4	0 of 4	2 of 4	1 of 4
Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) - Reactome	↑↓	10 of 24	2 of 8	2 of 7	1 of 7	4 of 9	3 of 9	10 of 24	2 of 8	4 of 9	1 of 9	4 of 10	4 of 10
Synthesis of bile acids and bile salts - Reactome	↓	6 of 27	1 of 6	1 of 6	1 of 6	0 of 6	0 of 6	15 of 27	3 of 15	4 of 15	1 of 15	6 of 15	2 of 15
Synthesis of bile acids and bile salts via 24-hydroxycholesterol - Reactome	↓	4 of 18	0 of 4	8 of 18	1 of 8	3 of 8	0 of 8	3 of 8	2 of 8				
Synthesis of bile acids and bile salts via 27-hydroxycholesterol - Reactome	↓	3 of 14	0 of 3	6 of 14	2 of 6	3 of 6	0 of 6	3 of 6	1 of 6				
Synthesis of bile acids and bile salts via 7alpha-hydroxycholesterol - Reactome	↓	5 of 23	1 of 5	1 of 5	1 of 5	0 of 5	0 of 5	13 of 23	2 of 13	4 of 13	1 of 13	5 of 13	2 of 13
Synthesis of epoxy (EET) and dihydroxyeicosatrienoic acids (DHET) - Reactome	↑↓	7 of 21	0 of 6	1 of 5	0 of 5	1 of 6	1 of 6	8 of 21	1 of 8	6 of 8	1 of 8	3 of 8	2 of 8
Synthesis of GDP-mannose - Reactome	↓	0 of 5						2 of 5	1 of 2	0 of 2	0 of 2	0 of 2	0 of 2
Synthesis of Ketone Bodies - Reactome	↓	0 of 4						3 of 4	0 of 3	2 of 3	0 of 3	1 of 3	1 of 3
Synthesis of PC - Reactome	↑	5 of 18	1 of 5	0 of 5	1 of 5	0 of 5	0 of 5	6 of 18	1 of 6	3 of 5	1 of 5	1 of 6	0 of 6
Synthesis of PE - Reactome	↑↓	5 of 12	2 of 5	0 of 5	1 of 5	0 of 5	0 of 5	7 of 12	2 of 6	2 of 6	1 of 6	1 of 7	0 of 7
Synthesis of substrates in N-glycan biosynthesis - Reactome	↓	2 of 17	0 of 2	0 of 2	1 of 2	0 of 2	0 of 2	7 of 17	2 of 7	2 of 7	0 of 7	0 of 7	1 of 7
Synthesis of very long-chain fatty acyl-CoAs - Reactome	↑	5 of 13	0 of 4	0 of 5	0 of 5	1 of 5	1 of 5	4 of 13	2 of 4	2 of 4	0 of 4	2 of 4	1 of 4
Targeted protein degradation - MouseCyc	↓	3 of 59	1 of 3	0 of 3	0 of 3	0 of 3	0 of 3	23 of 59	3 of 21	8 of 23	0 of 23	6 of 23	5 of 23
TCA cycle - MouseCyc	↑	9 of 13	1 of 9	0 of 9									
TCA Cycle - Wikipathways	↑	18 of 31	3 of 18	0 of 18	1 of 18	0 of 18	1 of 18	2 of 31	1 of 2	2 of 2	1 of 2	0 of 2	0 of 2
TCA cycle variation III (eukaryotic) - MouseCyc	↑	10 of 16	2 of 10	0 of 16									

Terminal pathway of complement - Reactome	↓	0 of 7					6 of 7	0 of 6	3 of 6	0 of 6	3 of 6	1 of 6	
Terpenoid backbone biosynthesis - Mus musculus (mouse) - KEGG	↑	8 of 21	6 of 7	3 of 8	1 of 8	0 of 8	0 of 8	6 of 21	1 of 6	2 of 5	0 of 5	2 of 6	3 of 6
tetrapyrrole biosynthesis II - MouseCyc	↓	1 of 5	1 of 1	1 of 1	1 of 1	0 of 1	0 of 1	4 of 5	1 of 4	3 of 4	1 of 4	2 of 4	2 of 4
TGF-beta Receptor Signaling Pathway - Wikipathways	↓	19 of 150	3 of 18	4 of 19	1 of 19	0 of 19	0 of 19	42 of 150	4 of 39	15 of 42	5 of 42	3 of 42	1 of 42
The citric acid (TCA) cycle and respiratory electron transport - Reactome	↑	46 of 122	3 of 42	0 of 44	1 of 44	1 of 46	1 of 46	8 of 122	1 of 7	2 of 7	2 of 7	5 of 7	1 of 7
TNF-alpha NF-kB Signaling Pathway - Wikipathways	↓	24 of 181	7 of 21	4 of 23	3 of 23	0 of 24	1 of 24	51 of 181	11 of 47	18 of 51	3 of 51	4 of 51	5 of 51
Toll Like Receptor 3 (TLR3) Cascade - Reactome	↓	12 of 88	4 of 11	2 of 10	2 of 10	1 of 11	0 of 11	28 of 88	2 of 26	16 of 28	4 of 28	1 of 28	0 of 28
Toll Like Receptor 4 (TLR4) Cascade - Reactome	↓	4 of 14	2 of 13	2 of 13	2 of 14	1 of 14	0 of 14	32 of 105	2 of 30	17 of 32	4 of 32	3 of 32	1 of 32
Toll Like Receptor signaling - Wikipathways	↓	4 of 12	1 of 4	0 of 4	0 of 4	1 of 4	2 of 4	12 of 33	2 of 12	8 of 12	0 of 12	1 of 12	0 of 12
Toll-like receptor signaling pathway - Wikipathways	↓	12 of 96	4 of 12	0 of 12	2 of 12	2 of 12	1 of 12	27 of 96	3 of 27	14 of 27	2 of 27	5 of 27	3 of 27
Toll-Like Receptors Cascades - Reactome	↓	18 of 125	5 of 17	2 of 16	2 of 16	2 of 17	1 of 17	38 of 125	3 of 36	22 of 38	5 of 38	6 of 38	3 of 38
Trafficking of dietary sterols - Reactome	↑	2 of 2	2 of 2	0 of 2									
trans, trans-farnesyl diphosphate biosynthesis - MouseCyc	↑	2 of 5	2 of 2	2 of 2	1 of 2	0 of 2	0 of 2	1 of 5	0 of 1	1 of 1	0 of 1	0 of 1	0 of 1
Transcription - Reactome	↓	19 of 173	1 of 17	0 of 17	1 of 17	0 of 18	0 of 18	48 of 173	9 of 43	16 of 41	1 of 41	5 of 45	3 of 45
trans-Golgi Network Vesicle Budding - Reactome	↓	6 of 60	2 of 6	2 of 6	0 of 6	0 of 6	0 of 6	19 of 60	2 of 16	7 of 19	1 of 19	3 of 19	1 of 19
Translation - Reactome	↑	34 of 141	1 of 19	1 of 19	0 of 19	3 of 33	1 of 33	26 of 141	4 of 24	8 of 24	0 of 24	3 of 26	0 of 26
Translation Factors - Wikipathways	↓	8 of 49	2 of 5	0 of 5	0 of 5	1 of 8	1 of 8	17 of 49	2 of 17	8 of 17	0 of 17	3 of 17	0 of 17
Transport of Mature mRNA derived from an Intron-Containing Transcript - Reactome	↓	5 of 51	0 of 5	0 of 5	1 of 5	0 of 5	0 of 5	18 of 51	1 of 18	6 of 17	0 of 17	2 of 18	2 of 18
Transport of Mature Transcript to Cytoplasm - Reactome	↓	5 of 51	0 of 5	0 of 5	1 of 5	0 of 5	0 of 5	19 of 55	1 of 19	7 of 18	0 of 18	2 of 19	2 of 19
Triacylglyceride Synthesis - Wikipathways	↓	4 of 23	2 of 4	0 of 4	0 of 4	0 of 4	0 of 4	9 of 23	3 of 9	3 of 9	2 of 9	2 of 9	1 of 9

↑ ↓

TRIF-mediated TLR3/TLR4 signaling - Reactome	↓	12 of 88	4 of 11	2 of 10	2 of 10	1 of 11	0 of 11	28 of 88	2 of 26	16 of 28	4 of 28	1 of 28	0 of 28
Triglyceride Biosynthesis - Reactome	↑	17 of 36	1 of 14	0 of 17	1 of 17	2 of 17	3 of 17	9 of 36	5 of 9	3 of 9	1 of 9	3 of 9	1 of 9
Tryptophan metabolism - Mus musculus (mouse) - KEGG	↓	12 of 44	1 of 12	1 of 11	1 of 11	3 of 12	3 of 12	12 of 44	3 of 11	7 of 11	0 of 11	3 of 12	3 of 12
Tryptophan metabolism - Wikipathways	↓	9 of 43	1 of 8	0 of 8	0 of 8	2 of 9	2 of 9	15 of 43	5 of 14	7 of 15	1 of 15	4 of 15	3 of 15
Type II interferon signaling (IFNG) - Wikipathways	↓	4 of 33	4 of 4	0 of 4	0 of 4	0 of 4	0 of 4	17 of 33	2 of 16	8 of 16	0 of 16	6 of 17	3 of 17
tyrosine degradation I - MouseCyc	↑	3 of 7	1 of 3	0 of 3	0 of 3	0 of 3	1 of 3	1 of 7	0 of 1	1 of 1	1 of 1	0 of 1	0 of 1
Tyrosine metabolism - Mus musculus (mouse) - KEGG	↓	6 of 42	2 of 6	2 of 6	2 of 6	0 of 6	2 of 6	8 of 42	3 of 8	5 of 8	1 of 8	2 of 8	2 of 8
Ubiquinone and other terpenoid-quinone biosynthesis - Mus musculus (mouse) - KEGG	↑	4 of 10	1 of 4	0 of 4	0 of 4	0 of 4	1 of 4	2 of 10	0 of 2	1 of 2	0 of 2	0 of 2	1 of 2
Ubiquitin mediated proteolysis - Mus musculus (mouse) - KEGG	↓	9 of 136	0 of 9	0 of 9	0 of 9	1 of 9	0 of 9	44 of 136	5 of 42	15 of 43	2 of 43	9 of 44	4 of 44
urea cycle - MouseCyc	↑	4 of 6	0 of 3	0 of 4	0 of 4	0 of 4	1 of 4	0 of 6					
Urea cycle - Reactome	↑	5 of 9	0 of 4	0 of 5	0 of 5	0 of 5	1 of 5	0 of 9					
Urea cycle and metabolism of amino groups - Wikipathways	↑	8 of 20	0 of 6	0 of 8	0 of 8	0 of 8	1 of 8	3 of 20	0 of 2	0 of 3	0 of 3	0 of 3	0 of 3
Valine, leucine and isoleucine degradation - Mus musculus (mouse) - KEGG	↓	11 of 51	3 of 11	1 of 11	1 of 11	1 of 11	1 of 11	17 of 51	3 of 16	11 of 16	0 of 16	5 of 17	5 of 17
Xenobiotics - Reactome	↑↓	18 of 37	4 of 13	1 of 11	3 of 11	3 of 17	4 of 17	15 of 37	2 of 12	9 of 14	3 of 14	9 of 15	8 of 15
Zinc influx into cells by the SLC39 gene family - Reactome	↓	1 of 10	0 of 1	1 of 1	0 of 1	0 of 1	0 of 1	6 of 10	1 of 6	1 of 6	0 of 6	0 of 6	1 of 6
γ-glutamyl cycle - MouseCyc	↓	2 of 10	0 of 2	1 of 2	0 of 2	0 of 2	0 of 2	6 of 10	2 of 6	3 of 6	0 of 6	2 of 6	1 of 6

E: Pathways which are excluded from CR analysis

upregulated pathways	downregulated pathways
Smooth Muscle Contraction - Reactome	Pathways in cancer - Mus musculus (mouse)
Disease - Reactome	Chagas disease (American trypanosomiasis) - Mus musculus (mouse)
Legionellosis - Mus musculus (mouse)	XPodNet - protein-protein interactions in the podocyte expanded by STRING - Wikipathways
Collecting duct acid secretion - Mus musculus (mouse)	PluriNetWork - Wikipathways
Defects in vitamin and cofactor metabolism	Toxoplasmosis - Mus musculus (mouse)
Defective AMN causes hereditary megaloblastic anemia 1	Osteoclast differentiation - Mus musculus (mouse)
Defective BTD causes biotidinase deficiency	Hepatitis B - Mus musculus (mouse)
Defective CD320 causes methylmalonic aciduria	Leishmaniasis - Mus musculus (mouse)
Defective CUBN causes hereditary megaloblastic anemia 1	Hepatitis C - Mus musculus (mouse)
Defective GIF causes intrinsic factor deficiency	Epstein-Barr virus infection - Mus musculus (mouse)
Defective HLCS causes multiple carboxylase deficiency	Influenza A - Mus musculus (mouse)
Defective LMBRD1 causes methylmalonic aciduria and homocystinuria type cbIF	Viral carcinogenesis - Mus musculus (mouse)
Defective MMAA causes methylmalonic aciduria type cbIA	Prion diseases - Mus musculus (mouse)
Defective MMAB causes methylmalonic aciduria type cbIB	Adipogenesis - Wikipathways
Defective MMACHC causes methylmalonic aciduria and homocystinuria type cbIC	Staphylococcus aureus infection - Mus musculus (mouse)
Defective MMADHC causes methylmalonic aciduria and homocystinuria type cbID	Herpes simplex infection - Mus musculus (mouse)
Defective MTR causes methylmalonic aciduria and homocystinuria type cbIG	Systemic lupus erythematosus - Mus musculus (mouse)
Defective MTRR causes methylmalonic aciduria and homocystinuria type cbIE	Amoebiasis - Mus musculus (mouse)
Defective MUT causes methylmalonic aciduria mut type	Amyloids - Reactome
Defective TCN2 causes hereditary megaloblastic anemia	Alcoholism - Mus musculus (mouse)
Defects in biotin (Btn) metabolism	
Defects in cobalamin (B12) metabolism	
Alzheimer,s disease - Mus musculus (mouse)	
Huntington,s disease - Mus musculus (mouse)	
Parkinson,s disease - Mus musculus (mouse)	
Non-alcoholic fatty liver disease (NAFLD) - Mus musculus (mouse)	
Proximal tubule bicarbonate reclamation - Mus musculus (mouse)	
Type II diabetes mellitus - Mus musculus (mouse)	

F: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and resveratrol

upregulated pathways	downregulated pathways
cholesterol biosynthesis II (via 24,25-dihydrolanosterol) - MouseCyc	Steroid hormone biosynthesis - Mus musculus (mouse) - KEGG
Pentose and glucuronate interconversions - Mus musculus (mouse) - KEGG	Metabolism of lipids and lipoproteins - Reactome
salvage pathways of pyrimidine ribonucleotides - MouseCyc	Retinol metabolism - Mus musculus (mouse) - KEGG
Metabolism - Reactome	Metabolism - Reactome
Biological oxidations - Reactome	Biological oxidations - Reactome
Diurnally Regulated Genes with Circadian Orthologs - Wikipathways	Protein processing in endoplasmic reticulum - Mus musculus (mouse) - KEGG
Chemical carcinogenesis - Mus musculus (mouse) - KEGG	Fatty acid, triacylglycerol, and ketone body metabolism - Reactome
Cholesterol biosynthesis - Reactome	glycerol-3-phosphate shuttle - MouseCyc
Cholesterol Biosynthesis - Wikipathways	L-cysteine degradation VI - MouseCyc
Metabolism of xenobiotics by cytochrome P450 - Mus musculus (mouse) - KEGG	One carbon metabolism and related pathways - Wikipathways
Cytochrome P450 - arranged by substrate type - Reactome	Drug metabolism - other enzymes - Mus musculus (mouse) - KEGG
Steroid biosynthesis - Mus musculus (mouse) - KEGG	Triglyceride Biosynthesis - Reactome
Porphyrin and chlorophyll metabolism - Mus musculus - KEGG	Taurine and hypotaurine metabolism - Mus musculus (mouse) - KEGG
mevalonate pathway I - MouseCyc	Sulfur amino acid metabolism - Reactome
Terpenoid backbone biosynthesis - Mus musculus - KEGG	nicotine degradation III - MouseCyc
cholesterol biosynthesis III (via desmosterol) - MouseCyc	Meiotic Recombination - Reactome
Phase II conjugation - Reactome	glutathione biosynthesis - MouseCyc
Exercise-induced Circadian Regulation - Wikipathways	
Phase 1 - Functionalization of compounds - Reactome	
cholesterol biosynthesis I - MouseCyc	
Metabolism of lipids and lipoproteins - Reactome	
Glutathione metabolism - Mus musculus (mouse) - KEGG	
superpathway of cholesterol biosynthesis - MouseCyc	
Glucuronidation - Reactome	
Circadian clock tutorial CGrove - Wikipathways	
Retinol metabolism - Mus musculus (mouse) - KEGG	
Circadian rhythm - Mus musculus (mouse) - KEGG	
Metapathway biotransformation - Wikipathways	
Trafficking of dietary sterols - Reactome	
Drug metabolism - cytochrome P450 - mouse - KEGG	
glutathione-mediated detoxification - MouseCyc	
Pyruvate metabolism - Mus musculus (mouse) - KEGG	
Glutathione conjugation - Reactome	
Starch and sucrose metabolism - Mus musculus - KEGG	
nicotine degradation II - MouseCyc	
Circadian Clock - Reactome	
Bile secretion - Mus musculus (mouse) - KEGG	
Glycerolipid metabolism - Mus musculus (mouse) - KEGG	
Drug metabolism - other enzymes - mouse - KEGG	
Nicotinate and nicotinamide metabolism - mouse - KEGG	
trans, trans-farnesyl diphosphate biosynthesis - MouseCyc	
PPAR signaling pathway - Wikipathways	
PPAR signaling pathway - Mus musculus (mouse) - KEGG	

G: Overlapping genes between CR and resveratrol assigned to the overrepresented pathways mentioned in separate and overlapped analysis

Pathway	upregulated genes	downregulated genes
Tryptophan metabolism	Dhcr24	Aldh1a1, Aox1, Cyp2f2, Cyp7b1, Ddc
Sulfur amino acid metabolism	Got1	Bhmt, Cdo1, Csad, Ethe1, Gclc, Gclm
One carbon metabolism and related pathways	Etnk2, Gpx3, Gpx6, Gpx7	Baat, Bhmt, Cdo1, Chka, Csad, Gclc, Gclm
Protein processing in endoplasmic reticulum		Derl3, Dnajb1, Dnajb2, Dnajb11, Ero1lb, Hspa5, Hspf1, Hyou1, Lman2, Nploc4, Pdia6, Sec23b, Sec24d, Stt3b, Syvn1, Vcp, Wfs1
Circadian regulation	Bhlhe40, Cebpb, Cry2, Gfra1, Gstm5, Idi1, Nr1d1, Per1, Per2, Per3, Prkab1, Stbd1, Ugp2	G0s2, Ppp1r3c, Tab2
biological oxidation	Acss2, Cyp2b9, Cyp2b13, Cyp3a13, Cyp4a10, Cyp4a14, Cyp7a1, Cyp17a1, Cyp51, Fmo3, Gstm5, Gstm7, Gsto1, Gstt2, Nnmt, Papss2, Sult1a1, Sult1c2, Ugp2, Ugt1a2	Aldh1a1, Cyp2d9, Cyp2f2, Cyp4a12a, Cyp4a12b, Cyp7b1, Gclc, Gclm, Slc35d1, Sult2a8, Ugt2b1, Ugt2b34, Ugt2b35
Glutathione metabolism	Gpx3, Gpx6, Gpx7, Gstm5, Gstm7, Gsto1, Gstt2, Idh1	
retinol metabolism	Cyp2b9, Cyp2b13, Cyp3a13, Cyp4a10, Cyp4a14, Rdh11, Ugt1a2	Aldh1a1, Aox1, Aox3, Cyp4a12a, Cyp4a12b, Retsat, Ugt2b1, Ugt2b34, Ugt2b35
Meiotic Recombination		H2afx, Hist1h3d, Hist2h2aa1, Hist2h2ac
PPAR signaling pathway	Acox1, Angptl4, Cyp4a10, Cyp4a14, Cyp7a1, Fabp5, Pck1, Ppara	Acaa1b, Cpt2, Cyp4a12a, Cyp4a12b, Rxra
Binding and uptake of ligands by scavenger receptors	Alb, Col4a2, Marco, Saa2	Hba-a1, Scara5
Porphyrin and chlorophyll metabolism	Alas1, Blvra, Gusb, Hmox1, Ugt1a2	Alas2, Ears2, Ugt2b1, Ugt2b34, Ugt2b35
salvage pathways of pyrimidine ribonucleotides	Nme4, Nmrk1	Cda
Nicotinate and nicotinamide metabolism	Enpp1, Nadsyn1, Nmrk1, Nnmt	Aox1, Aox3
Metapathway biotransformation	Akr1b3, Cyp7a1, Cyp17a1, Cyp51, Fmo3, Fmo4, Gal3st1, Gpx3, Gstm5, Gstm7, Gsto1, Gstt2, Nnmt, Sult1a1, Sult1c2, Ugt1a2	Baat, Cyp2f2, Cyp4v3, Cyp7b1, Ephx2, Nat6, Nat8
glycerol-3-phosphate shuttle		Gpd1, Gpd2
Taurine / hypotaurine metab.		Baat, Cdo1, Csad
Pentose and glucuronate interconversions	Akr1b3, Akr1b7, Aldh1b1, Gusb, Ugp2, Ugt1a2	Ugt2b1, Ugt2b34, Ugt2b35, Xylb
Starch and sucrose metabolism	Enpp1, Gusb, Ugp2, Ugt1a2	G6pc, Gck, Ugt2b1, Ugt2b34, Ugt2b35
Pyruvat metabolism	Acss2, Akr1b3, Akr1b7, Aldh1b1, Pck1, Pkm	Glo1
Propanoate metabolism	Acss2, Aldh1b1, Suclg2	
lipid metabolism , Fatty acid degradation, Triglyceride Biosynthesis, Glycerolipid metabolism	Abcg1, Abcg5, Abcg8, Acox1, Alb, Akr1b3, Akr1b7, Aldh1b1, Asah1, Cyp4a10, Cyp4a14, Cyp7a1, Etnk2, Gal3st1, Gpcpd1, Idh1, Lpin2, Pld3, Pnpla2, Ppap2a, Ppara, Ptdss2, Scap, Slco1a4, Ugcg	Abcd1, Acaa1b, Agpat2, Agpat3, Akr1c14, Apoa4, Baat, Chka, Cpt2, Cyp4a12a, Cyp4a12b, Cyp7b1, Elovl3, Ephx2, Ept1, Glyctk, Gpd1, Gpd2, Hsd17b12, Kpnb1, Lpgat1, Mtmr4m, Pla2g6, Rxra, Sgms2, Sgpl1, Sgpp1, Srd5a1, Tbl1x, Hsd3b2, Hsd3b5, Lipa
steroid and cholesterol biosynthesis, Terpenoid backbone biosynthesis, mevalonate pathway, trans, trans-farnesyl diphosphate biosynthesis	Cyp17a1, Cyp51, Dhcr24, Dhcr7, Fdps, Hmgcr, Hmgcs1, Idi1, Lss, Mvk, Nsdhl, Pmvk, Sqle	Cyp7b1, Hsd3b2, Hsd3b5, Hsd17b12, Srd5a1, Ugt2b1, Ugt2b34, Ugt2b35
steroid hormone biosynthesis	Cyp3a13, Cyp17a1, Cyp7a1, Srd5a2, Ugt1a2	Abcb4, Aqp8, Baat, Slc22a7, Rxra
bile secretion; Trafficking of dietary sterols	Abcc4; Abcg5, Abcg8, Atp1b1, Cyp7a1, Hmgcr, Slco1a4	

H: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and phloridzin

upregulated pathways	downregulated pathways
superpathway of cholesterol biosynthesis - MouseCyc	Retinol metabolism - Mus musculus (mouse) – KEGG
Cholesterol biosynthesis - Reactome	nicotine degradation III - MouseCyc
Cholesterol Biosynthesis - Wikipathways	Toll-Like Receptors Cascades - Reactome
cholesterol biosynthesis I - MouseCyc	nicotine degradation II - MouseCyc
cholesterol biosynthesis III (via desmosterol) - MouseCyc	Butanoate metabolism - Mus musculus (mouse) – KEGG
Exercise-induced Circadian Regulation - Wikipathways	Exercise-induced Circadian Regulation – Wikipathways
Metabolism - Reactome	Xenobiotics – Reactome
salvage pathways of pyrimidine ribonucleotides - MouseCyc	Drug metabolism - other enzymes - Mus musculus (mouse) – KEGG
Metabolism of lipids and lipoproteins - Reactome	TRIF-mediated TLR3/TLR4 signaling - Reactome
Pyrimidine salvage reactions - Reactome	MyD88-independent cascade - Reactome
Terpenoid backbone biosynthesis - Mus musculus (mouse) - KEGG	Toll Like Receptor 3 (TLR3) Cascade - Reactome
Fatty Acid Biosynthesis - Wikipathways	mRNA processing – Wikipathways
oleate biosynthesis II (animals) - MouseCyc	bupropion degradation - MouseCyc
Biosynthesis of unsaturated fatty acids - Mus musculus (mouse) - KEGG	Diurnally Regulated Genes with Circadian Orthologs – Wikipathways
Phenylalanine, tyrosine and tryptophan biosynthesis - Mus musculus (mouse) - KEGG	Destabilization of mRNA by AUF1 (hnRNP D0) – Reactome
Destabilization of mRNA by AUF1 (hnRNP D0) - Reactome	MAP kinase activation in TLR cascade – Reactome
cholesterol biosynthesis II (via 24,25-dihydrolanosterol) - MouseCyc	Toll Like Receptor 4 (TLR4) Cascade - Reactome
trans, trans-farnesyl diphosphate biosynthesis - MouseCyc	Valine, leucine and isoleucine degradation - Mus musculus (mouse) – KEGG
(deoxy)ribose phosphate degradation - MouseCyc	Protein processing in endoplasmic reticulum - Mus musculus (mouse) – KEGG
	Activated TLR4 signalling - Reactome
	Toll Like Receptor 9 (TLR9) Cascade - Reactome
	Synthesis of epoxy (EET) and dihydroxyeicosatrienoic acids (DHET) - Reactome
	Porphyrin and chlorophyll metabolism - Mus musculus (mouse) – KEGG
	TRAF6 Mediated Induction of proinflammatory cytokines – Reactome
	Toll Like Receptor TLR6:TLR2 Cascade – Reactome
	Toll Like Receptor 2 (TLR2) Cascade - Reactome
	Type II interferon signaling (IFNG) - Wikipathways
	2-methylbutyrate biosynthesis - MouseCyc
	MyD88 dependent cascade initiated on endosome – Reactome

Toll Like Receptor 7/8 (TLR7/8) Cascade – Reactome
fatty acid β-oxidation IV (unsaturated, even number) – MouseCyc
Toll Like Receptor signaling - Wikipathways
PPAR signaling pathway - Wikipathways
PPAR signaling pathway - <i>Mus musculus</i> (mouse) – KEGG
Fatty Acid Beta Oxidation - Wikipathways
fatty acid β-oxidation I - MouseCyc
MyD88 cascade initiated on plasma membrane – Reactome
Toll Like Receptor 10 (TLR10) Cascade – Reactome
Toll Like Receptor 5 (TLR5) Cascade - Reactome
MyD88:Mal cascade initiated on plasma membrane – Reactome
Toll Like Receptor TLR1:TLR2 Cascade – Reactome
Fatty acid degradation - <i>Mus musculus</i> (mouse) – KEGG
Complement Activation, Classical Pathway – Wikipathways
Regulation of mRNA Stability by Proteins that Bind AU-rich Elements - Reactome

I: Overlapping genes between CR and phloridzin assigned to the overrepresented pathways mentioned in separate and overlapped analysis

Pathway	upregulated genes	downregulated genes
IL-6 signaling	Erbb3, Hspb1	Hsp90aa1, Jak1, Jun, Map2k4, Mapk1, Pik3r1, Ppp2ca, Ppp2r1b, Ppp2r5e, Stat5a
Growth hormone receptor signaling	Cish, Socs2	Adam17, Ghr, Irs1, Stat5a
Androgen Receptor Signaling	Ccne1, Cdk1	Atf2, Casp8, Ctnnb1, Ghr, Gtf2f2, Gtf2h1, Jun, Mapk1, Nr3c1, Rchy1
RNA transport		Eif1a, Eif1b, Eif2b3, Eif2s3x, Eif4a2, Eif4g1, Eif5, Eif5b, Gemin2, Kpnb1, Ncbp2, Nup35, Nxt2, Pabpc1, Pnn, Rbm8a, Rpp14, Rpp40, Smn1
Butanoate metabolism		Abat, Acat1, Acsm3, Acsm5, Aldh5a1, Bdh1, Ehhadh, Hadha
mRNA processing & regulation of mRNA stability	Cirbp, Hspb1, Rps13, Ttc39a	A1cf, Abcb4, Akap1, Anp32a, Apobec1, Cpeb4, Cpsf2, Csad, Csde1, Eif2ak2, Eif4a2, Eif4g1, Ergic2, Fus, G3bp1, Gatc, Ggcx, Gtf2f2, Gtf2h1, Gtf2h2, Hnrnpa3, Hnrnpu, Hspa1b, Lsm2, Ltv1, Mettl3, Mki67ip, Ncbp2, Ncl, Nol8, Nop9, Npm1, Nup35, Oasl2, Pabpc1, Papola, Parn, Pcbp4, Pum1, Rad21, Rbm8a, Rbmx1, Rpp14, Rps4l, Smn1, Spop, Srp68, Srsf1, Srsf10, Srsf3, Srsf7, Ssb, Syncrip, Tardbp, Vprbp, Xrn2, Zfp36l1
lipid metabolism and lipo-protein metabolism, Fatty acid biosynthesis; Fatty	Acacb, Acot3, Acss2, Dhcr7, Fads1, Fdps, Gal3st1, Gpcpd1, Idi1, Nsdhl, Pmvk, Scd1, Sqle	9130409l23Rik, Abca1, Abcc3, Acaa1b, Acad11, Acadm, Acat1, Acot1, Acox2, Akr1c14, Akr1c20, Akr1d1, Bdh1, Chkb,

acid, triacylglycerol, and ketone body metabolism; beta-oxidation		Cpt1a, Decr1, Eci2, Ehhadh, Elovl3, Gyk, Hadha, Hsd3b2, Hsd3b3, Hsd17b12, Kpnb1, Ldlr, Pik3r1, Pitpn, Pla2g12a, Ppap2b, Ppara, Psap, Sc5d, Sgms2, Sgpp1, Slco1a1, Slco1b2, Srd5a1, Sumf1, Tbl1x
arachidonic acid metabo- lism	Cyp2c39, Cyp4a14	Cyp2c37, Cyp2c50, Cyp2c54, Cyp2j5, Cyp2j6
propanoate metabolism	Acacb, Acss2	Abat, Acadm, Acat1, Aldh3a2, Ehhadh, Hadha
Phenylalanine, tyrosine/ tryptophan biosynthesis	Got1	
Tryptophan metabolism, Valine, leucine and isoleu- cine degradation		Aadat, Abat, Acaa1b, Acadm, Acat1, Aldh3a2, Aox1, Aox3, Cyp2j6, Ehhadh, Hadha
Protein processing in en- doplasmic reticulum		Atf6, Atf6b, Canx, Capn2, Derl2, Dnaja1, Dnajb1, Dnajc5, Eif2ak2, Hsp90aa1, Hsp90ab1, Hsp90b1, Hspa1b, Hspf1, Nfe2l2, Nsf1l1c, Rnf5, Sec62, Sec63, Ube2d1, Ube2e3, Ube2g2, Ufd1
circadian clock	Idi1, Per1, Tubb4a	Azin1, Clock, Dnaja1, G0s2, Klf9, Nr1d1, Ppp1r3c, Psma4, Stbd1, Tab2
Bile secretion		Abcb4, Abcc2, Abcc3, Abcg2, Aqp4, Aqp9, Ephx1, Ldlr, Slc22a1, Slc22a7, Slco1a1, Slco1b2
retinol metabolism	Cyp2c39, Cyp4a14	Adh4, Aox1, Aox3, Cyp2a12, Cyp2c37, Cyp2c50, Cyp2c54, Cyp3a13, Cyp3a25, Dhrs4, Rdh16, Retsat, Ugt2b34, Ugt2b35
glutathione metabolism and γ -glutamyl cycle	Ggt6	Gclc, Gclm, Oplah
Cholesterol biosynthesis	Dhcr7, Fdps, Idi1, Nsdhl, Pmvk, Sqle	Acat1, Nudt12, Sc5d
Steroid hormone biosyn- thesis		Akr1d1, Cyp3a13, Cyp3a25, Hsd17b12, Hsd17b2, Hsd3b2, Srd5a1, Ugt2b34, Ugt2b35
PPAR signaling pathway	Cyp4a14, Fabp5, Scd1	Acaa1b, Acadm, Acox2, Cd36, Cpt1a, Ehhadh, Fabp2, Fabp7, Gyk, Ppara, Sorbs1
Porphyrin and chlorophyll metabolism	Alas1	Acaa1b, Acot1, Decr1, Hadha, Hsd17b12
Kennedy pathway		Chka, Chkb, Chpt1, Sgpl1
biological oxidation	Acss2, Cyp2c39, Cyp4a14, Nnmt	Adh4, Cyp2a12, Cyp2c37, Cyp2c50, Cyp2c54, Cyp2d12, Cyp2d40, Cyp2j5, Cyp2j6, Cyp3a13, Cyp3a25, Gclc, Gclm, Sult1c2, Tpmt, Ephx1, Ugt2b34, Ugt2b35
Toll like receptor cascade	Cdk1	Atf2, Casp8, Cd36, Chuk, Ctss, Dhx36, Eea1, Eif2ak2, Hsp90b1, Ikbke, Ikbkg, Irak2, Jun, Ly86, Map2k4, Mapk1, Ppp2ca, Ppp2r1b, Tab2, Tbk1, Ube2d1 Cxcl10, Cxcl9, Cybb, Eif2ak2, Gbp2b, Ifit2, Jak1, Stat2
Type II interferon signaling (IFNG)		Tk2
Pyrimidine salvage reac- tions		
Complement Activation		C1qb, C1s, C6, C8a, C8b
Class I MHC mediated antigen processing & presentation		Anapc2, Anapc4, Asb13, Canx, Cd36, Ctss, Cul2, Cul3, Cybb, Pja2, Rchy1, Rnf138, Rnf6, Smurf2, Tceb1, Uba3, Uba5, Vprbp
(deoxy)ribose phosphate degradation		Apobec1
2-methylbutyrate biosyn- thesis		Acat1, Ehhadh

J: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and dietary quercetin

upregulated pathways	downregulated pathways
Cytochrome P450 - arranged by substrate type - Reactome	Destabilization of mRNA by AUF1 (hnRNP D0) – Reactome
Biological oxidations - Reactome	
Macrophage markers - Wikipathways	
Diurnally Regulated Genes with Circadian Orthologs - Wikipathways	
Exercise-induced Circadian Regulation - Wikipathways	
Phase 1 - Functionalization of compounds - Reactome	
Retinol metabolism - <i>Mus musculus</i> - KEGG	
Hypertrophy Model - Wikipathways	

K: Overlapped genes between CR and dietary quercetin assigned to the overrepresented pathways mentioned in separate and overlapped analysis

Pathway	upregulated genes	downregulated genes
Kennedy pathway		Chka, Sgpl1
Destabilization of mRNA by AUF1	Hspb1	Hspa1b
Circadian clock	Cebpb, Per1	G0s2, Nr1d1, Ppp1r3c
biological oxidation	Cyp2b10, Cyp4a14, Nnmt, Sult1e1	Cyp2d12, Cyp2d40, Slc26a1, Sult1c2
retinol metabolism	Cyp2b10, Cyp4a14	
Macrophage markers	Cd14, Cd163	

L: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and quercetin administered intraperitoneally

upregulated pathways	downregulated pathways
Biological oxidations - Reactome	Biological oxidations - Reactome
Metabolism - Reactome	Metabolism - Reactome
Phase 1 - Functionalization of compounds - Reactome	Phase 1 - Functionalization of compounds – Reactome
Cytochrome P450 - arranged by substrate type - Reactome	Cytochrome P450 - arranged by substrate type – Reactome
Chemical carcinogenesis - <i>Mus musculus</i> (mouse) - KEGG	Chemical carcinogenesis - <i>Mus musculus</i> (mouse) - KEGG
Drug metabolism - cytochrome P450 - <i>Mus musculus</i> (mouse) - KEGG	Drug metabolism - cytochrome P450 - <i>Mus musculus</i> (mouse) - KEGG
Metabolism of xenobiotics by cytochrome P450 - <i>Mus musculus</i> (mouse) - KEGG	Metabolism of xenobiotics by cytochrome P450 - <i>Mus musculus</i> (mouse) - KEGG
Retinol metabolism - <i>Mus musculus</i> (mouse) - KEGG	Retinol metabolism - <i>Mus musculus</i> (mouse) – KEGG
Metabolism of lipids and lipoproteins - Reactome	Metabolism of lipids and lipoproteins - Reactome
Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) - Reactome	Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) - Reactome
Biosynthesis of unsaturated fatty acids - <i>Mus musculus</i> (mouse) - KEGG	Complement and coagulation cascades - <i>Mus musculus</i> (mouse) - KEGG
Growth hormone receptor signaling - Reactome	Growth hormone receptor signaling - Reactome
Fatty acids - Reactome	Fatty acids - Reactome

Metapathway biotransformation - Wiki-pathways	Complement Activation, Classical Pathway - Wikipathways
Arachidonic acid metabolism - <i>Mus musculus</i> (mouse) - KEGG	Arachidonic acid metabolism - <i>Mus musculus</i> (mouse) - KEGG
nicotine degradation II - MouseCyc	nicotine degradation II - MouseCyc
Leptin Insulin Overlap - Wikipathways	Bile acid and bile salt metabolism - Reactome
nicotine degradation III - MouseCyc	nicotine degradation III - MouseCyc
Glycolysis and Gluconeogenesis - Wiki-pathways	Synthesis of bile acids and bile salts - Reactome
glutathione-mediated detoxification - MouseCyc	Primary bile acid biosynthesis - <i>Mus musculus</i> (mouse) - KEGG
Phase II conjugation - Reactome	Beta-oxidation of pristanoyl-CoA - Reactome
Arachidonic acid metabolism - Reactome	Complement cascade - Reactome
Linoleic acid (LA) metabolism - Reactome	Arachidonic acid metabolism - Reactome
Adipocytokine signaling pathway - <i>Mus musculus</i> (mouse) - KEGG	Synthesis of bile acids and bile salts via 7alpha-hydroxycholesterol - Reactome
PPAR signaling pathway - <i>Mus musculus</i> (mouse) - KEGG	PPAR signaling pathway - <i>Mus musculus</i> (mouse) - KEGG
PPAR signaling pathway - Wikipathways	PPAR signaling pathway - Wikipathways
Insulin signaling pathway - <i>Mus musculus</i> (mouse) - KEGG	Complement and Coagulation Cascades - Wikipathways
oleate biosynthesis II (animals) - MouseCyc	Terminal pathway of complement - Reactome
Xenobiotics - Reactome	Xenobiotics - Reactome
Fatty acid degradation - <i>Mus musculus</i> (mouse) - KEGG	Type II interferon signaling (IFNG) - Wikipathways
alpha-linolenic (omega3) and linoleic (omega6) acid metabolism - Reactome	Fatty acid degradation - <i>Mus musculus</i> (mouse) - KEGG
Signaling by Leptin - Reactome	Retinol metabolism - Wikipathways
Estrogen metabolism - Wikipathways	Peroxisomal lipid metabolism - Reactome
biosynthesis of estrogens - MouseCyc	Interferon Signaling - Reactome
alpha-linolenic acid (ALA) metabolism - Reactome	Regulation of Complement cascade - Reactome
Methylation - Wikipathways	ISG15 antiviral mechanism - Reactome
Glutathione metabolism - <i>Mus musculus</i> (mouse) - KEGG	Antiviral mechanism by IFN-stimulated genes - Reactome
Cytosolic sulfonation of small molecules - Reactome	Steroid hormone biosynthesis - <i>Mus musculus</i> (mouse) - KEGG
glycolysis V (<i>Pyrococcus</i>) - MouseCyc	Steroid Biosynthesis - Wikipathways
Steroid hormone biosynthesis - <i>Mus musculus</i> (mouse) - KEGG	Synthesis of bile acids and bile salts via 27-hydroxycholesterol - Reactome
Fatty Acyl-CoA Biosynthesis - Reactome	Peroxisome - <i>Mus musculus</i> (mouse) - KEGG
bupropion degradation - MouseCyc	Immune System - Reactome
Glutathione metabolism - Wikipathways	Initial triggering of complement - Reactome
Selenium - Wikipathways	tetrapyrrole biosynthesis II - MouseCyc
glycolysis III - MouseCyc	bupropion degradation - MouseCyc
Glycolysis / Gluconeogenesis - <i>Mus musculus</i> (mouse) - KEGG	Synthesis of bile acids and bile salts via 24-hydroxycholesterol - Reactome
Prolactin signaling pathway - <i>Mus musculus</i> (mouse) - KEGG	Synthesis of epoxy (EET) and dihydroxyeicosatrienoic acids (DHET) - Reactome
Fatty acid elongation - <i>Mus musculus</i> (mouse) - KEGG	Classical antibody-mediated complement activation - Reactome
glycolysis I - MouseCyc	Fatty Acid Omega Oxidation - Wikipathways
Drug metabolism - other enzymes - <i>Mus musculus</i> (mouse) - KEGG	Bile secretion - <i>Mus musculus</i> (mouse) - KEGG
Glycolysis - Reactome	ketogenesis - MouseCyc

Regulation of Insulin-like Growth Factor (IGF) Transport and Uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs) - Reactome	Antigen processing and presentation - Mus musculus (mouse) - KEGG
Triglyceride Biosynthesis - Reactome	Ethanol oxidation - Reactome
Glucose transport - Reactome	Synthesis and degradation of ketone bodies - Mus musculus (mouse) - KEGG
Hexose transport - Reactome	Heme Biosynthesis - Wikipathways
EPO Receptor Signaling - Wikipathways	heme biosynthesis II - MouseCyc
Metabolism of carbohydrates - Reactome	sphingosine and sphingosine-1-phosphate metabolism - MouseCyc
	Cytokine Signaling in Immune system - Reactome
	Lipoprotein metabolism - Reactome
	Interleukin-6 signaling - Reactome
	Creation of C4 and C2 activators - Reactome
	mevalonate pathway I - MouseCyc
	Heme biosynthesis - Reactome
	Innate Immune System - Reactome
	Porphyrin and chlorophyll metabolism - Mus musculus (mouse) - KEGG
	Recycling of bile acids and salts - Reactome
	Transport of organic anions - Reactome
	Bile salt and organic anion SLC transporters – Reactome
	Protein processing in endoplasmic reticulum - Mus musculus (mouse) - KEGG
	Androgen biosynthesis - Reactome
	Phagosome - Mus musculus (mouse) - KEGG
	Metabolism of porphyrins - Reactome
	MHC class II antigen presentation - Reactome
	Fat digestion and absorption - Mus musculus (mouse) - KEGG
	Tryptophan metabolism - Wikipathways
	Chylomicron-mediated lipid transport - Reactome
	Lipid digestion, mobilization, and transport – Reactome
	Proteasome - Mus musculus (mouse) - KEGG
	Statin Pathway - Wikipathways
	Retinoid metabolism and transport - Reactome
	Linoleic acid metabolism - Mus musculus (mouse) – KEGG
	Exercise-induced Circadian Regulation - Wikipathways
	Diurnally Regulated Genes with Circadian Orthologs - Wikipathways
	One carbon metabolism and related pathways - Wikipathways
	Terpenoid backbone biosynthesis - Mus musculus (mouse) - KEGG
	Synthesis of Prostaglandins (PG) and Thromboxanes (TX) - Reactome
	Valine, leucine and isoleucine degradation - Mus musculus (mouse) - KEGG

M: Overlapping genes between CR and quercetin administered intraperitoneally assigned to the overrepresented pathways mentioned in separate and overlapped analysis

pathway	upregulated genes	downregulated genes
Tryptophan metabolism	Inmt, Tdo2	Adat, Aldh1a1, Aox1, Cyp7b1
Valine, leucine and isoleucine degradation	Ehhadh	Abat, Acat2, Aox1, Hadhb, Hmgcs2
One carbon metabolism and related pathways		Bhmt2, Chkb, Csad, Gclc, Gsr, Gss
Protein processing in endoplasmic reticulum		Bag1, Derl1, Derl3, Dnaja1, Dnajb2, Edem3, Eif2ak2, Hspa5, Nfe2l2, Pdia4, Sec23b, Stt3b, Ube2e1, Ube2e3
Circadian Regulation	Tob1	Cebpb, Dnaja1, G0s2, Psma4, Qk, Stbd1, Sumo3
Bile acid and bile salt metabolism; Synthesis of bile acids and bile salts from cholesterol, recycling of bile acids and salts	Adcy6, Sult2a3	Acox2, Akr1c14, Amacr, Aqp8, Cyp7b1, Ephx1, Hsd17b4, Hsd3b7, Slc22a7, Slco1a1, Slco1b2
Bile salt and organic anion SLC transporters		Slc13a3, Slc16a1, Slc16a7
biological oxidation	Cyp17a1, Cyp2b13, Cyp2c39, Cyp4a10, Cyp4a14, Cyp4a31, Fmo3, Gsta4, Gstm2, Nnmt, Sult1e1, Sult2a3	Adh4, Aldh1a1, Aldh1a7, Cyp2a12, Cyp2c29, Cyp2c50, Cyp2c70, Cyp2d10, Cyp2d12, Cyp2d40, Cyp2d9, Cyp2j5, Cyp3a13, Cyp3a25, Cyp4a12a, Cyp4a12b, Cyp7b1, Ephx1, Gclc, Ggh, Glyat, Gss, Gstm4, Mgst1, Ugdh, Ugt2b35, Ugt2b37
Steroid hormone biosynthesis, androgen and estrogen biosynthesis	Cyp17a1, Sts, Sult1e1	Cyp3a13, Cyp3a25, Cyp7b1, Hsd17b2, Hsd3b5, Srd5a1, Ugt2b35, Ugt2b37
Steroid biosynthesis	Cyp17a1, Lss	Hsd17b2, Hsd17b4, Hsd3b5, Sc4mol, Sc5d
Retinol / Retinoid metabolism and transport	Cyp2b13, Cyp2c39, Cyp4a10, Cyp4a14, Cyp4a31	Adh4, Akr1c14, Aldh1a1, Aldh1a7, Aox1, Cd36, Cyp2a12, Cyp2c29, Cyp2c50, Cyp2c70, Cyp3a13, Cyp3a25, Cyp4a12a, Cyp4a12b, Dhrs3, Gpc4, Sdc1, Sdc2, Sdc4, Ugt2b35, Ugt2b37
PPAR signaling pathway	Cyp4a10, Cyp4a14, Cyp4a31, Ehhadh, Fads2, Slc27a1	Acox2, Cd36, Cyp4a12a, Cyp4a12b, Hmgcs2, Scd1
Porphyrin / chlorophyll metab.		Alad, Ugt2b35, Ugt2b37
Type II interferon signaling (IFNG)		Cxcl10, Cxcl9, Eif2ak2, Ifngr2, Isg15, Soc3
Metabolism of lipids and lipoproteins; biosynthesis of fatty acids, fatty acyl-CoA biosynthesis; Linoleic acid (LA) and ALA metabolism, Fatty acid elongation, beta or omega oxidation, Lipoprotein metabolism , Fat digestion, absorption, mobilization, and transport, ketogenesis, statin pathway, sphingosine and sphingosine-1-phosphate metabolism, arachidonic acid metabolism	Acly, Acot7, Cyp17a1, Cyp2b13, Cyp2c39, Cyp4a10, Cyp4a14, Cyp4a31, Elovl6, Fads1, Fads2, Lss	Abca1, Acat2, Acox2, Adh4, Agpat2, Akr1c14, Aldh1a1, Amacr, Bmp1, Cd36, Cdipt, Chkb, Crot, Cyb5, Cyp2c29, Cyp2c50, Cyp2c70, Cyp2j5, Cyp3a13, Cyp3a25, Cyp4a12a, Cyp4a12b, Cyp7b1, Elovl3, Elovl5, Hadhb, Hmgcs2, Hpgd, Hsd17b4, Hsd3b5, Hsd3b7, Lcat, Lpgat1, Ppp1ca, Ppp1cb, Psap, Ran, Sc4mol, Sc5d, Sdc1, Sgp1, Sgpp1, Slco1a1, Slco1b2, Sqle, Srd5a1
Metabolism of carbohydrates, Glycolysis and Gluconeogenesis	Eno1, Pklr, Slc2a4	Adh4, Amy1, B3gat3, Csgalact2, Gpc4, Ldha, Mpc1, Sdc1, Sdc2, Sdc4, Slc25a13
Glucose/ Hexose transport	Slc2a4	Gclc, Gsr, Gss, Gstm4, Mgst1
Glutathione metabolism	Gsta4, Gstm2	

Methylation	Inmt, Nnmt	
Interferon Signaling		Ifit1, Ifngr2, Isg15, Socs3, U-be2e1, Ube2l6, Usp18
Immune System, Complement and coagulation cascades, Innate immune system, Creation of C4 and C2 activators	Adcy6, Cish, Irf7, Irs2, Nrg4, Rnf4, Socs2, Tirap	Actg1, Atg5, B2m, C1qa, C1s, C2, C4b, C4bp, C8a, C8b, C9, Capza2, Cd36, Cd74, Cdh1, Cfh, Crp, Ctnnb1, Ctss, Cyba, Dhx36, Egfr, F8, Ghr, H2-Aa, Hras, Ifit1, Ifngr2, Il6st, Isg15, Lbp, Ly86, Map2k4, Mbl1, Rap1a, Rnf135, Serping1, Socs3, Stat5a, Trex1, Tyrobp, Uba3, Ube2a, Ube2e1, Ube2l6, Usp18
Peroxisome, Peroxisomal lipid metabolism	Ehhadh	Acox2, Amacr, Crot, Ech1, Hsd17b4, Pex13, Pex26
Heme Biosynthesis		Alad
Selenium	Fads1, Fads2	Gsr
Phagosome		Actg1, Atp6v1h, Cd36, Clec7a, Ctss, Cyba, Dync1i2, H2-Aa, H2-Ab1, Mbl1, Sec22b
Proteasome		Psma4, Psmb8, Psmc2, Psmd1, Psmd6
Growth hormone receptor signaling	Cish, Irs2, Socs2	Ghr, Socs3, Stat5a
Adipocytokine signaling pathway, insulin signaling pathway, Leptin Insulin Overlap	Irs2, Pklr, Lepr, Slc2a4, Socs2	Cd36, Hras, Ppp1ca, Ppp1cb, Rheb, Socs3, Stat5a
Regulation of Insulin-like Growth Factor (IGF) Transport and Uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)	Igfbp1	Igfals
Transport of organic anions		Slco1a1, Slco1b2
Interleukin-6 signaling		Il6st, Socs3
Ethanol oxidation		Adh4, Aldh1a1, Aldh1a7

N: Pathways mentioned in overrepresentation analysis for overlapping genes between CR and genistein administered intraperitoneally

upregulated pathways	downregulated pathways
Biological oxidations - Reactome	Biological oxidations - Reactome
Metabolism - Reactome	Metabolism - Reactome
Phase 1 - Functionalization of compounds - Reactome	Phase 1 - Functionalization of compounds – Reactome
Cytochrome P450 - arranged by substrate type - Reactome	Cytochrome P450 - arranged by substrate type – Reactome
Chemical carcinogenesis - Mus musculus (mouse) - KEGG	Chemical carcinogenesis - Mus musculus (mouse) - KEGG
Drug metabolism - cytochrome P450 - Mus musculus (mouse) - KEGG	Drug metabolism - cytochrome P450 - Mus musculus (mouse) - KEGG
Metabolism of xenobiotics by cytochrome P450 - Mus musculus (mouse) - KEGG	Metabolism of xenobiotics by cytochrome P450 - Mus musculus (mouse) - KEGG
Retinol metabolism - Mus musculus (mouse) - KEGG	Retinol metabolism - Mus musculus (mouse) – KEGG
nicotine degradation II - MouseCyc	nicotine degradation II - MouseCyc
Arachidonic acid metabolism - Mus musculus (mouse) - KEGG	Complement and Coagulation Cascades - Wikopathways
Growth hormone receptor signaling - Reactome	Complement and coagulation cascades - Mus musculus (mouse) - KEGG
Xenobiotics - Reactome	Initial triggering of complement - Reactome
Metabolism of lipids and lipoproteins - Reactome	Metabolism of lipids and lipoproteins - Reactome

Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) - Reactome	Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) - Reactome
Biosynthesis of unsaturated fatty acids - Mus musculus (mouse) - KEGG	Complement Activation, Classical Pathway - Wikipathways
Fatty acid elongation - Mus musculus - KEGG	Complement cascade - Reactome
Arachidonic acid metabolism - Reactome	ISG15 antiviral mechanism - Reactome
Metapathway biotransformation - Wikipathways	Antiviral mechanism by IFN-stimulated genes – Reactome
Estrogen metabolism - Wikipathways	mevalonate pathway I - MouseCyc
bupropion degradation - MouseCyc	Interferon Signaling - Reactome
Triglyceride Biosynthesis - Reactome	nicotine degradation III - MouseCyc
Phenylalanine, tyrosine and tryptophan biosynthesis - Mus musculus (mouse) - KEGG	Transport of glucose and other sugars, bile salts and organic acids, metal ions and amine compounds - Reactome
nicotine degradation III - MouseCyc	tetrapyrrole biosynthesis II - MouseCyc
Phenylalanine metabolism - Mus musculus (mouse) - KEGG	Terpenoid backbone biosynthesis - Mus musculus (mouse) - KEGG
biosynthesis of estrogens - MouseCyc	Fatty acids - Reactome
Glycolysis and Gluconeogenesis - Wikipathways	Fatty acid degradation - Mus musculus (mouse) – KEGG
Cytosolic sulfonation of small molecules - Reactome	Destabilization of mRNA by AUF1 (hnRNP D0) – Reactome
Prolactin signaling pathway - Mus musculus (mouse) - KEGG	Classical antibody-mediated complement activation - Reactome
Fatty acid, triacylglycerol, and ketone body metabolism - Reactome	superpathway of cholesterol biosynthesis – MouseCyc
Phase II conjugation - Reactome	Metabolism of porphyrins - Reactome
Leptin Insulin Overlap - Wikipathways	ketogenesis - MouseCyc
acyl-CoA hydrolysis - MouseCyc	Beta-oxidation of pristanoyl-CoA - Reactome
Fatty acids - Reactome	Lipoprotein metabolism - Reactome
glutathione-mediated detoxification - MouseCyc	Synthesis and degradation of ketone bodies - Mus musculus (mouse) - KEGG
EPO Receptor Signaling - Wikipathways	Heme Biosynthesis - Wikipathways
Fatty Acyl-CoA Biosynthesis - Reactome	heme biosynthesis II - MouseCyc
Gluconeogenesis - Reactome	Abacavir transport and metabolism - Reactome
Cysteine and methionine metabolism - Mus musculus (mouse) - KEGG	Porphyrin and chlorophyll metabolism - Mus musculus (mouse) - KEGG
IL-3 Signaling Pathway - Wikipathways	Creation of C4 and C2 activators - Reactome
	Heme biosynthesis - Reactome
	SLC-mediated transmembrane transport – Reactome
	Organic cation/anion/zwitterion transport – Reactome
	Bile salt and organic anion SLC transporters – Reactome
	Bile acid and bile salt metabolism - Reactome
	Bile secretion - Mus musculus (mouse) - KEGG
	Retinol metabolism - Wikipathways
	PPAR signaling pathway - Mus musculus (mouse) – KEGG
	PPAR signaling pathway - Wikipathways
	Protein processing in endoplasmic reticulum - Mus musculus (mouse) - KEGG
	Peroxisome - Mus musculus (mouse) - KEGG
	Lipid digestion, mobilization, and transport – Reactome
	Proteasome - Mus musculus (mouse) - KEGG
	Primary bile acid biosynthesis - mouse - KEGG

O: Overlapping genes between CR and genistein administered intraperitoneally assigned to the overrepresented pathways mentioned in separate and overlapped analysis

Pathway	upregulation	downregulation
Growth hormone receptor signaling, Leptin Insulin Overlap	Cish, Irs2, Socs2	
biological oxidation	Cyp17a1, Cyp2b13, Cyp2b9, Cyp2c39, Cyp4a14, Cyp4a31, Fmo3, Gstt3, Sult1e1, Sult2a3	Adh4, Cyp2c29, Cyp2c55, Cyp2c70, Cyp2d10, Cyp2d12, Cyp2d40, Cyp2d9, Cyp3a13, Cyp3a25, Cyp4a12a, Cyp4a12b, Ephx1, Glyat, Gss, Gstm4, Mgst1, Ugt2b35, Ugt2b37
Protein processing in endoplasmic reticulum	Map3k5	Derl1, Dnaja1, Dnajb1, Dnajb2, Hspa5, Hspf1, Nfe2l2, Stt3b, Ube2e1
Bile secretion	Sult2a3	Ephx1, Slc10a2, Slc22a1, Slc22a7, Slco1b2
Retinol metabolism	Cyp2b13, Cyp2b9, Cyp2c39, Cyp4a14, Cyp4a31	Adh4, Aox1, Cyp2c29, Cyp2c55, Cyp2c70, Cyp3a13, Cyp3a25, Cyp4a12a, Cyp4a12b, Dhrs3, Ugt2b35, Ugt2b37
Cholesterol biosynthesis	Lss	Acat2, Hmgcs2
Metabolism of lipids and lipoproteins, Fatty acid, triacylglycerol, and ketone body metabolism, biosynthesis of fatty acids, beta oxidation, ketogenesis, Lipid digestion, mobilization, and transport, acyl-CoA hydrolysis, sphingosine and sphingosine-1-phosphate metabolism	Acly, Acot3, Agpat9, Cyp17a1, Cyp2c39, Cyp4a14, Cyp4a31, Elovl6, Gpcpd1, Lss, Ptgds	Acat2, Amacr, Cris1, Crot, Cyp2c29, Cyp2c55, Cyp4a12a, Cyp4a12b, Elovl3, Hadhb, Hmgcs2, Hsd3b5, Hsd3b7, Lcat, Ppp1ca, Psap, Sdc1, Sgpp1, Slc10a2, Slco1b2
Proteasome		Psmb8, Psmc2, Psmd1, Psmd4, Psmd6
Estrogen metabolism	Cyp17a1, Sult1e1	
Bile acid and bile salt metabolism; Primary bile acid biosynthesis		Amacr, Hsd3b7, Slc10a2, Slco1b2
Complement and coagulation cascades		C1qa, C1s, C4b, C4bp, C9, Cfd, Mbl1, Serping1
Interferon Signaling		Ifit1, Ifngr2, Isg15, Ube2e1, Ub2l6, Usp18
Peroxisome	Ehhadh	Amacr, Crot, Ech1, Pex26
Heme Biosynthesis		Alad, Alas2
Cysteine and methionine metabolism	Got1, Tat	Ldha, Mpst
Phenylalanine metabolism	Got1, Tat	Glyat
Retinol metabolism	Rarb	Adh4, Dhrs3
IL-3 Signaling Pathway	Cish, Id1, Socs2	Shc1
Transport of glucose and other sugars, bile salts and organic acids, metal ions and amine compounds, SLC-mediated transmembrane transport	Slc27a1, Slc43a1	Rhbg, Slc13a3, Slc16a7, Slc1a4, Slc22a1, Slc22a7, Slc39a4, Slc6a13, Slco1b2

P: All significantly regulated genes by caloric restriction with a fold change > 1.5 and the regulation of these genes by the different phytonutrients
 (grey bars indicate that the gene is significantly co-regulated by the phytonutrient ($q < 0.05$))

Upregulated genes

Entrez ID	gene symbol	FC CR	FC R	FC P	FC Qd	FC Qi	FC Gi
100503969	1810008I18Rik	0,54	NA	0,8441*	1,19	NA	NA
109163	3010003L21Rik	0,65	1,09	NA	NA	NA	NA
107476	Acaca	0,52	1,10	0,95	1,08	0,74	0,87
100705	Acacb	0,55	1,01	0,8314*	1,09	1,01	1,06
266645	Acmsd	0,67	0,99	NA	NA	1,04	1,05
101494	Al132709	0,34	NA	1,03	1,04	NA	NA
27053	Asns	0,65	1,01	0,8699*	1,00	0,88	0,96
216454	BC089597	0,65	0,689*	0,89	0,93	0,76	0,93
100038725	Cep85l	0,65	NA	NA	NA	NA	NA
56398	Chp1	0,57	0,6905*	NA	NA	1,60	1,21
226744	Cnst	0,64	0,98	1,05	1,13	0,83	0,82
74161	Ctcflos	0,61	NA	0,91	0,94	NA	NA
13088	Cyp2b10	0,65	1,04	1,42	0,8061*	1,27	1,25
13089	Cyp2b13	0,50	0,4516*	0,93	1,17	0,7212*	0,349*
243881	Cyp2b23	0,32	NA	NA	NA	1,11	1,14
13094	Cyp2b9	0,37	0,3154*	1,06	1,20	0,79	0,7088*
56050	Cyp39a1	0,51	1,00	0,96	1,16	1,66	1,36
13117	Cyp4a10	0,63	0,5301*	NA	NA	0,682*	0,93
320187	E230001N04Rik	0,61	0,90	NA	NA	NA	NA
20918	Eif1	0,65	1,06	NA	NA	1,16	1,11
108112	Eif4ebp3	0,21	0,5482*	NA	NA	0,94	0,621*
170439	Elovl6	0,66	1,26	1,04	1,19	0,7564*	0,6802*
16592	Fabp5	0,63	0,7866*	0,8948*	1,00	4,25	1,90
73385	Fam177a	0,44	NA	NA	NA	0,98	1,01
208164	Fam180a	0,65	0,734*	NA	NA	1,13	1,20
14104	Fasn	0,46	1,42	0,98	1,34	1,22	1,18
14229	Fkbp5	0,67	0,3323*	0,88	0,7626*	1,01	1,16
14262	Fmo3	0,20	0,5247*	0,87	0,95	0,1682*	0,2886*
14381	G6pdx	0,61	0,94	0,99	1,03	1,27	1,34
654467	Gm10052	0,62	NA	NA	NA	1,55	1,17
667250	Gm12657	0,61	NA	NA	NA	NA	NA
433520	Gm14403	0,66	NA	NA	NA	0,86	0,93
100043257	Gm15453	0,21	NA	NA	NA	1,10	0,96
65956	Gm21541	0,65	NA	NA	NA	1,33	1,76
100041932	Gm3579	0,35	NA	NA	NA	NA	NA
333331	Gm5131	0,60	NA	NA	NA	0,72	0,85
629557	Gm6981	0,65	NA	NA	NA	1,13	1,04
14718	Got1	0,53	0,6449*	0,8357*	0,8839*	2,18	0,6836*
14860	Gsta4	0,65	1,46	1,27	1,18	0,7017*	0,85
14864	Gstm3	0,40	1,04	1,11	1,05	1,35	1,66
232493	Gys2	0,65	NA	1,08	1,08	1,18	1,06
73738	Haus7	0,67	0,94	1,09	1,07	1,32	1,25

217845	Ifl27l2b	0,57	0,7565*	0,8721*	0,8796*	1,29	1,03
16006	Igfbp1	0,60	1,02	2,24	0,8266*	0,4983*	0,6048*
100044087	LOC100044087	0,57	NA	NA	NA	NA	NA
100045887	LOC100045887	0,57	NA	NA	NA	NA	NA
100046044	LOC100046044	0,53	NA	NA	NA	NA	NA
100046080	LOC100046080	0,59	NA	NA	NA	NA	NA
100047619	LOC100047619	0,65	NA	NA	NA	0,87	0,92
100047794	LOC100047794	0,66	NA	NA	NA	NA	NA
100048187	LOC100048187	0,60	NA	NA	NA	NA	NA
100048445	LOC100048445	0,66	NA	NA	NA	0,94	1,05
100048622	LOC100048622	0,61	NA	NA	NA	NA	NA
14245	Lpin1	0,54	0,99	1,06	0,5138*	1,14	1,10
70701	Nipal1	0,60	0,3599*	0,8128*	0,98	0,87	1,12
18113	Nnmt	0,41	0,7332*	0,6272*	0,7328*	0,5037*	0,69
83961	Nrg4	0,50	0,3985*	0,93	0,6556*	0,6885*	0,7205*
18194	Nsdhl	0,63	0,3863*	0,9121*	1,11	1,18	1,12
212503	Paox	0,62	1,05	0,97	1,06	1,10	1,03
110208	Pgd	0,66	1,07	0,8929*	1,15	1,20	1,14
628705	Phf11c	0,35	NA	NA	NA	0,89	0,98
68603	Pmvk	0,50	0,6176*	0,8738*	1,13	1,40	1,12
76487	Ppp1r3g	0,43	NA	1,03	1,07	0,5905*	0,87
320717	Pptc7	0,66	0,7002*	NA	NA	1,07	0,93
96875	Prg4	0,62	0,8444*	1,03	1,01	3,50	2,15
67103	Ptgr1	0,62	1,04	0,90	1,05	1,12	0,92
19734	Rgs16	0,63	0,3785*	1,62	1,54	0,93	1,24
68193	Rpl24	0,61	NA	NA	NA	1,34	1,31
20341	Selenbp1	0,65	0,8142*	0,92	0,98	1,08	1,07
72281	Sh2d4a	0,65	1,00	0,94	0,90	0,6742*	0,5954*
171405	Slc22a27	0,48	NA	1,02	1,05	0,6701*	0,4399*
236293	Slc22a29	0,60	0,99	NA	NA	1,07	0,99
20528	Slc2a4	0,65	1,12	0,88	0,97	0,6296*	0,79
20532	Slc3a1	0,63	0,7509*	0,8365*	1,10	2,03	0,97
72401	Slc43a1	0,58	NA	1,05	0,8808*	0,82	0,5474*
100216536	Snord1c	0,64	NA	NA	NA	NA	NA
27211	Snord35a	0,63	NA	NA	NA	0,5887*	0,72
100217454	Snord37	0,65	NA	NA	NA	NA	NA
20860	Sult1e1	0,59	1,07	0,91	0,8534*	0,6242*	0,6738*
629203	Sult2a3	0,50	NA	NA	NA	0,2972*	0,4026*
54610	Tbc1d8	0,65	0,86	1,01	0,8357*	1,02	1,03

Downregulated genes

Entrez ID	gene symbol	FC CR	FC R	FC P	FC Qd	FC Qi	FC Gi
74988	4930480G23Rik	1,81	NA	1,09	1,1384*	NA	NA
78934	4930581F22Rik	1,53	NA	0,99	0,92	NA	NA
70524	5730414N17Rik	1,73	NA	1,07	1,2184*	NA	NA
619326	9130409I23Rik	1,62	NA	1,2095*	0,97	0,97	1,06

230161	Acnat1	2,32	NA	NA	NA	0,72	0,88
26876	Adh4	1,63	1,21	1,1113*	1,16	2,9548*	2,0389*
11656	Alas2	1,63	2,2155*	1,1262*	1,02	1,02	1,5495*
71724	Aox3	1,75	1,4157*	1,4434*	1,04	1,59	1,50
71898	Apol9b	3,36	0,86	NA	NA	0,75	1,12
28194	Apon	1,55	1,06	1,08	1,02	1,20	1,5479*
74008	Arsg	1,77	1,09	1,135*	1,0948*	1,6881*	1,35
12116	Bhmt	1,57	1,5703*	1,04	0,95	0,90	1,07
320926	C730029A08Rik	2,04	NA	1,00	1,4113*	NA	NA
100038579	C730037M02Rik	1,58	NA	NA	NA	NA	NA
230558	C8a	1,76	1,18	1,1542*	0,91	1,9244*	1,52
110382	C8b	1,55	0,97	1,1587*	0,96	1,7269*	1,26
12279	C9	1,85	1,05	1,12	1,01	2,1596*	2,7851*
12350	Car3	1,71	1,654*	1,0056*	1,2897*	0,95	0,97
110794	Cebpe	2,04	NA	NA	NA	0,66	0,82
13897	Ces1e	1,57	1,4903*	1,2295*	1,00	0,75	0,91
382053	Ces3a	1,66	1,1807*	NA	NA	1,6871*	1,5197*
11537	Cfd	1,57	2,0347*	0,92	0,90	1,26	1,4692*
110902	Chrna2	1,83	1,13	1,1781*	1,0565*	0,98	1,17
93673	Cml2	1,82	0,85	1,2088*	0,94	1,15	1,13
246277	Cсад	1,79	1,5578*	1,3108*	0,89	2,0712*	1,35
114564	Csprs	1,93	0,65	NA	NA	1,02	1,08
13009	Csrp3	1,89	0,46	1,00	1,03	2,0498*	1,6468*
627860	Cyp2d37-ps	2,17	NA	NA	NA	NA	NA
71519	Cyp2u1	1,54	NA	1,06	1,11	0,99	1,16
277753	Cyp4a12a	3,26	2,4161*	1,10	0,95	1,7058*	2,4006*
13118	Cyp4a12b	2,28	2,7108*	NA	NA	2,1124*	2,827*
13123	Cyp7b1	1,72	1,4917*	1,05	0,99	2,7255*	1,34
13124	Cyp8b1	1,62	0,77	0,93	0,95	0,74	0,89
320751	D830014E11Rik	1,92	1,14	NA	NA	1,40	1,08
13195	Ddc	1,65	2,2572*	0,89	0,98	1,55	1,41
13370	Dio1	2,14	0,87	0,99	0,97	2,7381*	2,2137*
13423	Dnase2a	1,81	1,9103*	1,0827*	0,75	1,22	1,14
12686	Elov13	2,03	8,3276*	1,0935*	0,86	2,3594*	3,686*
245050	Fam198a	1,61	1,2857*	NA	NA	1,13	1,36
234199	Fgl1	1,51	0,59	0,66	0,82	1,24	1,21
68680	Fitm1	1,90	1,6911*	1,0739*	1,05	1,15	1,08
15377	Foxa3	1,67	0,65	0,95	0,77	1,21	1,4945*
14373	G0s2	1,63	2,3226*	1,1144*	1,4318*	4,0458*	1,7323*
100041085	G630090E17Rik	1,91	NA	NA	NA	NA	NA
626578	Gbp10	2,10	NA	NA	NA	NA	NA
100702	Gbp6	1,82	1,02	1,295*	0,98	1,09	0,97
23886	Gdf15	1,54	1,8197*	2,1827*	1,3954*	2,7155*	0,43
14619	Gjb2	1,60	0,86	0,98	0,99	1,42	1,41
216871	Gltpd2	2,21	1,11	NA	NA	1,23	1,06
100038514	Gm11837	2,40	NA	NA	NA	0,86	0,80
631323	Gm12250	1,75	NA	NA	NA	0,96	0,86
100040465	Gm2788	2,40	NA	0,91	1,2223*	NA	NA
215895	Gm4794	2,14	NA	NA	NA	0,86	0,65
240327	Gm4951	1,63	NA	NA	NA	1,23	1,39

241041	Gm4956	2,59	NA	NA	NA	0,97	1,13
626534	Gm6682	1,66	NA	NA	NA	1,10	0,91
668108	Gm8979	3,14	NA	NA	NA	NA	NA
668139	Gm8995	2,15	0,99	NA	NA	1,17	1,24
14675	Gna14	1,61	1,15	1,1426*	1,1567*	0,74	0,84
72748	Hdhd3	1,71	1,6199*	1,1382*	1,04	1,40	1,17
319165	Hist1h2ad	1,89	NA	NA	NA	NA	NA
319166	Hist1h2ae	1,54	1,23	NA	NA	1,38	1,22
319173	Hist1h2af	1,69	0,94	NA	NA	0,96	0,86
319168	Hist1h2ah	1,81	0,94	NA	NA	1,12	0,93
319169	Hist1h2ak	1,58	NA	NA	NA	0,82	0,75
319170	Hist1h2an	1,73	0,90	NA	NA	0,85	0,77
319171	Hist1h2ap	1,55	NA	NA	NA	NA	NA
15486	Hsd17b2	1,50	1,10	1,1485*	0,99	3,0345*	1,41
15496	Hsd3b5	3,81	5,0466*	1,08	1,14	3,8766*	3,4387*
15505	Hspf1	1,55	2,1185*	1,143*	1,00	0,96	1,531*
15957	Ifit1	2,14	1,01	1,2065*	0,98	2,9278*	3,0535*
15959	Ifit3	1,69	0,87	1,10	1,03	0,97	1,51
667370	Ifit3b	2,08	0,68	NA	NA	0,95	1,16
16145	Igtp	1,69	0,85	1,00	0,95	1,7593*	1,5083*
60440	ligrp1	1,54	0,96	1,1325*	1,01	1,25	1,06
16325	Inhbc	1,96	1,8439*	1,1772*	1,16	1,58	1,44
16326	Inhbe	2,05	1,4144*	1,0898*	1,1534*	1,19	1,12
54396	Irgm2	1,68	0,88	1,00	0,88	2,408*	1,4883*
100038882	Isg15	1,94	NA	NA	NA	2,4079*	3,2528*
64697	Keg1	1,75	1,16	1,03	1,12	3,0228*	2,0137*
100040592	LOC100040592	2,52	NA	NA	NA	NA	NA
100044576	LOC100044576	1,55	NA	NA	NA	1,18	1,07
100047173	LOC100047173	2,06	NA	NA	NA	0,52	0,58
100047200	LOC100047200	1,62	NA	NA	NA	NA	NA
100047427	LOC100047427	2,14	NA	NA	NA	NA	NA
100047651	LOC100047651	1,93	NA	NA	NA	NA	NA
100047674	LOC100047674	1,90	NA	NA	NA	NA	NA
100047707	LOC100047707	1,58	NA	NA	NA	NA	NA
641240	LOC641240	1,90	0,41	NA	NA	NA	NA
17194	Mbl1	1,81	1,20	1,00	1,03	1,9748*	2,049*
59012	Moxd1	1,62	0,91	1,2851*	0,97	0,69	0,82
17837	Mug2	1,67	1,02	NA	NA	1,16	0,88
100039008	Mup10	1,51	NA	1,0621*	1,147*	NA	NA
17841	Mup2	2,50	2,1592*	NA	NA	1,46	1,03
381530	Mup20	4,57	NA	NA	NA	4,5176*	3,4253*
381531	Mup21	3,32	NA	NA	NA	2,4537*	2,0524*
17842	Mup3	1,59	1,314*	0,99	0,85	1,11	1,02
17844	Mup5	1,62	0,97	0,82	0,96	1,38	1,43
620807	Mup6	2,78	NA	NA	NA	2,7742*	1,6157*
68396	Nat8	1,67	1,446*	0,99	0,93	1,65	2,1476*
50490	Nox4	1,61	1,0491*	1,3627*	0,99	1,27	1,34
231655	Oasl1	1,52	1,01	0,87	1,04	1,16	1,5838*
227627	Obp2a	1,96	1,3868*	1,6838*	1,1688*	1,00	1,10
259026	Olfr378	1,59	1,07	NA	NA	1,10	1,34

258590	Olf702	1,57	1,04	NA	NA	0,70	0,85
225631	Onecut2	1,59	1,3412*	1,07	1,11	0,96	0,89
75552	Paqr9	2,24	1,21	0,99	1,2768*	0,88	1,03
18559	Pctp	1,55	0,72	0,93	0,92	0,97	1,05
71830	Pdilt	1,62	2,5484*	1,3829*	1,05	1,15	1,05
75985	Rab30	1,59	1,04	1,177*	1,2221*	1,34	0,89
19369	Raet1d	3,43	NA	NA	NA	NA	NA
109222	Rarres1	1,86	0,88	1,03	1,04	2,2351*	1,6241*
58185	Rsad2	1,56	1,0615*	1,144*	0,97	0,87	0,96
20208	Saa1	2,43	0,45	0,55	0,53	0,95	0,99
20209	Saa2	1,79	0,44	0,28	0,44	0,99	1,07
20211	Saa4	1,74	0,91	1,5963*	0,87	1,7338*	1,30
74032	Sdr42e1	1,57	0,91	1,1449*	0,99	1,6722*	1,19
70061	Sdr9c7	1,80	1,5587*	1,3952*	1,05	1,23	1,13
20342	Selenbp2	2,16	NA	NA	NA	1,08	1,05
68054	Serpina12	2,14	1,2818*	1,1342*	1,04	0,87	0,91
20704	Serpina1e	2,49	0,95	NA	NA	0,82	1,02
238393	Serpina3f	2,69	0,69	NA	NA	0,99	1,01
321018	Serpina4-ps1	3,53	7,8202*	1,3025*	0,70	NA	NA
241877	Slc10a5	1,94	NA	NA	NA	0,89	0,75
319848	Slc17a4	2,49	1,6482*	NA	NA	1,15	1,27
108114	Slc22a7	1,78	1,3457*	1,1246*	1,05	1,7508*	2,0526*
226781	Slc30a10	1,61	NA	1,1206*	1,2085*	1,54	1,31
28248	Slco1a1	2,30	1,13	1,1611*	1,04	2,1771*	1,49
68469	Sox6os	1,78	1,4263*	0,94	1,01	1,12	1,09
78925	Srd5a1	1,66	2,855*	1,1092*	1,1425*	2,1705*	1,20
76971	Sult2a8	1,99	1,34*	1,08	1,01	1,12	1,04
96935	Susd4	1,59	0,93	0,95	1,02	0,55	0,68
21786	Tff3	1,55	0,13	0,87	0,90	0,49	0,28
66039	Tmem254a	2,13	1,04	NA	NA	2,7768*	2,2451*
21807	Tsc22d1	1,52	1,4446*	1,2676*	1,2241*	2,0054*	1,5902*
50500	Ttpa	1,53	0,87	1,03	0,95	4,5041*	2,2968*
71773	Ugt2b1	1,66	1,6716*	1,05	1,00	1,44	1,32
243085	Ugt2b35	1,62	1,2066*	1,1428*	1,02	1,9637*	1,7566*
105887	Ugt3a1	2,13	NA	1,1838*	0,98	2,011*	1,5946*
223337	Ugt3a2	1,53	0,95	1,10	0,97	1,39	1,25
24110	Usp18	1,60	0,72	1,126*	0,92	2,1211*	2,3037*
378472	Vaultrc5	3,99	NA	NA	NA	0,33	0,54
433070	Vmn2r96	1,55	NA	NA	NA	0,93	0,97
327956	Vmo1	1,84	NA	NA	NA	1,29	1,24
22361	Vnn1	2,13	1,02	0,93	1,0863*	0,96	0,88
67701	Wfdc2	1,62	2,2296*	0,95	1,07	0,86	1,36
381246	Xkr9	1,88	NA	NA	NA	1,06	1,01
668923	Zfp442	1,80	NA	NA	NA	1,13	1,05


Q: Overrepresented pathways identified by the regulated CR genes with fold change >1.5


upregulated pathways	downregulated pathways
Phase 1 - Functionalization of compounds - Reactome	Systemic lupus erythematosus - Mus musculus (mouse) - KEGG
Biological oxidations - Reactome	Meiotic Recombination - Reactome
Metabolism - Reactome	nicotine degradation III - MouseCyc
Xenobiotics - Reactome	nicotine degradation II - MouseCyc
Drug metabolism - cytochrome P450 - Mus musculus (mouse) - KEGG	Steroid hormone biosynthesis - Mus musculus (mouse) - KEGG
Metabolism of lipids and lipoproteins - Reactome	Terminal pathway of complement - Reactome
Cytochrome P450 - arranged by substrate type - Reactome	Complement and coagulation cascades - Mus musculus (mouse) - KEGG
Fatty acid biosynthesis - Mus musculus (mouse) - KEGG	Retinol metabolism - Mus musculus (mouse) - KEGG
Triglyceride Biosynthesis - Reactome	Alcoholism - Mus musculus (mouse) - KEGG
Chemical carcinogenesis - Mus musculus (mouse) - KEGG	Phase 1 - Functionalization of compounds - Reactome
Retinol metabolism - Mus musculus - KEGG	Biological oxidations - Reactome
Fatty acid, triacylglycerol, and ketone body metabolism - Reactome	Complement cascade - Reactome
Metabolism of xenobiotics by cytochrome P450 - Mus musculus (mouse) - KEGG	Cytochrome P450 - arranged by substrate type - Reactome
Arachidonic acid metabolism - Mus musculus (mouse) - KEGG	Complement Activation, Classical Pathway - Wikipathways
Fatty Acyl-CoA Biosynthesis - Reactome	Regulation of Complement cascade - Reactome
Glutathione metabolism - Mus musculus (mouse) - KEGG	Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) - Reactome
biotin-carboxyl carrier protein assembly - MouseCyc	Scavenging by Class B Receptors - Reactome
Fatty Acid Biosynthesis - Wikipathways	catecholamine biosynthesis - MouseCyc
Phase II conjugation - Reactome	Metabolism - Reactome
pentose phosphate pathway - MouseCyc	Metabolism of lipids and lipoproteins - Reactome
Metapathway biotransformation - Wikipathways	Glycoprotein hormones - Reactome
Insulin signaling pathway - Mus musculus - KEGG	Prion diseases - Mus musculus (mouse) - KEGG
Pentose Phosphate Pathway - Wikipathways	Bile acid and bile salt metabolism - Reactome
Pentose phosphate pathway (hexose monophosphate shunt) - Reactome	Porphyrin and chlorophyll metabolism - Mus musculus (mouse) - KEGG
nicotine degradation II - MouseCyc	Cytochrome P450 - Wikipathways
Biotin transport and metabolism - Reactome	Peptide hormone biosynthesis - Reactome
pentose phosphate pathway (oxidative branch) - MouseCyc	Drug metabolism - cytochrome P450 - Mus musculus (mouse) - KEGG
Import of palmitoyl-CoA into the mitochondrial matrix - Reactome	Tyrosine metabolism - Mus musculus (mouse) - KEGG
Metabolism of water-soluble vitamins and cofactors - Reactome	Formyl peptide receptors bind formyl peptides and many other ligands - Reactome
Defective TCN2 causes hereditary megaloblastic anemia - Reactome	Fatty acid degradation - Mus musculus (mouse) - KEGG
Cytosolic sulfonation of small molecules - Reactome	Steroid Biosynthesis - Wikipathways
Amino acid synthesis and interconversion (transamination) - Reactome	Advanced glycosylation endproduct receptor signaling - Reactome
Metabolism of amino acids and derivatives - Reactome	DEx/H-box helicases activate type I IFN and inflammatory cytokines production - Reactome
Metabolism of carbohydrates - Reactome	Androgen biosynthesis - Reactome
Metabolism of polyamines - Reactome	Endogenous sterols - Reactome
Defective LMBRD1 causes methylmalonic aciduria and homocystinuria type cbIF - Reactome	Synthesis of bile acids and bile salts via 27-hydroxycholesterol - Reactome
Cholesterol Biosynthesis - Wikipathways	Amine-derived hormones - Reactome


Defective AMN causes hereditary megaloblastic anemia 1 - Reactome	Drug metabolism - other enzymes - <i>Mus musculus</i> (mouse) - KEGG
Defective CUBN causes hereditary megaloblastic anemia 1 - Reactome	Primary bile acid biosynthesis - <i>Mus musculus</i> (mouse) - KEGG
Metabolism of vitamins and cofactors - Reactome	Fatty acids - Reactome
Defective GIF causes intrinsic factor deficiency - Reactome	Complement and Coagulation Cascades - Wikipathways
Defects in cobalamin (B12) metabolism - Reactome	Interferon Signaling - Reactome
Defective MMADHC causes methylmalonic aciduria and homocystinuria type cbID - Reactome	PPAR signaling pathway - <i>Mus musculus</i> (mouse) - KEGG
Defective MTRR causes methylmalonic aciduria and homocystinuria type cbIE - Reactome	Metabolism of steroid hormones and vitamin D - Reactome
Defective MTR causes methylmalonic aciduria and homocystinuria type cbIG - Reactome	Ascorbate and aldarate metabolism - <i>Mus musculus</i> (mouse) - KEGG
Defective MMAB causes methylmalonic aciduria type cbIB - Reactome	Metabolism of xenobiotics by cytochrome P450 - <i>Mus musculus</i> (mouse) - KEGG
Defective MMAA causes methylmalonic aciduria type cbIA - Reactome	Arachidonic acid metabolism - <i>Mus musculus</i> (mouse) - KEGG
Defective MUT causes methylmalonic aciduria mut type - Reactome	Binding and Uptake of Ligands by Scavenger Receptors - Reactome
Defective CD320 causes methylmalonic aciduria - Reactome	ISG15 antiviral mechanism - Reactome
Defective MMACHC causes methylmalonic aciduria and homocystinuria type cbIC - Reactome	Antiviral mechanism by IFN-stimulated genes - Reactome
Defective HLCS causes multiple carboxylase deficiency - Reactome	Metapathway biotransformation - Wikipathways
Defective BTD causes biotidinase deficiency - Reactome	TAK1 activates NFkB by phosphorylation and activation of IKKs complex - Reactome
Defects in biotin (Btn) metabolism - Reactome	Synthesis of bile acids and bile salts - Reactome
Defects in vitamin and cofactor metabolism - Reactome	Metabolism of amino acids and derivatives - Reactome
Cholesterol biosynthesis - Reactome	Amoebiasis - <i>Mus musculus</i> (mouse) - KEGG
superpathway of cholesterol biosynthesis - MouseCyc	PPAR signaling pathway - Wikipathways
glutathione-mediated detoxification - MouseCyc	RIP-mediated NFkB activation via ZBP1 - Reactome
Pentose phosphate pathway - <i>Mus musculus</i> (mouse) - KEGG	Steroid hormones - Reactome
Propanoate metabolism - <i>Mus musculus</i> (mouse) - KEGG	Pentose and glucuronate interconversions - <i>Mus musculus</i> (mouse) - KEGG
Amino acid transport across the plasma membrane - Reactome	ZBP1(DAI) mediated induction of type I IFNs - Reactome
Alanine, aspartate and glutamate metabolism - <i>Mus musculus</i> (mouse) - KEGG	Chemical carcinogenesis - <i>Mus musculus</i> (mouse) - KEGG
bupropion degradation - MouseCyc	TRAF6 mediated NF-kB activation - Reactome
Pyruvate metabolism - <i>Mus musculus</i> (mouse)	Arachidonic acid metabolism - Reactome
nicotine degradation III - MouseCyc	Sulfur amino acid metabolism - Reactome
Glycolysis and Gluconeogenesis - Wikipathways	Peptide hormone metabolism - Reactome
Translation Factors - Wikipathways	bupropion degradation - MouseCyc
Amino acid and oligopeptide SLC transporters - Reactome	Glycine, serine and threonine metabolism - <i>Mus musculus</i> (mouse) - KEGG
Glucose metabolism - Reactome	
Adipocytokine signaling pathway - <i>Mus musculus</i> (mouse) - KEGG	
Arachidonic acid metabolism - Reactome	

R: Overlap between the upregulated CR genes (*q*-value < 0.05) based on the entrez IDs which are investigated by all dietary interventions (15552 IDs)
 (black squares indicate that the gene is regulated by the phytonutrient as well)

		R	P	Q	RP	RQ	PQ	RPQ		R	P	Q	RP	RQ	PQ	RPQ
58520	0610007P14Rik								26408	Map3k5						
66117	1110001J03Rik								53608	Map3k6						
68480	1110007C09Rik								29857	Mapk12						
68618	1110012L19Rik								69546	Mapk1ip1						
68531	1110020A21Rik								212307	Mapre2						
73747	1110034G24Rik								223989	Marf1						
68861	1190002N15Rik								546071	Mast3						
67885	1500011K16Rik								108645	Mat2b						
73287	1700040L02Rik								110962	Mbd6						
69066	1810010H24Rik								17195	Mbl2						
66293	1810032O08Rik								84004	Mcam						
69809	1810046K07Rik								72039	Mccc1						
72056	1810055G02Rik								17210	Mcl1						
72282	1810062G17Rik								94178	Mcoln1						
66439	2010012O05Rik								17448	Mdh2						
70257	2010107E04Rik								28077	Med10						
381792	2310040G24Rik								59024	Med12						
69657	2310047D07Rik								75613	Med25						
69662	2310061I04Rik								69790	Med30						
73667	2410004P03Rik								67381	Med4						
77034	2510039O18Rik								17289	Mertk						
67513	2610002J02Rik								70152	Mettl7a1						
77014	2700079J08Rik								73822	Mfsd12						
73205	3110043O21Rik								76574	Mfsd2a						
109292	4631423B10Rik								232714	Mgam						
67392	4833420G17Rik								17313	Mgp						
74430	4930452B06Rik								66447	Mgst3						
75255	4930562F07Rik								27008	Micall1						
78817	4930588G05Rik								68041	Mid1ip1						
319582	6430573F11Rik								28010	Miip						
213393	8430408G22Rik								17347	Mknk2						
74525	8430419L09Rik								70291	Mkrn2os						
319977	A530079E22Rik								21428	Mlx						
433752	AA415398								208104	Mlxip						
78894	Aacs								109129	Mmadhc						
67758	Adadac								75104	Mmd2						
239435	Aard								17393	Mmp7						
30956	Aass								72199	Mms19						
27410	Abca3								93734	Mpv17l						
217258	Abca8a								234384	Mpv17l2						
27413	Abcb11								15064	Mr1						
18671	Abcb1a								17533	Mrc1						
239273	Abcc4								18100	Mrpl40						
27421	Abcc6								52443	Mrpl48						
11307	Abcg1								66230	Mrps28						
27409	Abcg5								232536	Mrps35						
67470	Abcg8								66128	Mrps36						
76491	Abhd14b								17698	Msn						








101565	Ccp110		18950	Pnp	
12475	Cd14		66853	Pnpla2	
93671	Cd163		116939	Pnpla3	
69142	Cd209f		75772	Pnpla5	
13136	Cd55		242608	Podn	
12516	Cd7		27205	Podxl	
12527	Cd9		69745	Pold4	
673094	Cd99		18973	Pole	
218294	Cdc14b		50776	Polg2	
12531	Cdc25b		67710	Polr2g	
240505	Cdc42bpg		245841	Polr2h	
104252	Cdc42ep2		74414	Polr3c	
58804	Cdc42ep5		107939	Pom121	
109332	Cdcp1		18984	Por	
66626	Cdip1		50706	Postn	
12534	Cdk1		74776	Ppa2	
18557	Cdk18		19012	Ppap2a	
66971	Cdk5rap1		19017	Ppargc1a	
264064	Cdk8		170826	Ppargc1b	
12577	Cdkn1c		231327	Ppat	
12580	Cdkn2c		19035	Ppib	
67177	Cdt1		19038	Ppic	
26365	Ceacam1		105675	Ppif	
12608	Cebpb		227399	Ppip5k2	
13706	Cela2a		19041	Ppl	
12614	Celsr1		26930	Ppnr	
12615	Cenpa		17872	Ppp1r15a	
83815	Cenpq		19057	Ppp3cc	
99100	Cep152		54637	Praf2	
241447	Cers6		21672	Prdx2	
72361	Ces2g		11757	Prdx3	
26370	Cetn2		77619	Prelid2	
75472	Cfap126		96875	Prg4	
14894	Cfap20		19089	Prkcsh	
12631	Cfl1		19125	Prodh	
69065	Chac1		19126	Prom1	
72170	Chchd4		19128	Pros1	
66433	Chchd7		114863	Prosc	
12648	Chd1		192159	Prpf8	
68058	Chd1l		19139	Prps1	
244059	Chd2		67763	Prpsap1	
320790	Chd7		75210	Prr3	
74277	Chic2		330657	Prss53	
68038	Chid1		76560	Prss8	
68953	Chmp2a		107272	Psat1	
11438	Chrna4		19171	Psmb10	
234421	Cib3		26445	Psmb2	
71722	Cic		67151	Psmd9	
12696	Cirbp		19188	Psme2	
68379	Ciz1		100678	Psph	
12747	Clk1		19201	PstPIP2	
12750	Clk4		19215	Ptgds	


94040	Clmn			19216	Ptger1	
69634	Clybl			67103	Ptgr1	
76469	Cmya5			545622	Ptpn3	
71994	Cnn3			110095	Pygl	
69171	Cnppd1			110391	Qdpr	
226744	Cnst			60507	Qtrt1	
26920	Cntrl			19329	Rab17	
67892	Coa6			19332	Rab20	
319876	Cobll1			19335	Rab23	
12825	Col3a1			19339	Rab3a	
12827	Col4a2			69834	Rab43	
12833	Col6a1			19341	Rab4a	
12834	Col6a2			56382	Rab9	
17846	Commd1			70314	Rabep2	
69456	Commd10			19352	Rabggtb	
66199	Commd4			19355	Rad1	
59042	Cope			241694	Ralgapa2	
66423	Coprs			19730	Ralgds	
67876	Coq10b			54409	Ramp2	
217707	Coq6			74334	Ranbp10	
12850	Coq7			223332	Ranbp3l	
12865	Cox7a1			56705	Ranbp9	
70574	Cpm			57785	Rangrf	
234577	Cpne2			56508	Rapgef4	
54188	Cpsf4			217944	Rapgef5	
12946	Cr1l			218772	Rarb	
12909	Crcp			19414	Rasa3	
12925	Crip1			320292	Rasgef1b	
56693	Crtap			19395	Rasgrp2	
12952	Cry1			192678	Rassf3	
12953	Cry2			54354	Rassf5	
12962	Crybb3			73246	Rassf6	
68631	Cryl1			56275	Rbm14	
12971	Crym			229700	Rbm15	
104318	Csnk1d			381626	Rbm33	
70425	Csnk1g3			71684	Rbm43	
13007	Csrp1			19659	Rbp1	
13008	Csrp2			71330	Rcbtb1	
14219	Ctgf			17252	Rdh11	
107869	Cth			105014	Rdh14	
13034	Ctse			19682	Rdh5	
13038	Ctsk			103142	Rdh9	
13039	Ctsl			53614	Reck	
13041	Ctsw			72549	Reep4	
13048	Cux2			68703	Rere	
72017	Cyb5r1			232441	Rerg	
109754	Cyb5r3			19733	Rgn	
66445	Cyc1			19734	Rgs16	
13074	Cyp17a1			19736	Rgs4	
77951	Cyp20a1			69159	Rhebl1	
13088	Cyp2b10			69288	Rhobtb1	
13089	Cyp2b13			11854	Rhod	


13094	Cyp2b9		52428	Rhpn2	
13097	Cyp2c38		72388	Ripk4	
13098	Cyp2c39		68477	Rmnd5a	
226143	Cyp2c44		58809	Rnase4	
56448	Cyp2d22		67153	Rnaseh2b	
13108	Cyp2g1		223881	Rnd1	
74519	Cyp2j9		11858	Rnd2	
244209	Cyp2r1		74315	Rnf145	
56050	Cyp39a1		75234	Rnf19b	
13112	Cyp3a11		105239	Rnf44	
13113	Cyp3a13		252967	Ropn1l	
64385	Cyp4f14		19885	Rorc	
13121	Cyp51		68240	Rpa3	
225995	D030056L22Rik		71919	Rpap3	
210108	D130043K22Rik		19892	Rpe65	
52392	D1Ert622e		19895	Rpia	
65111	Dap3		67115	Rpl14	
69635	Dapk1		68028	Rpl22l1	
70248	Dazap1		20103	Rps5	
13171	Dbt		20130	Rras	
66686	Dcbld1		20133	Rrm1	
245038	Dclk3		20135	Rrm2	
13191	Dctn1		27981	Rsrp1	
66422	Dctpp1		20166	Rtnk	
13194	Ddb1		242819	Rundc3b	
107986	Ddb2		20196	S100a13	
72108	Ddh2		20200	S100a6	
13200	Ddost		224903	Safb	
13202	Ddt		224902	Safb2	
13214	Defb1		58198	Sall1	
74754	Dhcr24		68653	Samm50	
13360	Dhcr7		269003	Sap130	
66375	Dhrs7		104175	Sbk1	
69192	Dhx16		107767	Scamp1	
71723	Dhx34		20778	Scarb1	
330938	Dixdc1		20249	Scd1	
235339	Dlat		20266	Scn1b	
53310	Dlg3		74617	Scpep1	
13388	Dll1		64136	Sdf2l1	
235380	Dmxl2		68002	Sdhaf4	
23950	Dnajb6		67680	Sdhb	
30045	Dnajc12		66925	Sdhd	
235567	Dnajc13		231691	Sds	
71972	Dnmbp		257635	Sdsl	
21673	Dntt		74136	Sec14l1	
319899	Dock6		20334	Sec23a	
208266	Dot1l		99683	Sec24b	
93838	Dqx1		69162	Sec31a	
13506	Dsc2		66212	Sec61b	
56405	Dusp14		20341	Selenbp1	
235661	Dync1li1		80795	Selk	
110350	Dync2h1		223776	Selo	


213575	Dync2li1			20351	Sema4a		
56455	Dynll1			26456	Sema4g		
103551	E130012A19Rik			93684	Sep15		
106389	Eaf2			20362	Sept8		
13595	Ebp			12401	Serpina6		
52665	Echdc1			140742	Sesn1		
52430	Echdc2			75747	Sesn3		
68545	Ecscr			208043	Setd1b		
192193	Edem1			224440	Setd4		
13614	Edn1			20390	Sftp4d		
13631	Eef2k			14057	Sfxn1		
81701	Egfl8			20393	Sgk1		
77683	Ehmt1			16923	Sh2b3		
58521	Eid1			27387	Sh2d3c		
13685	Eif4ebp1			72281	Sh2d4a		
54326	Elov12			57028	Sh3bp1		
170439	Elov16			66938	Sh3d21		
59308	Emcn			58234	Shank3		
72205	Eml2			20437	Siah1a		
13731	Emp2			20439	Siah2		
13803	Enc1			17691	Sik1		
13808	Eno3			57738	Slc15a2		
13809	Enpep			100561	Slc15a4		
18605	Enpp1			72472	Slc16a10		
12496	Entpd2			217316	Slc16a5		
13819	Epas1			76306	Slc18b1		
269587	Epb4.1			20509	Slc19a1		
17075	Epcam			116914	Slc19a2		
1E+08	Epg5			20515	Slc20a1		
13842	Epha8			70840	Slc22a16		
13846	Ephb4			18400	Slc22a18		
223650	Eppk1			236149	Slc22a26		
13867	Erbb3			20522	Slc23a1		
170942	Erdr1			54338	Slc23a2		
74155	Errfi1			13358	Slc25a1		
69524	Esam			67863	Slc25a11		
26379	Esrra			78830	Slc25a12		
214253	Etnk2			20523	Slc25a14		
71760	Etnppl			18408	Slc25a15		
14058	F10			68267	Slc25a22		
14068	F7			214663	Slc25a29		
16592	Fabp5			69906	Slc25a32		
76267	Fads1			104910	Slc25a47		
56473	Fads2			230125	Slc25a51		
104943	Fam110c			26457	Slc27a1		
215512	Fam117a			63959	Slc29a1		
218236	Fam120a			20526	Slc2a2		
84652	Fam126a			20528	Slc2a4		
66270	Fam134b			56485	Slc2a5		
58909	Fam13a			56017	Slc2a8		
230766	Fam167b			22782	Slc30a1		
320159	Fam179a			108652	Slc35b3		


212483	Fam193b		70484	Slc35d2	
218850	Fam208a		14385	Slc37a4	
27999	Fam3c	[solid black]	67760	Slc38a2	
77938	Fam53b		76257	Slc38a3	
56306	Fam60a		69354	Slc38a4	
235461	Fam63b	[solid black]	72027	Slc39a4	
72826	Fam76b	[solid black]	20532	Slc3a1	
69955	Fars2	[solid black]	213603	Slc44a3	
14104	Fasn		71706	Slc46a3	
320720	Fastkd1		330962	Slc51b	
69577	Fastkd3		330064	Slc5a6	
217335	Fbf1	[solid black]	14411	Slc6a12	
269514	Fbxl4		102857	Slc6a8	
50755	Fbxo18		11988	Slc7a2	
231670	Fbxo21	[solid black]	224022	Slc7a4	
76454	Fbxo31	[solid black]	170756	Slc8b1	
70611	Fbxo33		28250	Slco1a4	
71538	Fbxo9		20568	Slpi	
215384	Fcgbp	[solid black]	55994	Smad9	
14130	Fcgr2b	[solid black]	69780	Smap2	
207278	Fchsd2		67155	Smarca2	
14137	Fdft1		226026	Smc5	
110196	Fdps	[solid black]	69029	Smdt1	
14149	Fdxr		64074	Smoc2	
14151	Fech		29856	Smtn	
14159	Fes	[solid black]	20613	Snai1	
13998	Fgd6		20583	Snai2	
14194	Fh1		319317	Snhg11	
14198	Fhit		20623	Snrk	
231997	Fkbp14		78372	Snrnp25	
14229	Fkbp5	[solid black]	20637	Snrnp70	
216805	Flcn		68981	Snrpa1	
14251	Flot1	[solid black]	20646	Snrpn	
14257	Flt4		268534	Sntg2	
14260	Fmn1		55988	Snx12	
55990	Fmo2	[solid black]	244962	Snx14	
14262	Fmo3		67804	Snx2	
226564	Fmo4	[solid black]	223920	Soat2	
14263	Fmo5		216233	Socs2	
15375	Foxa1	[solid black]	20657	Sod3	
56458	Foxo1		20410	Sorbs3	
212398	Frat2	[solid black]	20671	Sox17	
232288	Frmd4b		20672	Sox18	
14313	Fst	[solid black]	20692	Sparc	
14317	Ftcd		13602	Sparcl1	
69064	Fuom	[solid black]	219140	Spata13	
18550	Furin		78779	Spata2l	
14366	Fzd4		67629	Spc24	
14367	Fzd5		66442	Spc25	
14381	G6pdx		71026	Speer3	
14388	Gab1		74288	Spem1	
53897	Gal3st1	[solid black]	56381	Spen	













S: Overlap between the downregulated CR genes (*q*-value < 0.05) based on the entrez IDs which are investigated by all dietary interventions (15552 IDs)

(black squares indicate that the gene is regulated by the phytonutrient as well)



67618	Aasdhppt			26403	Map3k11		
56321	Aatf			245877	Map7d1		
11302	Aatk			332110	Mapk15		
268860	Abat			218975	Mapk1ip1l		
11303	Abca1			17165	Mapkapk5		
11305	Abca2			71779	March8		
18670	Abcb4			17167	Marco		
12780	Abcc2			232087	Mat2a		
76408	Abcc3			17184	Matr3		
11666	Abcd1			17192	Mbd3		
24015	Abce1			17194	Mbl1		
27406	Abcf3			193813	Mcfd2		
26357	Abcg2			66405	Mcts2		
57742	Abhd1			16543	Mdfic		
226016	Abhd17b			108098	Med21		
54608	Abhd2			230316	Megf9		
106861	Abhd3			104362	Meig1		
66082	Abhd6			75178	Meiob		
11308	Abi1			17299	Mettl1		
319713	Ablim3			72096	Mettl10		
113868	Acaa1a			74319	Mettl23		
235674	Acaa1b			75422	Mettl5		
102632	Acad11			69572	Mfsd3		
229211	Acad9			23945	Mgll		
11364	Acadm			17314	Mgmt		
110446	Acat1			338320	Mia2		
67131	Acbd4			338366	Mia3		
74159	Acbd5			218613	Mier3		
209186	Acnat2			69674	Mif4gd		
171210	Acot2			109154	Mlec		
171282	Acot4			58805	Mlxipl		
170789	Acot8			67468	Mmd		
93732	Acox2			17380	Mme		
11433	Acp5			236792	Mmgt1		
264895	Acsf2			17388	Mmp15		
14081	Acsf1			58223	Mmp19		
20216	Acsm3			19070	Mob4		
272428	Acsm5			17434	Mocs2		
11461	Actb			68393	Mogat1		
11465	Actg1			72825	Mon1a		
11472	Actn2			56397	Morf4l2		
71670	Acy3			68929	Mospd3		
11488	Adam11			59012	Moxd1		
11491	Adam17			55951	Mpc1		
58860	Adamdec1			67973	Mphosph10		
216991	Adap2			66308	Mplkip		
56417	Adar			56524	Mpp6		
268822	Adck5			246221	Mpst		
52614	Adgre4			17527	Mpv17		
54672	Adgrg3			14012	Mpzl2		
110789	Adgrv1			77037	Mrap		
26876	Adh4			107732	Mrpl10		




68839	Ankrd46		66208	Nenf	
56503	Ankrd49		18021	Nfatc3	
99696	Ankrd50		18023	Nfe2l1	
72074	Anks4b		18024	Nfe2l2	
11737	Anp32a		18029	Nfic	
67628	Anp32b		18030	Nfil3	
66471	Anp32e		18032	Nfix	
16790	Anpep		105193	Nhlrc1	
11761	Aox1		66257	Nicn1	
71724	Aox3		67949	Nifk	
11772	Ap2a2		18081	Ninj1	
11777	Ap3s1		93790	Nipa2	
57267	Apba3		66536	Nipsnap3b	
11808	Apoa4		69721	Nkiras1	
238055	Apob		75805	Nln	
11810	Apobec1		101613	Nlrp6	
75761	Apol7a		270151	Nlrx1	
28194	Apon		64685	Nmi	
66408	Apxt		70769	Nolc1	
11829	Aqp4		55989	Nop58	
64008	Aqp9		50490	Nox4	
11835	Ar		217365	Nploc4	
68497	Arel1		230103	Npr2	
11841	Arf2		13864	Nr2f6	
228998	Arfgap1		26424	Nr5a2	
11855	Arhgap5		192292	Nrbp1	
213649	Arhgef19		18186	Nrp1	
11861	Arl4a		18193	Nsd1	
75423	Arl5a		386649	Nsfl1c	
54208	Arl6ip1		56876	Nsmf	
11865	Arntl		319638	Nt5dc1	
59046	Arpp19		70021	Nt5dc2	
11883	Arsa		74137	Nuak2	
74008	Arsg		26426	Nubp2	
142688	Asb13		52653	Nudcd2	
65256	Asb2		17766	Nudt1	
77987	Ascc3		67993	Nudt12	
58170	Asic5		66911	Nudt16l1	
70396	Asnsd1		213484	Nudt18	
11484	Aspa		67528	Nudt7	
320817	Atad2b		66387	Nudt8	
107503	Atf5		68564	Nufip2	
226641	Atf6		71805	Nup93	
12915	Atf6b		78252	Nxpe2	
11793	Atg5		237082	Nxt2	
74244	Atg7		231655	Oasl1	
212974	Athl1		227627	Obp2a	
71093	Atoh8		77090	Ocel1	
320940	Atp11c		70113	Odfl3b	
11937	Atp2a1		72075	Ogrfr	
11964	Atp6v1a		18295	Ogn	
11981	Atp9a		50914	Olig1	

230649	Atpaf1		27047	Omd	
235610	Atrip		15379	Onecut1	
20239	Atxn2		225631	Onecut2	
54140	Avpr1a		403187	Opa3	
54375	Azin1		75475	Oplah	
26877	B3galt1		26428	Orc4	
72727	B3gat3		227102	Ormdl1	
101148	B630005N14Rik		71720	Osbpl3	
12012	Baat		79196	Osbpl5	
23821	Bace1		237542	Osbpl8	
12013	Bach1		71839	Osgin1	
12017	Bag1		72201	Otud6b	
29810	Bag3		18813	Pa2g4	
224727	Bag6		18458	Pabpc1	
12018	Bak1		18472	Pafah1b1	
170442	Bbox1		67869	Paip2	
72569	Bbs5		72333	Palld	
80748	BC004004		214627	Papd5	
227707	BC005624		18789	Papola	
667597	BC023105		231474	Paqr3	
223631	BC025446		71904	Paqr7	
12033	Bcap29		75552	Paqr9	
12927	Bcar1		58220	Pard6b	
12041	Bckdk		26430	Parg	
12051	Bcl3		243771	Parp12	
71911	Bdh1		235587	Parp3	
12116	Bhmt		80285	Parp9	
64918	Bhmt2		67683	Pbdc1	
12122	Bid		67307	Pbld2	
53619	Blcap		68371	Pbld2	
12151	Bmi1		66923	Pbrm1	
12153	Bmp1		23983	Pcbp1	
12159	Bmp4		59092	Pcbp4	
12162	Bmp7		75599	Pcdh1	
12183	Bpgm		18553	Pcsk6	
14312	Brd2		18559	Pctp	
76809	Bri3bp		18563	Pcx	
12215	Bsg		13026	Pcyt1a	
107182	Btaf1		18567	Pdcd2	
68815	Btbd10		67466	Pdcl	
26363	Btd		211948	Pde12	
53414	Bysl		18582	Pde6d	
57316	C1d		14827	Pdia3	
94192	C1galt1		12304	Pdia4	
59048	C1galt1c1		71853	Pdia6	
12259	C1qa		71830	Pdilt	
12260	C1qb		56376	Pdlim5	
12262	C1qc		319518	Pdpr	
50908	C1s1		59020	Pdzk1	
77644	C330007P06Rik		105083	Pelo	
12269	C4bp		64058	Perp	
12274	C6		64934	Pes1	

230558	C8a					18631	Pex11a		
110382	C8b					72129	Pex13		
69379	C8g					19298	Pex19		
12279	C9					74043	Pex26		
67770	Caap1					67199	Pfdn1		
12283	Cab39					18645	Pfn2		
12301	Cacybp					72542	Pgam5		
69719	Cad					241062	Pgap1		
54725	Cadm1					233575	Pgap2		
260299	Cadm4					57757	Pglyrp2		
109624	Cald1					109785	Pgm3		
140904	Calm1					66522	Pgpep1		
12317	Calr					239510	Phf20l1		
12330	Canx					78246	Phf23		
12333	Capn1					102093	Phkb		
12334	Capn2					21664	Phlda1		
12339	Capn7					208177	Phldb2		
53872	Caprin1					233489	Picalm		
12340	Capza1					98496	Pid1		
12343	Capza2					327942	Pigl		
12350	Car3					78928	Pigt		
12319	Car8					66268	Pigyl		
213819	Casd1					18708	Pik3r1		
12370	Casp8					18710	Pik3r3		
26885	Casp8ap2					170741	Pilrb1		
12389	Cav1					18712	Pim1		
104836	Cbl11					72400	Pinx1		
212139	Cc2d1a					56305	Pitpnb		
75565	Ccdc101					106522	Pkdcc		
69668	Ccdc115					18771	Pknox1		
104479	Ccdc117					227937	Pkp4		
108811	Ccdc122					66350	Pla2g12a		
57895	Ccdc126					69836	Pla2g12b		
67119	Ccdc159					192654	Pla2g15		
52713	Ccdc59					53357	Pla2g6		
108673	Ccdc86					18786	Plaa		
66365	Ccdc90b					104759	Pld4		
20304	Ccl5					27276	Plekhb1		
20305	Ccl6					72287	Plekhf1		
216527	Ccm2					11520	Plin2		
12450	Ccng1					66968	Plin5		
12453	Ccni					67801	Pllp		
12455	Ccnt1					53317	Plrg1		
12460	Ccs					18830	Pltp		
12479	Cd1d1					18845	Plxna2		
246278	Cd207					29858	Pmm1		
12484	Cd24a					56695	Pnkd		
60533	Cd274					59047	Pnkp		
12488	Cd2ap					18949	Pnn		
217305	Cd300ld					66249	Pno1		
66205	Cd302					241274	Pnpla7		
12491	Cd36					103711	Pnpo		

70333	Cd3eap				18970	Polb		
12506	Cd48	■	■	■	66979	Pole4		
23833	Cd52				64424	Polr1e		
12508	Cd53				69920	Polr2i		
11801	Cd5l	■		■	235036	Ppan		
12514	Cd68				67916	Ppap2b		
16149	Cd74	■			19016	Pparg		
12520	Cd81				66496	Ppdpf		
12522	Cd83				223828	Pphln1		
12524	Cd86				67905	Ppm1m		
98828	Cdc123	■	■	■	19046	Ppp1cb		
216150	Cdc34	■	■	■	76497	Ppp1r11		
12558	Cdh2	■	■	■	17931	Ppp1r12a		
72040	Cdhr5				68458	Ppp1r14a		
52858	Cdip1t				244416	Ppp1r3b		
12575	Cdkn1a	■	■	■	53412	Ppp1r3c		
12583	Cdo1	■	■	■	243725	Ppp1r9a		
12606	Cebpa				19052	Ppp2ca		
109901	Cela1	■	■	■	235542	Ppp2r3a		
12616	Cenpb	■	■	■	226849	Ppp2r5a		
73139	Cenpv				26931	Ppp2r5c		
99712	Cept1				26932	Ppp2r5e		
104158	Ces1d				70351	Ppp4r1		
13897	Ces1e	■	■	■	232314	Ppp4r2		
234564	Ces1f	■	■	■	67857	Ppp6c		
12623	Ces1g	■	■	■	226169	Pprc1		
102022	Ces2a	■	■	■	110593	Prdm2		
216618	Cfap36	■	■	■	11758	Prdx6		
11537	Cfd	■	■	■	116847	Prelp		
23837	Cfdp1	■	■	■	213760	Prepl		
12628	Cfh				18747	Prkaca		
50702	Cfhr1				75292	Prkd3		
12633	Cflar	■	■	■	72981	Prkrir		
18636	Cfp				19116	Prlr		
68567	Cgref1				15469	Prmt1		
68044	Chac2	■	■	■	67939	Prorsd1		
107932	Chd4				56194	Prpf40a		
109151	Chd9				66151	Prr13		
218865	Chdh	■	■	■	19152	Prtn3		
12660	Chka				19156	Psap		
234852	Chmp1a				19165	Psen2		
75608	Chmp4b				26442	Psma5		
69993	Chn2	■	■	■	16913	Psmb8		
66917	Chordc1				16912	Psmb9		
100910	Chpf2				19181	Psmc2		
212862	Chpt1	■	■	■	23996	Psmc4		
110902	Chrna2				67089	Psmc6		
66511	Chtop				66997	Psmd12		
12683	Cidea				59029	Psmd14		
12684	Cideb				19185	Psmd4		
14311	Cidec				66413	Psmd6		
54124	Cks1b				19192	Psme3		



208677	Creb3l3		19692	Reg1	
233490	Crebzf		100532	Rell1	
433375	Creg1		67442	Retsat	
76737	Creld2		54391	Rfk	
68337	Crip2		242406	Rgp1	
66586	Crls1		50780	Rgs3	
66877	Crnkl1		214951	Rhbdl1	
74114	Crot		58176	Rhbg	
12944	Crp		19744	Rheb	
214301	Crygn		69581	Rhou	
12972	Cryz		217835	Rin3	
246277	Cсад		93726	Rnase2a	
229663	Csde1		29864	Rnf11	
110750	Cse1l		75212	Rnf121	
78752	Csgalnact2		71956	Rnf135	
211660	Cspp1		56515	Rnf138	
13009	Csrp3		66825	Rnf186	
13010	Cst3		268749	Rnf31	
83410	Cstf2t		207742	Rnf43	
13016	Ctbp1		19823	Rnf7	
13018	Ctcf		76892	Rnft1	
67181	Ctdnep1		215615	Rnpep	
67655	Ctdp1		19877	Rock1	
227292	Ctdsp1		66049	Rogdi	
66642	Ctnnbl1		19934	Rpl22	
51797	Ctps		66229	Rpl7l1	
13040	Ctss		67053	Rpp14	
64138	Ctsz		208366	Rpp40	
71745	Cul2		68441	Rraga	
26554	Cul3		66922	Rras2	
75717	Cul5		81910	Rrbp1	
244672	Cwf19l2		107094	Rrp12	
14825	Cxcl1		224823	Rrp36	
15945	Cxcl10		101867	Rrp8	
55985	Cxcl13		59014	Rrs1	
13056	Cyb561		58185	Rsd2	
66427	Cyb5b		170728	Rtn4ip1	
230582	Cyb5rl		237847	Rtn4rl1	
13058	Cybb		235636	Rtp3	
13095	Cyp2c29		67775	Rtp4	
13096	Cyp2c37		216724	Rufy1	
107141	Cyp2c50		192174	Rwdd4a	
404195	Cyp2c54		20194	S100a10	
68444	Cyp2d13		20201	S100a8	
13105	Cyp2d9		20202	S100a9	
13107	Cyp2f2		13609	S1pr1	
13109	Cyp2j5		94226	S1pr5	
13110	Cyp2j6		20208	Saa1	
13116	Cyp46a1		20209	Saa2	
277753	Cyp4a12a		20210	Saa3	
102294	Cyp4v3		20211	Saa4	
13123	Cyp7b1		209086	Samd9l	

13124	Cyp8b1		72193	Scaf11	
16007	Cyr61	[■]	24044	Scamp2	
242484	D630039A03Rik	[■■■■■]	56807	Scamp5	
208846	Daam1	[■■■■■]	71145	Scara5	[■]
13143	Dapk2		56367	Scoc	[■]
13163	Daxx		240880	Scyl3	[■]
83703	Dbr1		20969	Sdc1	
242418	Dcaf10	[■]	20971	Sdc4	[■]
28199	Dcaf11	[■■■]	68332	Sdhaf1	
212123	Dcaf15	[■■■]	74032	Sdr42e1	[■■■]
98193	Dcaf8		70061	Sdr9c7	[■]
68087	Dcakd	[■■■]	67815	Sec14l2	[■■■]
13179	Dcn		20333	Sec22b	
13190	Dct	[■]	27054	Sec23b	[■]
114893	Dcun1d1		69608	Sec24d	[■]
100737	Dcun1d4		69276	Sec62	
13195	Ddc	[■■■]	140740	Sec63	
77591	Ddx10	[■■■]	75420	Secisbp2	
67040	Ddx17	[■■■]	70354	Secisbp2l	[■■■]
56200	Ddx21	[■■■]	114679	Selm	
71986	Ddx28	[■■■]	20345	Selplg	
13205	Ddx3x	[■■■■■]	320213	Senp5	
26900	Ddx3y	[■■■■■]	109079	Sephs1	
78394	Ddx52	[■■■■■]	235072	Sept7	
13209	Ddx6	[■■■■■]	53860	Sept9	
234311	Ddx60	[■■■■■]	20364	Sepw1	[■]
67460	Decr1	[■■■■■]	66870	Serb1	[■]
26378	Decr2	[■■■■■]	56442	Serinc1	[■]
23854	Def8		230779	Serinc2	
329877	Dennd4c	[■■■■■]	218442	Serinc5	
70377	Derl3	[■■■■■]	217847	Serpina10	
58239	Dexi	[■■■■■]	380780	Serpina11	
71755	Dhdh	[■■■■■]	68054	Serpina12	[■■■■■]
70451	Dhrs13	[■■■■■]	74069	Serpina3a	[■■■■■]
20148	Dhrs3	[■■■■■]	20716	Serpina3n	
28200	Dhrs4	[■■■■■]	321018	Serpina4-ps1	[■]
216820	Dhrs7b	[■■■■■]	71907	Serpina9	[■■■■■]
13204	Dhx15	[■■■■■]	66222	Serpib1a	
218629	Dhx29	[■■■■■]	20708	Serpib6b	
72162	Dhx36	[■■■■■]	18787	Serpine1	
80861	Dhx58	[■■■■■]	12258	Serpingle1	
13370	Dio1		81898	Sf3b1	
72662	Dis3	[■■■■■]	66055	Sf3b6	
234130	Dkk4		67788	Sfr1	
74129	Dmgdh		27219	Sgk2	
68922	Dnaic1		74442	Sgms2	
15502	Dnaja1	[■■■■■]	20397	Sgpl1	
81489	Dnajb1	[■■■■■]	81535	Sgpp1	
67838	Dnajb11	[■■■■■]	56726	Sh3bgrl	
70604	Dnajb14	[■■■■■]	212531	Sh3bgrl2	
56812	Dnajb2	[■■■■■]	24055	Sh3bp2	
67035	Dnajb4	[■■■■■]	56392	Shoc2	



13682	Eif4a2					28253	Slco1b2			
75705	Eif4b					380773	Slrp			
13684	Eif4e					66390	Slmo2			
208643	Eif4g1					207818	Smagp			
217869	Eif5					20586	Smarca4			
226982	Eif5b					54380	Smarcal1			
244548	Elmod2					20588	Smarcc1			
12686	Elovl3					68094	Smarcc2			
68801	Elovl5					70099	Smc4			
193838	Eme2					170748	Smco4			
71946	Endod1					104570	Smek2			
209558	Enpp3					226517	Smg7			
224794	Enpp4					68552	Smim14			
83965	Enpp5					66487	Smim4			
12499	Entpd5					20595	Smn1			
223527	Eny2					319757	Smo			
227867	Epc2					66313	Smurf2			
13836	Epha2					319822	Smyd4			
13849	Ephx1					20619	Snap23			
13850	Ephx2					67826	Snap47			
77781	Epm2aip1					66618	Snrnp27			
13854	Epn1					76167	Snrnp35			
98845	Eps8l2					20643	Snrpe			
28042	Ept1					20649	Sntb1			
67458	Ergic1					266781	Snx17			
67456	Ergic2					102607	Snx19			
226090	Ermp1					69226	Snx24			
66580	Esf1					12702	Socs3			
77411	Esrp2					68469	Sox6os			
66841	Etfdh					20683	Sp1			
232146	Eva1a					20687	Sp3			
14020	Evi5					64406	Sp5			
66482	Exoc2					50850	Spast			
211446	Exoc3					66624	Spcs2			
20336	Exoc4					229285	Spg20			
14071	F9					20729	Spin1			
14079	Fabp2					100689	Spon2			
12140	Fabp7					20747	Spop			
68636	Fahd1					76857	Spopl			
23873	Faim					66674	Spryd7			
98952	Fam102a					79043	Spsb3			
66540	Fam107b					211949	Spsb4			
213056	Fam126b					18412	Sqstm1			
227737	Fam129b					78925	Srd5a1			
225358	Fam13b					19073	Srgn			
212326	Fam149a					20810	Srm			
105428	Fam149b					217337	Srp68			
73942	Fam151b					20818	Srprb			
319604	Fam168a					27364	Srr			
106014	Fam19a5					110809	Srsf1			
105203	Fam208b					14105	Srsf10			
208659	Fam20a					20382	Srsf2			

67017	Fam210b		20383	Srsf3				
235493	Fam214a		225027	Srsf7				
78323	Fam219b		76650	Srxn1				
67238	Fam220a		20823	Ssb				
69134	Fam25c		72475	Ssbp3				
215900	Fam26f		20442	St3gal1				
67922	Fam32a		20441	St3gal3				
75698	Fam35a		20443	St3gal4				
108958	Fam73b		20440	St6gal1				
213956	Fam83f		50935	St6galnac6				
105732	Fam83h		56792	Stap1				
69627	Fam89a		56018	Stard10				
68250	Fam96a		20846	Stat1				
72722	Fam98a		20847	Stat2				
68215	Fam98b		20850	Stat5a				
14102	Fas		20852	Stat6				
380601	Fastkd5		20867	Stip1				
246256	Fcgr4		20872	Stk16				
14132	Fcgrt		56274	Stk3				
218503	Fcho2		58231	Stk4				
14133	Fcna		13830	Stom				
218952	Fermt2		74152	Stra6l				
225020	Fez2		94186	Strn3				
14164	Fgf1		68292	Stt3b				
56636	Fgf21		20909	Stx4a				
75296	Fgfr1op		20024	Sub1				
234199	Fgl1		84112	Sucnr1				
231630	Ficd		192136	Sugct				
68680	Fitm1		76971	Sult2a8				
14228	Fkbp4		57429	Sult5a1				
54418	Fmn2		20610	Sumo3				
207854	Fmr1nb		211389	Suox				
214459	Fnbp1l		114741	Supt16				
319448	Fndc3a		96935	Susd4				
72007	Fndc3b		66962	Swsap1				
14276	Folr2		67043	Syap1				
15376	Foxa2		71846	Syce2				
15377	Foxa3		20962	Sycp3				
14236	Foxn2		56403	Syncrip				
14287	Fpgs		217030	Synrg				
75540	Fpgt		74126	Syvn1				
14289	Fpr2		213491	Szrd1				
56095	Ftsj3		231151	Tada2b				
51886	Fubp1		21339	Taf1a				
320267	Fubp3		21354	Tap1				
233908	Fus		230908	Tardbp				
14297	Fxn		110960	Tars				
17281	Fyco1		52440	Tax1bp1				
14369	Fzd7		56480	Tbk1				
14370	Fzd8		21372	Tbl1x				
14373	G0s2		21376	Tbrg1				
27041	G3bp1		21393	Tcap				

23881	G3bp2		21399	Tceal1	
57436	Gabarapl1		67923	Tceb1	
13197	Gadd45a		23965	Tenm3	
17873	Gadd45b		21754	Tesk1	
14635	Galk1		66654	Tex12	
14423	Galnt1		21763	Tex2	
108148	Galnt2		75623	Tex30	
14455	Gas5		21780	Tfam	
14456	Gas6		21786	Tff3	
234366	Gata2a		50765	Tfr2	
14469	Gbp2		22042	Tfrc	
14468	Gbp2b		21812	Tgfbr1	
55932	Gbp3		21817	Tgm2	
100702	Gbp6		74088	Them7	
229900	Gbp7		331401	Thoc2	
14527	Gcgr		21832	Thpo	
14629	Gclc		21838	Thy1	
14630	Gclm		21846	Tie1	
68133	Gcsh		68140	Tigd2	
14544	Gda		276891	Timd4	
1E+08	Gdap10		76916	Timmdc1	
23886	Gdf15		66131	Tipin	
66603	Gemin2		27375	Tjp3	
11692	Gfer		21885	Tle1	
70575	Gfod2		21886	Tle2	
223267	Ggact		114606	Tle6	
110175	Ggct		384059	Tlr12	
14600	Ghr		53791	Tlr5	
76425	Gid8		73166	Tm7sf2	
317758	Gimap9		68059	Tm9sf2	
215751	Ginm1		107358	Tm9sf3	
14618	Gjb1		69660	Tmbim1	
14619	Gjb2		330401	Tmcc1	
104174	Gldc		68944	Tmc01	
109801	Glo1		67511	Tmed9	
107146	Glyat		76547	Tmem101	
235582	Glyctk		215210	Tmem120a	
63985	Gmfb		71929	Tmem123	
331026	Gmppb		230678	Tmem125	
229363	Gmps		66990	Tmem134	
14673	Gna12		232086	Tmem150a	
14675	Gna14		66074	Tmem167	
14677	Gnai1		66058	Tmem176a	
30877	Gnl3		66343	Tmem177	
54342	Gnppnat1		75146	Tmem180	
54214	Golga4		231832	Tmem184a	
57437	Golga7		67226	Tmem19	
66629	Golph3		70612	Tmem230	
73274	Gpbp1		652925	Tmem243	
14733	Gpc1		69981	Tmem30a	
14735	Gpc4		238257	Tmem30b	
14555	Gpd1		67878	Tmem33	

14571	Gpd2					66079	Tmem42		
268566	Gphn		■	■	■	99887	Tmem56		
107173	Gpr137					100201	Tmem64		
80290	Gpr146					67893	Tmem86a		
76238	Grhpr					103743	Tmem98		
227753	Gsn		■	■	■	20776	Tmie		
14852	Gspt1		■	■	■	19241	Tmsb4x		
14782	Gsr					72736	Tmx1		
14854	Gss					21928	Tnfaip2		
14859	Gsta3					66443	Tnfaip8l1		
14865	Gstm4					21925	Tnncl2		
98053	Gtf2f1					21955	Tnnt1		
68705	Gtf2f2	■				21957	Tnnt3		
14884	Gtf2h1		■	■	■	67952	Tomm20		
23894	Gtf2h2		■	■	■	28185	Tomm70a		
70239	Gtf3c5					72960	Top1mt		
207704	Gtpbp10		■	■	■	240832	Tor1aip2		
69237	Gtpbp4		■	■	■	30933	Tor2a		
14923	Guk1		■	■	■	110012	Tpgs1		
14933	Gyk		■	■	■	22004	Tpm2		
14958	H1f0		■	■	■	22019	Tpp2		
14960	H2-Aa		■	■	■	67808	Tprgl		
14961	H2-Ab1		■	■	■	22031	Traf3		
15270	H2afx		■	■	■	66975	Trappc13		
14969	H2-Eb1		■	■	■	58866	Treh		
15042	H2-T24		■	■	■	224840	Treml4		
30963	Hacd1					319236	Trim12c		
70757	Hacd2					80890	Trim2		
56794	Hacl1		■	■	■	20821	Trim21		
97212	Hadha		■	■	■	20128	Trim30a		
231086	Hadhb					209387	Trim30d		
15130	Hbb-b2					69807	Trim32		
353502	Hcfc1r1					79263	Trim39		
433759	Hdac1					108943	Trmt10a		
232232	Hdac11					66926	Trmt6		
72748	Hdhd3	■	■	■	■	328162	Trmt61a		
207304	Hectd1		■	■	■	60599	Trp53inp1		
67138	Herc6					68728	Trp53inp2		
15205	Hes1					58800	Trpm7		
55927	Hes6					22368	Trpv2		
192231	Hexim1					227682	Trub2		
15234	Hgf					76927	Tsacc		
54426	Hgfac					21807	Tsc22d1		
68917	Hint2					232670	Tspan33		
15258	Hipk2		■	■	■	22110	Tspyl1		
15259	Hipk3		■	■	■	69499	Tsr2		
50708	Hist1h1c					56844	Tssc4		
319158	Hist1h4i					234875	Ttc13		
319190	Hist2h2be					75516	Ttc32		
110521	Hivep1					72747	Ttc39c		
217082	Hif					104718	Ttc7b		
15354	Hmgb3					75425	Tti1		

15360	Hmgcs2		50500	Ttpa	
21410	Hnf1b		117160	Ttyh2	
15378	Hnf4a		227613	Tubb4b	
68314	Hnf4aos		67510	Tvp23b	
229279	Hnrnpa3		65960	Twsg1	
15384	Hnrnpab		52700	Txndc17	
15381	Hnrnpc		105245	Txndc5	
11991	Hnrnpd		234723	Txnl4b	
98758	Hnrnpf		22177	Tyrobp	
56258	Hnrnph2		71767	Tysnd1	
15387	Hnrnpk		68736	Tyw5	
74326	Hnrnpr		22200	Uba3	
51810	Hnrnpu		66663	Uba5	
74318	Hopx		24108	Ubd	
15439	Hp		22209	Ube2a	
15446	Hpgd		56550	Ube2d2a	
15460	Hr		66105	Ube2d3	
15108	Hsd17b10		22194	Ube2e1	
56348	Hsd17b12		218793	Ube2e2	
243168	Hsd17b13		67921	Ube2f	
15486	Hsd17b2		67128	Ube2g1	
15488	Hsd17b4		22213	Ube2g2	
15493	Hsd3b2		56791	Ube2l6	
15494	Hsd3b3		109161	Ube2q2	
15496	Hsd3b5		70620	Ube2v2	
101502	Hsd3b7		22215	Ube3a	
72552	Hsdl1		27643	Ubl4	
72479	Hsdl2		69459	Ubl7	
15519	Hsp90aa1		79560	Ublcp1	
15516	Hsp90ab1		225896	Ubxn1	
22027	Hsp90b1		217379	Ubxn2a	
110920	Hspa13		67812	Ubxn4	
15525	Hspa4		22245	Uck1	
14828	Hspa5		22230	Ufd1l	
80888	Hspb8		67490	Ufl1	
15505	Hspf1		71773	Ugt2b1	
12282	Hyou1		100727	Ugt2b34	
216792	Iba57		243085	Ugt2b35	
15902	Id2		22238	Ugt2b5	
15903	Id3		223337	Ugt3a2	
15925	Ide		29869	Ulk2	
15936	Ier2		22248	Unc119	
26388	Ifi202b		19704	Upf1	
76933	Ifi27l2a		22276	Uros	
70110	Ifi35		56041	Uso1	
99899	Ifi44		230484	Usp1	
15953	Ifi47		59025	Usp14	
71586	Ifih1		24110	Usp18	
15957	Ifit1		170822	Usp33	
15958	Ifit2		22284	Usp9x	
15959	Ifit3		231915	Uspl1	
15980	Ifngr2		217109	Utp18	

55978	Ift20		216987	Utp6		
67286	Ift22		22330	Vcl		
16005	Igfals		269523	Vcp		
16009	Igfbp3		22361	Vnn1		
207683	Igsf11		26464	Vnn3		
16145	Igtp		52348	Vps37a		
60440	Ilgp1		330192	Vps37b		
56489	Ikbke		218035	Vps41		
16151	Ikbkg		21427	Vps72		
67143	Ikzf5		53612	Vti1b		
16164	Il13ra1		219189	Vwa8		
16195	Il6st		225131	Wac		
66541	Immp1l		94254	Wbscr16		
242291	Impad1		216156	Wdr18		
103844	Inca1		70465	Wdr77		
16325	Inhbc		230796	Wdtc1		
16326	Inhbe		192200	Wfdc12		
233752	Insc		67701	Wfdc2		
72999	Insig2		66107	Wfdc21		
70422	Ints2		232341	Wnk1		
230673	Ipo13		22413	Wnt2		
75751	Ipo4		59002	Wrap73		
320727	Ipo8		78903	Wrnip1		
74239	Iqce		60532	Wtap		
16179	Irak1		22433	Xbp1		
108960	Irak2		11798	Xiap		
27056	Irf5		321003	Xpnpep3		
54139	Irf6		73192	Xpot		
16391	Irf9		74335	Xrcc3		
15944	Irgm1		102448	Xylb		
54396	Irgm2		217119	Xylt2		
57444	Isg20		67008	Yae1d1		
26968	Islr		67057	Yaf2		
67441	Isoc2b		56449	Ybx3		
71927	Itfg1		67180	Yipf5		
106581	Itfg3		77929	Yipf6		
241226	Itga8		56418	Ykt6		
217837	Itpk1		27377	Yme1l1		
269424	Jade1		240255	Ythdc2		
67767	Jagn1		213541	Ythdf2		
16453	Jak3		229096	Ythdf3		
16069	Jchain		22627	Ywhae		
66124	Josd2		22628	Ywhag		
16476	Jun		58203	Zbp1		
16477	Junb		66391	Zbtb11os1		
16478	Jund		30928	Zbtb18		
235041	Kank2		56490	Zbtb20		
14534	Kat2a		81630	Zbtb22		
67773	Kat8		56805	Zbtb33		
72425	Katnbl1		226470	Zbtb41		
16529	Kcnk5		237256	Zc3h12d		
84036	Kcnn1		76014	Zc3h18		

233107	Kctd15			78781	Zc3hav1		
218214	Kdm1b			227449	Zcchc2		
70750	Kdsr			214290	Zcchc6		
50868	Keap1			66361	Zfand1		
64697	Keg1			100494	Zfand2a		
16573	Kif5b			11906	Zfhx3		
21847	Klf10			22640	Zfp1		
16601	Klf9			65020	Zfp110		
242721	Klhd7a			72465	Zfp131		
77113	Klh12			26465	Zfp146		
242521	Klh19			22661	Zfp148		
16646	Kpna1			26466	Zfp260		
16648	Kpna3			74120	Zfp263		
16649	Kpna4			30046	Zfp292		
16211	Kpnb1			218100	Zfp322a		
16691	Krt8			54367	Zfp326		
100087	Kti12			230761	Zfp362		
214669	L3mbtl2			12192	Zfp3611		
16763	Lad1			22701	Zfp41		
16768	Lag3			235682	Zfp445		
16774	Lama3			224691	Zfp472		
16780	Lamb3			22709	Zfp51		
16784	Lamp2			72672	Zfp518a		
68576	Lamtor5			240068	Zfp563		
217980	Larp4b			170938	Zfp617		
16796	Lasp1			22720	Zfp62		
16803	Lbp			72611	Zfp655		
16816	Lcat			319475	Zfp672		
212391	Lcor			1E+08	Zfp729b		
18826	Lcp1			233056	Zfp790		
16828	Ldha			240064	Zfp799		
52815	Ldhd			235047	Zfp809		
259301	Leap2			235956	Zfp825		
68192	Leprotl1			69504	Zfp932		
16854	Lgals3			71508	Zfp935		
19039	Lgals3bp			74149	Zfp946		
56048	Lgals8			22763	Zfr		
76429	Lhpp			218441	Zfyve16		
16880	Lifr			72739	Zkscan3		
319583	Lig4			22401	Zmat3		
108030	Lin7a			228880	Zmynd8		
22343	Lin7c			98999	Znfx1		
16889	Lipa			70103	Znhit1		
15450	Lipc			29805	Znhit2		
16891	Lipg			229937	Znhit6		
56722	Litaf			387524	Znrf2		
214895	Lman2l			99334	Zscan29		
232685	Lncpint			71971	Zswim1		
66887	Lonp2			69747	Zswim7		
16949	Loxl1			52696	Zwint		
226856	Lpgat1						

T: Pearson correlation values between CR and the plant bioactives (including synergistic analysis) for all pathways which include at least one CR gene with a fold change > 1.5

Pathway	set size (genes in bg)	CR genes	R	P	Q	RP	RQ	PQ	RPQ
4-hydroxyproline degradation I - MouseCyc	4(4)	1 (25.0%)	0,90	0,96	0,99	0,95	0,94	0,98	0,96
Acetylcholine Binding And Downstream Events - Reactome	14(14)	1 (7.1%)	0,55	0,87	0,53	0,82	0,67	0,87	0,85
Activated TLR4 signalling - Reactome	95(83)	2 (2.4%)	-0,39	-0,52	-0,57	-0,54	-0,52	-0,63	-0,60
Activation of Gene Expression by SREBP (SREBF) - Reactome	7(7)	1 (14.3%)	-0,30	-0,17	-0,74	-0,27	-0,62	-0,63	-0,53
Activation of Nicotinic Acetylcholine Receptors - Reactome	14(14)	1 (7.1%)	0,55	0,87	0,53	0,82	0,67	0,87	0,85
acyl-ACP thioesterase pathway - MouseCyc	1(1)	1 (100%)	-	-	-	-	-	-	-
Adaptive Immune System - Reactome	457 (399)	1 (0.3%)	0,02	0,14	0,14	0,08	0,06	0,18	0,11
Adipocytokine signaling pathway - Mus musculus (mouse) - KEGG	71(66)	2 (3.0%)	0,08	0,32	0,00	0,20	0,07	0,22	0,17
Adipogenesis - Wikopathways	133 (125)	3 (2.4%)	0,14	0,10	0,07	0,17	0,15	0,11	0,17
ADP signalling through P2Y purinoceptor 1 - Reactome	9(9)	1 (11.1%)	0,51	0,52	0,90	0,58	0,69	0,72	0,68
Advanced glycosylation endproduct receptor signaling - Reactome	13(12)	2 (16.7%)	-0,64	-0,78	-0,88	-0,79	-0,79	-0,85	-0,83
Alanine and aspartate metabolism - Wikopathways	12(12)	1 (8.3%)	0,49	0,60	0,29	0,78	0,46	0,66	0,73
Alanine, aspartate and glutamate metabolism - Mus musculus (mouse) - KEGG	35(29)	2 (6.9%)	0,26	0,46	0,37	0,43	0,31	0,53	0,45
Alcoholism - Mus musculus (mouse) - KEGG	202 (105)	1 (1.0%)	0,34	0,16	0,08	0,41	0,35	0,17	0,41
Alternative complement activation - Reactome	3(2)	1 (50.0%)	1,00	-1,00	-1,00	1,00	1,00	-1,00	1,00
Amine Oxidase reactions - Reactome	4(4)	1 (25.0%)	-1,00	0,70	0,34	0,19	-0,54	0,72	0,35
Amine-derived hormones - Reactome	14(14)	2 (14.3%)	0,33	0,02	-0,35	0,35	0,30	-0,09	0,32
Amino acid and oligopeptide SLC transporters - Reactome	50(46)	1 (2.2%)	0,38	0,28	0,16	0,51	0,38	0,37	0,50
Amino acid synthesis and interconversion (transamination) - Reactome	17(15)	2 (13.3%)	0,38	0,58	0,35	0,49	0,41	0,66	0,51
Amino acid transport across the plasma membrane - Reactome	32(30)	1 (3.3%)	0,46	0,35	0,03	0,60	0,40	0,35	0,54
Amoebiasis - Mus musculus (mouse) - KEGG	120 (108)	4 (3.7%)	0,16	0,35	0,19	0,33	0,20	0,38	0,35
Amphetamine addiction - Mus musculus (mouse) - KEGG	68(63)	1 (1.6%)	0,66	0,07	0,10	0,68	0,67	0,11	0,68
Androgen biosynthesis - Reactome	15(11)	2 (18.2%)	0,97	0,31	0,52	0,97	0,97	0,46	0,97
Antiviral mechanism by IFN-stimulated genes - Reactome	25(21)	1 (4.8%)	-0,04	0,33	-0,16	0,12	-0,08	0,20	0,08
Arachidonic acid metabolism - Mus musculus (mouse) - KEGG	98(78)	4 (5.1%)	0,69	0,17	-0,17	0,67	0,63	0,06	0,61
Arachidonic acid metabolism - Reactome	76(57)	2 (3.5%)	0,61	0,30	-0,19	0,63	0,55	0,14	0,57
Arginine and proline metabolism - Mus musculus (mouse) - KEGG	57(48)	1 (2.1%)	0,04	0,40	0,20	0,18	0,08	0,41	0,21
Ascorbate and aldarate metabolism - Mus musculus (mouse) - KEGG	27(14)	2 (14.3%)	0,75	0,44	0,09	0,80	0,69	0,34	0,71
asparagine biosynthesis I - MouseCyc	2(2)	1 (50.0%)	-1,00	1,00	1,00	1,00	-1,00	1,00	1,00
Bile acid and bile salt metabolism - Reactome	36(36)	4 (11.1%)	0,18	0,22	-0,18	0,23	0,11	0,08	0,17
bile acid biosynthesis, neutral pathway - MouseCyc	6(6)	1 (16.7%)	-0,35	-0,28	-0,82	-0,34	-0,49	-0,61	-0,45
Bile secretion - Mus musculus (mouse) - KEGG	72(68)	2 (2.9%)	0,25	0,22	0,03	0,31	0,23	0,17	0,28
Binding and Uptake of Ligands by Scavenger Receptors - Reactome	39(31)	2 (6.5%)	-0,46	-0,69	-0,65	-0,66	-0,61	-0,72	-0,69

Biogenic Amine Synthesis - Wikipathways	15(14)	1 (7.1%)	0,90	-0,09	-0,10	0,91	0,87	-0,11	0,86
Biological oxidations - Reactome	168 (128)	14 (10.9%)	0,54	0,21	-0,01	0,54	0,51	0,15	0,51
biosynthesis of estrogens - MouseCyc	12(11)	2 (18.2%)	0,35	0,80	0,50	0,53	0,47	0,80	0,61
biosynthesis of serotonin and melatonin - MouseCyc	5(4)	1 (25.0%)	1,00	-0,38	-0,38	0,98	1,00	-0,39	0,97
Biosynthesis of unsaturated fatty acids - Mus musculus (mouse) - KEGG	25(24)	1 (4.2%)	0,37	0,53	-0,41	0,49	0,18	0,15	0,37
Biotin transport and metabolism - Reactome	10(10)	2 (20.0%)	0,04	0,49	-0,60	0,18	-0,02	0,35	0,12
biotin-carboxyl carrier protein assembly - MouseCyc	3(3)	2 (66.7%)	-1,00	0,76	-0,99	-0,94	-1,00	0,46	-0,96
bupropion degradation - MouseCyc	38(29)	3 (10.3%)	0,81	-0,01	-0,24	0,75	0,78	-0,15	0,74
Calcium Regulation in the Cardiac Cell - Wikipathways	150 (135)	2 (1.5%)	0,21	-0,18	-0,23	0,09	0,10	-0,22	0,00
Calcium signalling pathway - Mus musculus (mouse) - KEGG	183 (165)	1 (0.6%)	0,06	0,20	0,19	0,14	0,10	0,26	0,17
catecholamine biosynthesis - MouseCyc	6(6)	2 (33.3%)	0,56	0,53	-0,81	0,78	0,52	0,41	0,75
Catecholamine biosynthesis - Reactome	4(4)	1 (25.0%)	0,99	-0,50	-0,69	0,99	0,98	-0,76	0,99
Cell cycle - Wikipathways	88(79)	1 (1.3%)	0,17	0,23	0,15	0,24	0,23	0,26	0,28
Chagas disease (American trypanosomiasis) - Mus musculus (mouse) - KEGG	104 (100)	1 (1.0%)	-0,02	0,28	0,15	0,12	0,04	0,30	0,15
Chemical carcinogenesis - Mus musculus (mouse)	95(70)	8 (11.4%)	0,36	0,09	-0,22	0,33	0,27	-0,04	0,26
Cholesterol biosynthesis - Reactome	22(22)	2 (9.1%)	0,40	0,44	0,00	0,49	0,38	0,33	0,46
Cholesterol Biosynthesis - Wikipathways	15(15)	2 (13.3%)	0,24	0,53	0,14	0,40	0,26	0,43	0,39
cholesterol biosynthesis I - MouseCyc	12(12)	1 (8.3%)	0,39	0,46	0,22	0,47	0,41	0,40	0,46
cholesterol biosynthesis III (via desmosterol) - MouseCyc	11(11)	1 (9.1%)	0,38	0,49	0,24	0,47	0,40	0,43	0,47
Class A/1 (Rhodopsin-like receptors) - Reactome	290 (234)	2 (0.9%)	-0,34	-0,25	-0,48	-0,39	-0,45	-0,42	-0,47
Clathrin derived vesicle budding - Reactome	61(52)	1 (1.9%)	0,62	0,15	-0,43	0,69	0,50	-0,09	0,56
Cocaine addiction - Mus musculus (mouse) - KEGG	50(46)	1 (2.2%)	0,68	0,14	0,04	0,73	0,67	0,12	0,72
Complement Activation, Classical Pathway - Wikipathway	17(13)	3 (23.1%)	0,54	0,34	-0,26	0,57	0,48	0,22	0,52
Complement and coagulation cascades - Mus musculus (mouse) - KEGG	77(65)	5 (7.7%)	0,27	0,25	0,02	0,38	0,26	0,22	0,36
Complement and Coagulation Cascades - Wikipathways	61(57)	3 (5.3%)	0,27	0,21	0,11	0,37	0,28	0,22	0,37
Complement cascade - Reactome	40(28)	4 (14.3%)	0,41	0,29	-0,13	0,52	0,38	0,16	0,49
Constitutive PI3K/AKT Signaling in Cancer - Reactome	81(77)	1 (1.3%)	0,23	0,28	0,31	0,32	0,33	0,37	0,40
Cysteine and methionine metabolism - Mus musculus (mouse) - KEGG	38(33)	2 (6.1%)	0,54	0,14	-0,01	0,58	0,48	0,12	0,53
Cytochrome P450 - arranged by substrate type - Reactome	84(55)	7 (12.7%)	0,64	0,07	-0,30	0,57	0,55	-0,10	0,49
Cytochrome P450 - Wikipathways	41(35)	3 (8.6%)	0,29	0,08	-0,25	0,25	0,21	-0,05	0,19
Cytokine Signaling in Immune system - Reactome	193 (168)	2 (1.2%)	0,00	0,27	0,01	0,13	0,00	0,19	0,11
Cytokine-cytokine receptor interaction - Mus musculus (mouse) - KEGG	266 (216)	2 (0.9%)	0,30	0,31	0,24	0,40	0,33	0,35	0,41
Cytosolic sensors of pathogen-associated DNA - Reactome	58(45)	2 (4.4%)	-0,64	-0,68	-0,77	-0,74	-0,76	-0,77	-0,78
Cytosolic sulfonation of small molecules - Reactome	18(14)	2 (14.3%)	0,34	0,41	0,44	0,61	0,49	0,54	0,73
DAP12 interactions - Reactome	158 (144)	1 (0.7%)	0,16	0,25	0,22	0,24	0,21	0,30	0,28
DAP12 signaling - Reactome	147 (138)	1 (0.7%)	0,16	0,25	0,23	0,24	0,22	0,30	0,28

Defective AMN causes hereditary megaloblastic anemia 1 - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective BTD causes biotidinase deficiency - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective CD320 causes methylmalonic aciduria - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective CUBN causes hereditary megaloblastic anemia 1 - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective GIF causes intrinsic factor deficiency - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective HLCS causes multiple carboxylase deficiency - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective LMBRD1 causes methylmalonic aciduria and homocystinuria type cbIF - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective MMAA causes methylmalonic aciduria type cbIA - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective MMAB causes methylmalonic aciduria type cbIB - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective MMACHC causes methylmalonic aciduria and homocystinuria type cbIC - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective MMADHC causes methylmalonic aciduria and homocystinuria type cbID - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective MTR causes methylmalonic aciduria and homocystinuria type cbIG - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective MTRR causes methylmalonic aciduria and homocystinuria type cbIE - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective MUT causes methylmalonic aciduria mut type - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defective TCN2 causes hereditary megaloblastic anemia - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defects in biotin (Btn) metabolism - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defects in cobalamin (B12) metabolism - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Defects in vitamin and cofactor metabolism - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Degradation of cysteine and homocysteine - Reactome	8(8)	1 (12.5%)	0,66	0,81	-0,47	0,76	0,58	0,89	0,74
DEx/H-box helicases activate type I IFN and inflammatory cytokines production - Reactome	13(12)	2 (16.7%)	-0,82	-0,81	-0,87	-0,86	-0,88	-0,84	-0,87
Disease - Reactome	608 (554)	4 (0.7%)	0,14	0,23	0,03	0,23	0,14	0,19	0,21
Diurnally Regulated Genes with Circadian Orthologs - Wikipathways	48(42)	1 (2.4%)	0,39	0,33	0,29	0,44	0,40	0,42	0,45
Dopaminergic synapse - Mus musculus (mouse) - KEGG	135 (120)	1 (0.8%)	0,27	0,18	0,15	0,32	0,28	0,21	0,32
Dopaminergic Neurogenesis - Wikipathways	30(30)	1 (3.3%)	0,90	0,01	-0,15	0,91	0,84	-0,11	0,86
Downstream signal transduction - Reactome	142 (134)	1 (0.7%)	0,17	0,26	0,24	0,26	0,23	0,31	0,30
Downstream Signaling Events Of B Cell Receptor (BCR) - Reactome	108 (104)	1 (1.0%)	0,21	0,24	0,28	0,29	0,30	0,32	0,35
Downstream signaling of activated FGFR - Reactome	130 (122)	1 (0.8%)	0,23	0,26	0,28	0,31	0,29	0,33	0,35
Drug metabolism - cytochrome P450 - Mus musculus (mouse) - KEGG	97(70)	10 (14.3%)	0,46	0,25	-0,17	0,48	0,40	0,12	0,43
Drug metabolism - other enzymes - Mus musculus (mouse) - KEGG	61(39)	3 (7.7%)	0,45	0,33	-0,17	0,53	0,35	0,15	0,42
EGFR1 Signaling Pathway - Wikipathways	176 (169)	1 (0.6%)	0,04	-0,17	-0,22	-0,07	-0,06	-0,23	-0,13
Endogenous sterols - Reactome	14(13)	3 (23.1%)	0,14	-0,14	-0,60	0,08	0,01	-0,41	-0,02
ErbB signaling pathway - Mus musculus (mouse) - KEGG	87(82)	1 (1.2%)	0,31	0,25	0,32	0,38	0,42	0,38	0,47
ErbB signaling pathway - Wikipathways	46(44)	1 (2.3%)	0,47	0,32	0,44	0,54	0,64	0,50	0,67

Estrogen metabolism - Wikipathways	14(8)	1 (12.5%)	0,42	0,71	0,84	0,60	0,64	0,89	0,74
Estrogen signaling pathway - <i>Mus musculus</i> (mouse) - KEGG	99(86)	1 (1.2%)	0,35	0,17	0,23	0,38	0,37	0,25	0,39
ethanol degradation II (cytosol) - MouseCyc	11(10)	1 (10.0%)	0,66	0,68	0,69	0,72	0,75	0,84	0,78
Ethanol oxidation - Reactome	10(9)	1 (11.1%)	0,81	0,76	0,37	0,83	0,87	0,78	0,88
Exercise-induced Circadian Regulation - Wikipathways	49(42)	2 (4.8%)	0,31	0,32	0,27	0,37	0,33	0,39	0,39
Facilitative Na ⁺ -independent glucose transporters - Reactome	11(10)	1 (10.0%)	-0,27	0,68	0,58	0,05	-0,15	0,77	0,21
Fatty acid biosynthesis - <i>Mus musculus</i> (mouse) - KEGG	6(6)	3 (50.0%)	-0,46	0,54	-0,76	-0,13	-0,62	-0,37	-0,43
Fatty Acid Biosynthesis - Wikipathways	22(21)	3 (14.3%)	-0,39	0,53	-0,37	0,03	-0,45	0,19	-0,17
fatty acid biosynthesis initiation II - MouseCyc	2(2)	1 (50.0%)	-1,00	1,00	-1,00	-1,00	-1,00	-1,00	-1,00
fatty acid biosynthesis initiation III - MouseCyc	2(2)	1 (50.0%)	-1,00	1,00	-1,00	-1,00	-1,00	-1,00	-1,00
Fatty acid degradation - <i>Mus musculus</i> (mouse) - KEGG	48(37)	2 (5.4%)	0,81	0,11	0,06	0,75	0,79	0,12	0,73
Fatty acid elongation - <i>Mus musculus</i> (mouse) - KEGG	24(24)	2 (8.3%)	0,78	0,38	-0,58	0,80	0,77	-0,06	0,78
Fatty Acid Omega Oxidation - Wikipathways	7(7)	1 (14.3%)	0,55	0,24	0,71	0,50	0,67	0,58	0,61
Fatty acid, triacylglycerol, and ketone body metabolism - Reactome	85(77)	6 (7.8%)	0,53	0,41	-0,10	0,57	0,52	0,21	0,56
Fatty acids - Reactome	18(9)	1 (11.1%)	0,90	0,15	-0,17	0,84	0,85	0,02	0,79
Fatty Acyl-CoA Biosynthesis - Reactome	17(16)	4 (25.0%)	0,77	0,38	-0,71	0,78	0,74	-0,36	0,75
Fc epsilon receptor (FCER1) signaling - Reactome	155(141)	1 (0.7%)	0,22	0,26	0,31	0,31	0,32	0,36	0,37
FMO oxidizes nucleophiles - Reactome	3(3)	1 (33.3%)	0,93	0,79	0,91	0,99	0,98	0,85	1,00
formaldehyde oxidation I - MouseCyc	5(5)	1 (20.0%)	0,23	-0,51	-0,53	-0,03	-0,01	-0,88	-0,25
Formyl peptide receptors bind formyl peptides and many other ligands - Reactome	13(11)	2 (18.2%)	-0,58	-0,77	-0,84	-0,80	-0,78	-0,81	-0,83
G alpha (i) signalling events - Reactome	222(156)	2 (1.3%)	-0,37	-0,36	-0,54	-0,46	-0,49	-0,50	-0,53
G alpha (q) signalling events - Reactome	167(141)	3 (2.1%)	-0,33	-0,37	-0,48	-0,44	-0,44	-0,47	-0,49
G Protein Signaling Pathways - Wikipathways	91(84)	1 (1.2%)	0,17	0,24	0,46	0,27	0,30	0,40	0,37
GAB1 signalosome - Reactome	90(86)	1 (1.2%)	0,24	0,30	0,34	0,34	0,34	0,39	0,41
Gap junction assembly - Reactome	3(3)	1 (33.3%)	0,00	-0,87	0,86	-0,25	0,13	-0,54	-0,13
Gap junction trafficking - Reactome	10(10)	1 (10.0%)	-0,06	-0,23	0,08	-0,16	-0,05	-0,20	-0,14
Gap junction trafficking and regulation - Reactome	12(12)	1 (8.3%)	-0,04	-0,18	0,21	-0,12	0,00	-0,09	-0,08
Gastrin-CREB signalling pathway via PKC and MAPK - Reactome	188(162)	3 (1.9%)	-0,29	-0,34	-0,46	-0,40	-0,40	-0,45	-0,45
Glucocorticoid & Mineralcorticoid Metabolism - Wikipathways	13(11)	1 (9.1%)	1,00	0,34	0,53	1,00	1,00	0,48	1,00
Glucocorticoid biosynthesis - Reactome	13(8)	1 (12.5%)	1,00	0,30	0,49	1,00	1,00	0,44	1,00
Gluconeogenesis - Reactome	35(30)	1 (3.3%)	0,19	0,42	0,12	0,33	0,20	0,37	0,33
Glucose metabolism - Reactome	69(60)	1 (1.7%)	0,26	0,48	0,08	0,42	0,26	0,41	0,39
Glucose transport - Reactome	40(37)	1 (2.7%)	0,24	0,40	0,18	0,31	0,26	0,37	0,32
Glucuronidation - Wikipathways	16(10)	1 (10.0%)	0,90	0,15	0,06	0,89	0,90	0,14	0,86
Glutathione and one carbon metabolism - Wikipathways	31(28)	2 (7.1%)	0,51	0,37	0,01	0,58	0,50	0,30	0,56
Glutathione conjugation - Reactome	26(25)	1 (4.0%)	0,37	0,11	0,13	0,35	0,37	0,14	0,35

Glutathione metabolism - Mus musculus (mouse) - KEGG	55(48)	4 (8.3%)	0,19	0,05	-,09	0,18	0,16	0,00	0,15
Glutathione metabolism - Wikipathways	19(17)	1 (5.9%)	0,59	0,38	0,50	0,60	0,61	0,53	0,62
glutathione-mediated detoxification - MouseCyc	27(22)	2 (9.1%)	-0,11	-0,43	-0,29	-0,22	-0,15	-0,44	-0,25
Glycerolipid metabolism - Mus musculus (mouse) - KEGG	56(48)	1 (2.1%)	0,18	0,31	0,43	0,28	0,31	0,50	0,39
Glycerophospholipid biosynthesis - Reactome	86(79)	1 (1.3%)	0,15	0,29	0,40	0,21	0,23	0,46	0,28
Glycerophospholipid metabolism - Mus musculus (mouse) - KEGG	91(81)	1 (1.2%)	0,12	0,25	0,31	0,18	0,18	0,38	0,23
glycine betaine degradation - MouseCyc	8(8)	1 (12.5%)	0,79	-0,29	0,16	0,75	0,81	-0,06	0,75
Glycine, serine and threonine metabolism - Mus musculus (mouse) - KEGG	39(33)	2 (6.1%)	0,53	0,27	0,18	0,60	0,55	0,32	0,61
Glycolysis / Gluconeogenesis - Mus musculus (mouse) - KEGG	66(60)	1 (1.7%)	0,21	0,33	0,14	0,29	0,22	0,31	0,29
Glycolysis and Gluconeogenesis - Wikipathways	50(47)	2 (4.3%)	0,24	0,30	0,05	0,29	0,22	0,24	0,27
Glycoprotein hormones - Reactome	9(8)	2 (25.0%)	0,86	0,89	0,91	0,89	0,89	0,94	0,91
GPCR downstream signaling - Reactome	1065 (420)	3 (0.7%)	-0,22	-0,13	-0,31	-0,24	-0,30	-0,25	-0,30
GPCR ligand binding - Reactome	403 (306)	2 (0.7%)	-0,27	-0,21	-0,42	-0,33	-0,38	-0,36	-0,40
GRB2 events in ERBB2 signaling - Reactome	21(21)	1 (4.8%)	0,68	0,43	0,75	0,76	0,74	0,77	0,80
Heme biosynthesis - Reactome	11(11)	1 (9.1%)	0,79	0,42	0,07	0,81	0,75	0,35	0,77
Heme Biosynthesis - Wikipathways	9(9)	1 (11.1%)	0,88	0,47	0,09	0,87	0,84	0,36	0,83
heme biosynthesis II - MouseCyc	9(9)	1 (11.1%)	0,88	0,47	0,09	0,87	0,84	0,36	0,83
Hemostasis - Reactome	412 (364)	2 (0.5%)	0,08	0,21	0,08	0,18	0,10	0,21	0,19
Hepatitis C - Mus musculus (mouse) - KEGG	136 (120)	1 (0.8%)	-0,07	0,32	0,00	0,08	-0,07	0,22	0,07
Herpes simplex infection - Mus musculus (mouse) - KEGG	212 (164)	1 (0.6%)	-0,10	0,28	-0,07	0,01	-0,10	0,19	0,00
Hexose transport - Reactome	42(39)	1 (2.6%)	0,25	0,41	0,20	0,32	0,27	0,39	0,33
Highly calcium permeable nicotinic acetylcholine receptors - Reactome	9(9)	1 (11.1%)	0,70	0,94	0,80	0,90	0,77	0,96	0,91
Highly calcium permeable postsynaptic nicotinic acetylcholine receptors - Reactome	11(11)	1 (9.1%)	0,64	0,93	0,74	0,89	0,74	0,93	0,90
Histidine metabolism - Mus musculus (mouse) - KEGG	29(27)	1 (3.7%)	0,75	0,02	-0,01	0,76	0,74	0,01	0,73
Immune System - Reactome	894 (750)	9 (1.2%)	-0,02	0,06	-0,14	0,01	-0,06	-0,03	-0,03
Import of palmitoyl-CoA into the mitochondrial matrix - Reactome	9(7)	2 (28.6%)	0,21	0,87	-0,69	0,60	-0,13	0,46	0,30
Influenza A - Mus musculus (mouse) - KEGG	172 (146)	1 (0.7%)	-0,14	0,31	-0,05	0,01	-0,13	0,20	0,00
Initial triggering of complement - Reactome	21(12)	1 (8.3%)	0,60	0,00	-0,32	0,65	0,54	-0,23	0,61
Innate Immune System - Reactome	481 (399)	7 (1.8%)	-0,03	-0,05	-0,26	-0,05	-0,12	-0,17	-0,12
Insulin Signaling - Wikipathways	157 (150)	1 (0.7%)	0,24	0,20	-0,04	0,30	0,19	0,12	0,24
Insulin signaling pathway - Mus musculus (mouse) - KEGG	143 (125)	4 (3.2%)	0,19	0,36	-0,24	0,27	0,10	0,15	0,20
Integration of energy metabolism - Reactome	90(79)	1 (1.3%)	0,07	0,22	0,22	0,16	0,12	0,26	0,19
Interconversion of polyamines - Reactome	3(3)	1 (33.3%)	-0,83	0,72	0,45	-0,85	-0,83	0,76	-0,84
Interferon alpha/beta signaling - Reactome	28(20)	1 (5.0%)	-0,48	0,36	-0,18	-0,21	-0,41	0,14	-0,21
Interferon Signaling - Reactome	61(49)	2 (4.1%)	-0,10	0,30	-0,14	0,04	-0,13	0,13	0,00
ISG15 antiviral mechanism - Reactome	25(21)	1 (4.8%)	-0,04	0,33	-0,16	0,12	-0,08	0,20	0,08

L-cysteine degradation I - MouseCyc	3(3)	1 (33.3%)	1,00	0,95	0,51	1,00	0,98	1,00	1,00
L-cysteine degradation VI - MouseCyc	2(2)	1 (50.0%)	1,00	1,00	-1,00	1,00	1,00	1,00	1,00
Leptin and adiponectin - Wikipathways	10(10)	1 (10.0%)	0,17	0,50	-0,21	0,34	0,05	0,20	0,19
Lysosome - Mus musculus (mouse) - KEGG	124 (117)	2 (1.7%)	0,35	0,12	-0,25	0,40	0,29	-0,02	0,34
Lysosome Vesicle Biogenesis - Reactome	23(21)	1 (4.8%)	0,83	0,19	-0,75	0,83	0,75	-0,28	0,77
Maturity onset diabetes of the young - Mus musculus (mouse) - KEGG	26(24)	1 (4.2%)	-0,12	0,29	-0,41	0,03	-0,26	-0,08	-0,11
Membrane Trafficking - Reactome	148(12 9)	2 (1.6%)	0,39	0,08	-0,29	0,41	0,28	-0,09	0,30
Metabolism - Reactome	1386 (1211)	38 (3.1%)	0,43	0,21	0,03	0,46	0,42	0,18	0,45
Metabolism of amino acids and derivatives - Reactome	143 (130)	7 (5.4%)	0,29	0,18	-0,03	0,34	0,27	0,15	0,32
Metabolism of carbohydrates - Reactome	242 (214)	4 (1.9%)	0,10	0,28	-0,07	0,21	0,06	0,18	0,16
Metabolism of lipids and lipoproteins - Reactome	491 (437)	16 (3.7%)	0,51	0,26	-0,03	0,53	0,49	0,17	0,52
Metabolism of polyamines - Reactome	16(12)	2 (16.7%)	0,33	0,66	0,34	0,58	0,38	0,75	0,61
Metabolism of porphyrins - Reactome	16(14)	1 (7.1%)	0,81	0,42	0,08	0,83	0,77	0,36	0,79
Metabolism of proteins - Reactome	521 (380)	3 (0.8%)	0,17	0,12	0,23	0,20	0,22	0,20	0,24
Metabolism of steroid hormones and vitamin D - Reactome	30(25)	2 (8.0%)	0,97	0,35	0,37	0,97	0,97	0,42	0,96
Metabolism of vitamins and cofactors - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Metabolism of water-soluble vitamins and cofactors - Reactome	73(68)	3 (4.4%)	0,06	0,32	-0,24	0,19	-0,03	0,13	0,11
Metabolism of xenobiotics by cytochrome P450 - Mus musculus (mouse)	96(71)	8 (11.3%)	0,40	0,09	-0,24	0,37	0,32	-0,05	0,30
Metapathway biotransformation - Wikipathways	142 (116)	8 (6.9%)	0,29	0,29	-0,05	0,33	0,26	0,20	0,31
methionine salvage II (mammalia) - MouseCyc	3(2)	1 (50.0%)	1,00	1,00	-1,00	1,00	1,00	1,00	1,00
Methionine salvage pathway - Reactome	6(6)	1 (16.7%)	0,98	0,78	0,78	0,93	0,97	0,94	0,97
Methylation - Reactome	10(9)	1 (11.1%)	0,59	0,71	0,81	0,73	0,75	0,78	0,78
Methylation - Wikipathways	9(9)	1 (11.1%)	0,75	0,70	0,67	0,77	0,76	0,72	0,77
mevalonate pathway I - MouseCyc	11(10)	1 (10.0%)	0,72	0,74	-0,45	0,86	0,64	0,54	0,80
Mineralocorticoid biosynthesis - Reactome	12(7)	1 (14.3%)	1,00	0,32	0,48	1,00	1,00	0,44	1,00
mRNA processing - Wikipathways	465 (395)	1 (0.3%)	0,13	0,21	-0,07	0,22	0,09	0,14	0,18
MyD88 cascade initiated on plasma membrane - Reactome	69(61)	2 (3.3%)	-0,48	-0,61	-0,63	-0,62	-0,60	-0,69	-0,67
MyD88 dependent cascade initiated on endosome - Reactome	75(67)	2 (3.0%)	-0,47	-0,59	-0,62	-0,61	-0,60	-0,68	-0,66
MyD88:Mal cascade initiated on plasma membrane - Reactome	69(61)	2 (3.3%)	-0,48	-0,61	-0,63	-0,62	-0,60	-0,69	-0,67
MyD88-independent cascade - Reactome	89(79)	2 (2.5%)	-0,40	-0,52	-0,58	-0,55	-0,53	-0,63	-0,60
Myometrial Relaxation and Contraction Pathways - Wikipathways	157 (147)	2 (1.4%)	0,10	-0,38	-0,05	-0,12	0,07	-0,33	-0,12
Neuroactive ligand-receptor interaction - Mus musculus (mouse) - KEGG	282 (239)	1 (0.4%)	0,06	0,41	0,10	0,27	0,09	0,36	0,27
Neuronal System - Reactome	254 (220)	1 (0.5%)	0,12	0,25	0,18	0,23	0,17	0,28	0,26
Neurotransmitter Receptor Binding And Downstream Transmission In The Postsynaptic Cell - Reactome	133 (116)	1 (0.9%)	0,12	0,22	0,20	0,21	0,17	0,26	0,25
NGF signalling via TRKA from the plasma membrane - Reactome	184 (171)	1 (0.6%)	0,20	0,26	0,30	0,29	0,28	0,35	0,35
Nicotinate and nicotinamide metabolism -	32(29)	2	0,56	0,74	0,42	0,74	0,61	0,74	0,75

Mus musculus (mouse) - KEGG		(6.9%)									
Nicotine Activity on Dopaminergic Neurons - Wikipathways	10(10)	1 (10.0%)	0,93	-0,22	0,18	0,92	0,95	-0,15	0,94		
nicotine degradation II - MouseCyc	56(40)	7 (17.5%)	0,79	0,13	-0,22	0,76	0,77	0,00	0,75		
nicotine degradation III - MouseCyc	48(34)	6 (17.6%)	0,79	0,00	-0,25	0,74	0,76	-0,14	0,72		
Nitrogen metabolism - Mus musculus (mouse) - KEGG	17(15)	1 (6.7%)	0,57	0,15	0,82	0,58	0,68	0,68	0,68		
Nuclear receptors in lipid metabolism and toxicity - Wikipathways	30(28)	2 (7.1%)	-0,03	-0,20	0,04	-0,10	-0,02	-0,16	-0,09		
Nuclear signaling by ERBB4 - Reactome	55(46)	1 (2.2%)	0,67	0,21	0,66	0,61	0,73	0,52	0,68		
Oligomerization of connexins into connexons - Reactome	3(3)	1 (33.3%)	0,00	-0,87	0,86	-0,25	0,13	-0,54	-0,13		
One Carbon Metabolism - Wikipathways	29(26)	1 (3.8%)	0,42	0,15	-0,13	0,46	0,39	0,04	0,43		
One carbon metabolism and related pathways - Wikipathways	49(46)	2 (4.3%)	0,36	0,52	-0,18	0,45	0,31	0,35	0,40		
Organic anion transport - Reactome	4(4)	1 (25.0%)	0,95	0,97	0,77	0,99	0,96	0,96	0,99		
Organic cation/anion/zwitterion transport - Reactome	12(12)	1 (8.3%)	0,79	0,38	0,61	0,84	0,82	0,54	0,85		
Ovarian steroidogenesis - Mus musculus (mouse) - KEGG	58(50)	2 (4.0%)	0,96	0,21	0,40	0,96	0,96	0,33	0,96		
palmitate biosynthesis I (animals) - MouseCyc	1(1)	1 (100.0 %)	-	-	-	-	-	-	-		
Pantothenate and CoA biosynthesis - Mus musculus (mouse) - KEGG	16(16)	1 (6.2%)	0,15	-0,05	0,20	0,10	0,24	0,06	0,18		
Pentose and glucuronate interconversions - Mus musculus (mouse) - KEGG	34(20)	2 (10.0%)	0,79	0,27	-0,12	0,77	0,74	0,11	0,68		
pentose phosphate pathway - MouseCyc	11(10)	2 (20.0%)	0,08	-0,12	-0,65	0,02	-0,24	-0,62	-0,27		
Pentose phosphate pathway - Mus musculus (mouse) - KEGG	30(28)	2 (7.1%)	0,02	0,25	-0,36	0,15	-0,11	-0,01	0,01		
Pentose Phosphate Pathway - Wikipathways	7(6)	2 (33.3%)	0,66	-0,03	-0,18	0,57	0,58	-0,13	0,52		
Pentose phosphate pathway (hexose monophosphate shunt) - Reactome	8(7)	2 (28.6%)	0,45	-0,21	-0,44	0,29	0,28	-0,42	0,16		
pentose phosphate pathway (oxidative branch) - MouseCyc	5(5)	2 (40.0%)	0,46	-0,30	-0,62	0,61	0,02	-0,92	-0,18		
Peptide hormone biosynthesis - Reactome	11(10)	2 (20.0%)	0,87	0,84	0,87	0,89	0,90	0,90	0,91		
Peptide hormone metabolism - Reactome	36(32)	2 (6.2%)	0,64	0,47	0,61	0,71	0,69	0,57	0,73		
Peptide ligand-binding receptors - Reactome	188 (145)	2 (1.4%)	-0,44	-0,37	-0,62	-0,52	-0,57	-0,55	-0,60		
Peroxisome - Mus musculus (mouse) - KEGG	81(76)	2 (2.6%)	0,36	0,27	-0,26	0,44	0,30	0,08	0,39		
Phagosome - Mus musculus (mouse) - KEGG	180 (137)	1 (0.7%)	-0,11	0,20	0,03	-0,02	-0,10	0,19	-0,01		
Phase 1 - Functionalization of compounds	102(72)	10 (13.9%)	0,63	0,17	-0,21	0,58	0,55	0,02	0,52		
Phase II conjugation - Reactome	67(57)	4 (7.0%)	0,36	0,34	0,40	0,45	0,45	0,44	0,50		
phenylalanine degradation III - MouseCyc	6(5)	1 (20.0%)	0,90	-0,26	0,84	0,65	0,93	0,26	0,75		
Phenylalanine metabolism - Mus musculus (mouse) - KEGG	24(19)	2 (10.5%)	0,73	0,01	0,28	0,75	0,76	0,13	0,76		
Phenylalanine, tyrosine and tryptophan biosynthesis - Mus musculus (mouse) - KEGG	9(7)	1 (14.3%)	0,85	-0,14	0,41	0,42	0,79	-0,01	0,44		
Phospholipid metabolism - Reactome	132 (120)	1 (0.8%)	0,12	0,25	0,30	0,19	0,18	0,35	0,24		
PI-3K cascade - Reactome	86(82)	1 (1.2%)	0,24	0,30	0,33	0,34	0,34	0,39	0,41		
PI3K events in ERBB2 signaling - Reactome	86(82)	1 (1.2%)	0,24	0,30	0,33	0,34	0,34	0,39	0,41		
PI3K events in ERBB4 signaling - Reactome	86(82)	1 (1.2%)	0,24	0,30	0,33	0,34	0,34	0,39	0,41		

PI3K/AKT activation - Reactome	89(85)	1 (1.2%)	0,24	0,30	0,33	0,34	0,34	0,39	0,41
PI3K/AKT Signaling in Cancer - Reactome	86(82)	1 (1.2%)	0,24	0,30	0,33	0,34	0,34	0,39	0,41
PIP3 activates AKT signaling - Reactome	86(82)	1 (1.2%)	0,24	0,30	0,33	0,34	0,34	0,39	0,41
Platelet activation, signaling and aggregation - Reactome	190 (166)	2 (1.2%)	0,18	0,14	0,13	0,24	0,21	0,17	0,26
Platelet degranulation - Reactome	86(70)	1 (1.4%)	0,28	0,09	-0,08	0,32	0,26	0,04	0,29
PodNet- protein-protein interactions in the podocyte - Wikipathways	315 (295)	1 (0.3%)	0,18	0,13	0,12	0,22	0,20	0,16	0,23
Polyamines are oxidized to amines, aldehydes and H2O2 by PAOs - Reactome	2(2)	1 (50.0%)	-1,00	1,00	1,00	-1,00	-1,00	1,00	-1,00
Porphyrin and chlorophyll metabolism - Mus musculus (mouse) - KEGG	43(27)	3 (11.1%)	0,75	0,27	-0,08	0,77	0,69	0,15	0,69
Postsynaptic nicotinic acetylcholine receptors - Reactome	14(14)	1 (7.1%)	0,55	0,87	0,53	0,82	0,67	0,87	0,85
PPAR signaling pathway - Mus musculus (mouse) - KEGG	81(67)	3 (4.5%)	0,43	0,15	-0,15	0,44	0,37	0,04	0,39
PPAR signaling pathway - Wikipathways	81(67)	3 (4.5%)	0,43	0,15	-0,15	0,44	0,37	0,04	0,39
Presynaptic nicotinic acetylcholine receptors - Reactome	12(12)	1 (8.3%)	0,61	0,88	0,55	0,84	0,70	0,89	0,86
Primary bile acid biosynthesis - Mus musculus (mouse) - KEGG	16(16)	3 (18.8%)	0,06	-0,03	-0,63	0,03	-0,08	-0,32	-0,07
Prion diseases - Mus musculus (mouse) - KEGG	35(32)	3 (9.4%)	0,25	0,44	-0,20	0,39	0,22	0,31	0,35
Propanoate metabolism - Mus musculus (mouse) - KEGG	31(28)	2 (7.1%)	-0,10	0,68	-0,48	0,39	-0,24	0,50	0,27
Protein digestion and absorption - Mus musculus (mouse) - KEGG	88(76)	1 (1.3%)	0,15	0,47	0,21	0,39	0,21	0,46	0,42
Protein processing in endoplasmic reticulum - Mus musculus (mouse) - KEGG	169 (156)	1 (0.6%)	0,25	0,11	-0,21	0,29	0,17	-0,02	0,21
Pyruvate metabolism - Mus musculus (mouse) - KEGG	42(39)	2 (5.1%)	0,15	0,51	-0,27	0,34	0,05	0,29	0,25
Recycling of bile acids and salts - Reactome	12(12)	1 (8.3%)	0,37	0,43	0,01	0,48	0,34	0,23	0,42
Regulation of Cholesterol Biosynthesis by SREBP (SREBF) - Reactome	12(11)	1 (9.1%)	0,04	0,07	-0,74	0,05	-0,27	-0,55	-0,21
Regulation of Complement cascade - Reactome	24(18)	3 (16.7%)	0,46	0,42	-0,05	0,57	0,42	0,28	0,52
Regulation of IFNA signaling - Reactome	27(19)	1 (5.3%)	-0,47	0,37	-0,22	-0,19	-0,41	0,12	-0,21
Regulation of Insulin Secretion - Reactome	69(60)	1 (1.7%)	0,02	0,21	0,36	0,10	0,10	0,32	0,17
Regulation of Insulin Secretion by Acetylcholine - Reactome	10(10)	1 (10.0%)	0,19	0,29	0,80	0,27	0,36	0,51	0,38
Regulation of Insulin Secretion by Fatty Acids Bound to GPR40 (FFAR1) - Reactome	8(7)	1 (14.3%)	0,46	0,44	0,86	0,50	0,73	0,64	0,63
Regulation of Insulin Secretion by Free Fatty Acids - Reactome	8(7)	1 (14.3%)	0,46	0,44	0,86	0,50	0,73	0,64	0,63
Regulation of Insulin-like Growth Factor (IGF) Transport and Uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs) - Reactome	31(28)	1 (3.6%)	-0,22	-0,43	0,57	-0,49	-0,03	-0,33	-0,40
Regulation of thyroid hormone activity - Reactome	2(2)	1 (50.0%)	-1,00	1,00	-1,00	-1,00	-1,00	1,00	-1,00
Response to elevated platelet cytosolic Ca2+ - Reactome	92(75)	1 (1.3%)	0,27	0,11	-0,07	0,32	0,26	0,07	0,30
Retinol metabolism - Mus musculus (mouse) - KEGG	87(56)	8 (14.3%)	0,81	0,21	-0,21	0,76	0,75	0,07	0,70
Retinol metabolism - Wikipathways	39(38)	1 (2.6%)	0,41	0,37	0,16	0,54	0,43	0,35	0,53
Reversible Hydration of Carbon Dioxide - Reactome	12(11)	1 (9.1%)	0,91	0,18	0,85	0,90	0,92	0,76	0,92
RIG-I/MDA5 mediated induction of IFN-alpha/beta pathways - Reactome	45(40)	2 (5.0%)	-0,61	-0,69	-0,80	-0,74	-0,76	-0,78	-0,79
RIP-mediated NFkB activation via ZBP1 - Reactome	21(20)	2 (10.0%)	-0,73	-0,76	-0,84	-0,79	-0,82	-0,82	-0,83
Role of LAT2/NTAL/LAB on calcium mobilization - Reactome	96(87)	1 (1.1%)	0,23	0,31	0,34	0,34	0,33	0,40	0,41


Scavenging by Class B Receptors - Reactome	6(6)	2 (33.3%)	-0,82	-0,78	-0,73	-0,81	-0,78	-0,77	-0,79
Selenium metabolism-Selenoproteins - Wikipathways	47(41)	2 (4.9%)	0,16	0,09	0,01	0,17	0,14	0,07	0,16
Serotonergic synapse - Mus musculus (mouse) - KEGG	130(116)	1 (0.9%)	0,44	0,01	-0,11	0,36	0,40	-0,03	0,31
serotonin and melatonin biosynthesis - MouseCyc	5(4)	1 (25.0%)	1,00	-0,38	-0,38	0,98	1,00	-0,39	0,97
Serotonin and melatonin biosynthesis - Reactome	4(4)	1 (25.0%)	1,00	-0,38	-0,38	0,98	1,00	-0,39	0,97
SHC1 events in ERBB2 signaling - Reactome	24(24)	1 (4.2%)	0,66	0,48	0,65	0,75	0,69	0,76	0,78
SHC1 events in ERBB4 signaling - Reactome	19(19)	1 (5.3%)	0,68	0,46	0,77	0,76	0,73	0,80	0,79
SIDS Susceptibility Pathways - Wikipathways	61(57)	1 (1.8%)	0,74	0,05	0,18	0,73	0,76	0,13	0,72
Signal amplification - Reactome	16(14)	1 (7.1%)	0,50	0,36	0,64	0,50	0,66	0,62	0,63
Signal Transduction - Reactome	1870 (1160)	4 (0.3%)	-0,01	0,04	-0,12	0,01	-0,05	-0,03	-0,03
Signaling by EGFR - Reactome	161 (150)	1 (0.7%)	0,23	0,26	0,23	0,31	0,28	0,31	0,34
Signaling by EGFR in Cancer - Reactome	163 (152)	1 (0.7%)	0,21	0,25	0,21	0,30	0,26	0,29	0,32
Signaling by ERBB2 - Reactome	144 (133)	1 (0.8%)	0,20	0,24	0,20	0,27	0,24	0,27	0,30
Signaling by ERBB4 - Reactome	154 (137)	1 (0.7%)	0,25	0,23	0,25	0,32	0,31	0,30	0,36
Signaling by FGFR - Reactome	144 (132)	1 (0.8%)	0,21	0,27	0,20	0,30	0,26	0,30	0,32
Signaling by FGFR in disease - Reactome	155 (143)	1 (0.7%)	0,15	0,30	0,16	0,26	0,19	0,30	0,28
Signaling by GPCR - Reactome	1162 (502)	3 (0.6%)	-0,18	-0,08	-0,28	-0,19	-0,25	-0,20	-0,24
Signaling by PDGF - Reactome	160 (152)	1 (0.7%)	0,14	0,27	0,26	0,23	0,21	0,33	0,28
Signaling by SCF-KIT - Reactome	125 (120)	1 (0.8%)	0,14	0,31	0,25	0,25	0,21	0,35	0,30
Signaling by the B Cell Receptor (BCR) - Reactome	134 (125)	1 (0.8%)	0,18	0,15	0,27	0,22	0,26	0,25	0,29
Signalling by NGF - Reactome	275 (254)	1 (0.4%)	0,20	0,27	0,21	0,29	0,25	0,31	0,32
SLC-mediated transmembrane transport - Reactome	249 (229)	4 (1.7%)	0,28	0,27	0,17	0,36	0,30	0,29	0,36
sphingosine and sphingosine-1-phosphate metabolism - MouseCyc	9(9)	1 (11.1%)	0,46	0,66	0,10	0,59	0,43	0,50	0,54
Spinal Cord Injury - Wikipathways	103(95)	1 (1.1%)	0,10	0,26	0,14	0,24	0,13	0,28	0,24
Staphylococcus aureus infection - Mus musculus (mouse) - KEGG	52(44)	2 (4.5%)	0,19	0,07	-0,03	0,22	0,17	0,04	0,21
Starch and sucrose metabolism - Mus musculus (mouse) - KEGG	54(36)	2 (5.6%)	0,43	0,13	-0,08	0,39	0,36	0,06	0,33
Steroid biosynthesis - Mus musculus (mouse) - KEGG	17(17)	1 (5.9%)	0,54	0,22	-0,11	0,57	0,52	0,12	0,55
Steroid Biosynthesis - Wikipathways	13(11)	2 (18.2%)	0,98	0,25	0,50	0,99	0,98	0,44	0,98
Steroid hormone biosynthesis - Mus musculus (mouse) - KEGG	58(40)	7 (17.5%)	0,86	0,17	0,32	0,86	0,86	0,28	0,85
Steroid hormones - Reactome	30(25)	2 (8.0%)	0,97	0,35	0,37	0,97	0,97	0,42	0,96
Sterols are 12-hydroxylated by CYP8B1 - Reactome	1(1)	1 (100.0%)	-	-	-	-	-	-	-
Sulfur amino acid metabolism - Reactome	25(24)	3 (12.5%)	0,71	0,60	-0,09	0,75	0,65	0,47	0,71
superpathway of cholesterol biosynthesis - MouseCyc	25(24)	2 (8.3%)	0,57	0,61	-0,13	0,66	0,53	0,47	0,63
superpathway of methionine degradation - MouseCyc	19(18)	2 (11.1%)	0,49	0,21	0,01	0,53	0,48	0,17	0,51
Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) - Reactome	26(16)	1 (6.2%)	0,84	0,22	-0,28	0,80	0,77	0,01	0,73


Synthesis of bile acids and bile salts - Reactome	27(27)	3 (11.1%)	0,07	0,11	-0,47	0,09	-0,04	-0,10	0,01
Synthesis of bile acids and bile salts via 24-hydroxycholesterol - Reactome	18(18)	2 (11.1%)	-0,09	0,27	-0,54	0,10	-0,28	0,01	-0,06
Synthesis of bile acids and bile salts via 27-hydroxycholesterol - Reactome	14(14)	2 (14.3%)	0,36	0,16	-0,07	0,38	0,34	0,14	0,37
Synthesis of bile acids and bile salts via Talpha-hydroxycholesterol - Reactome	23(23)	1 (4.3%)	-0,15	0,00	-0,24	-0,12	-0,18	-0,08	-0,14
Synthesis of Leukotrienes (LT) and Eoxins (EX) - Reactome	16(13)	1 (7.7%)	0,08	0,70	0,18	0,26	0,11	0,58	0,27
Synthesis of Lipoxins (LX) - Reactome	6(5)	1 (20.0%)	-0,18	0,56	0,25	-0,13	-0,13	0,63	-0,07
Synthesis of PC - Reactome	18(16)	1 (6.2%)	0,06	0,06	0,71	0,07	0,22	0,58	0,22
Synthesis of PE - Reactome	12(10)	1 (10.0%)	0,27	0,07	0,75	0,28	0,41	0,70	0,41
Synthesis of very long-chain fatty acyl-CoAs - Reactome	13(12)	2 (16.7%)	0,88	0,22	-0,67	0,88	0,87	-0,28	0,87
Systemic lupus erythematosus - Mus musculus (mouse) - KEGG	147(56)	3 (5.4%)	-0,14	0,53	-0,22	0,04	-0,16	0,40	0,00
TAK1 activates NFkB by phosphorylation and activation of IKKs complex - Reactome	26(23)	2 (8.7%)	-0,63	-0,72	-0,88	-0,72	-0,78	-0,81	-0,79
Taurine and hypotaurine metabolism - Mus musculus (mouse) - KEGG	10(9)	1 (11.1%)	0,95	0,92	-0,71	0,98	0,92	0,84	0,99
Terminal pathway of complement - Reactome	7(6)	3 (50.0%)	0,16	0,44	-0,41	0,46	0,01	0,27	0,40
Terpenoid backbone biosynthesis - Mus musculus (mouse) - KEGG	22(18)	1 (5.6%)	0,71	0,64	-0,36	0,83	0,66	0,43	0,78
tetrapyrrole biosynthesis II - MouseCyc	5(5)	1 (20.0%)	0,98	0,65	0,32	0,99	0,96	0,57	0,97
TGF-beta signaling pathway - Mus musculus (mouse) - KEGG	82(75)	2 (2.7%)	0,59	0,25	0,37	0,65	0,66	0,34	0,67
Thrombin signalling through proteinase activated receptors (PARs) - Reactome	17(17)	1 (5.9%)	0,22	0,33	0,66	0,40	0,38	0,53	0,51
Thromboxane signalling through TP receptor - Reactome	8(8)	1 (12.5%)	0,69	0,50	0,78	0,66	0,80	0,68	0,74
thyroid hormone metabolism I (via de-iodination) - MouseCyc	3(2)	1 (50.0%)	-1,00	1,00	-1,00	-1,00	-1,00	1,00	-1,00
Thyroxine biosynthesis - Reactome	7(7)	1 (14.3%)	-0,70	0,19	-0,29	-0,37	-0,88	0,06	-0,44
TNF-alpha NF-kB Signaling Pathway - WikiPathways	184(169)	1 (0.6%)	0,30	0,34	0,00	0,41	0,28	0,27	0,38
Toll Like Receptor 10 (TLR10) Cascade - Reactome	69(61)	2 (3.3%)	-0,48	-0,61	-0,63	-0,62	-0,60	-0,69	-0,67
Toll Like Receptor 2 (TLR2) Cascade - Reactome	73(65)	2 (3.1%)	-0,41	-0,59	-0,60	-0,58	-0,54	-0,67	-0,63
Toll Like Receptor 3 (TLR3) Cascade - Reactome	89(79)	2 (2.5%)	-0,40	-0,52	-0,58	-0,55	-0,53	-0,63	-0,60
Toll Like Receptor 4 (TLR4) Cascade - Reactome	106(94)	2 (2.1%)	-0,39	-0,48	-0,56	-0,54	-0,52	-0,61	-0,60
Toll Like Receptor 5 (TLR5) Cascade - Reactome	69(61)	2 (3.3%)	-0,48	-0,61	-0,63	-0,62	-0,60	-0,69	-0,67
Toll Like Receptor 7/8 (TLR7/8) Cascade - Reactome	75(67)	2 (3.0%)	-0,47	-0,59	-0,62	-0,61	-0,60	-0,68	-0,66
Toll Like Receptor 9 (TLR9) Cascade - Reactome	79(71)	2 (2.8%)	-0,46	-0,58	-0,62	-0,60	-0,59	-0,67	-0,65
Toll Like Receptor TLR1:TLR2 Cascade - Reactome	69(61)	2 (3.3%)	-0,48	-0,61	-0,63	-0,62	-0,60	-0,69	-0,67
Toll Like Receptor TLR6:TLR2 Cascade - Reactome	73(65)	2 (3.1%)	-0,41	-0,59	-0,60	-0,58	-0,54	-0,67	-0,63
Toll-Like Receptors Cascades - Reactome	126(114)	2 (1.8%)	-0,32	-0,44	-0,54	-0,48	-0,45	-0,58	-0,55
Toxoplasmosis - Mus musculus (mouse) - KEGG	113(105)	1 (1.0%)	-0,13	0,32	-0,11	0,04	-0,15	0,22	0,01
TRAF6 mediated induction of NFkB and MAP kinases upon TLR7/8 or 9 activation - Reactome	73(65)	2 (3.1%)	-0,46	-0,60	-0,63	-0,61	-0,60	-0,69	-0,66
TRAF6 Mediated Induction of proinflammatory cytokines - Reactome	63(57)	2 (3.5%)	-0,50	-0,61	-0,64	-0,63	-0,62	-0,69	-0,68
TRAF6 mediated NF-kB activation - Reactome	24(22)	2 (9.1%)	-0,76	-0,74	-0,85	-0,81	-0,85	-0,81	-0,84
trans-Golgi Network Vesicle Budding - Reactome	61(52)	1	0,62	0,15	-0,43	0,69	0,50	-0,09	0,56


tome		(1.9%)									
Transmembrane transport of small molecules - Reactome	499 (443)	4 (0.9%)	0,21	0,29	0,13	0,31	0,23	0,28	0,32		
Transmission across Chemical Synapses - Reactome	178 (158)	1 (0.6%)	0,14	0,29	0,21	0,26	0,19	0,32	0,29		
Transport of connexins along the secretory pathway - Reactome	3(3)	1 (33.3%)	0,00	-0,87	0,86	-0,25	0,13	-0,54	-0,13		
Transport of connexons to the plasma membrane - Reactome	2(2)	1 (50.0%)	1,00	-1,00	1,00	-1,00	1,00	-1,00	-1,00		
Transport of glucose and other sugars, bile salts and organic acids, metal ions and amine compounds - Reactome	95(91)	2 (2.2%)	0,26	0,25	0,24	0,37	0,33	0,31	0,41		
Transport of inorganic cations/anions and amino acids/oligopeptides - Reactome	93(85)	1 (1.2%)	0,35	0,23	0,24	0,46	0,38	0,35	0,48		
Transport of organic anions - Reactome	12(11)	1 (9.1%)	0,46	0,52	-0,03	0,64	0,46	0,36	0,60		
Transport of vitamins, nucleosides, and related molecules - Reactome	33(28)	1 (3.6%)	0,32	0,38	0,02	0,42	0,32	0,28	0,40		
TRIF-mediated TLR3/TLR4 signaling - Reactome	89(79)	2 (2.5%)	-0,40	-0,52	-0,58	-0,55	-0,53	-0,63	-0,60		
Triglyceride Biosynthesis - Reactome	36(33)	5 (15.2%)	0,70	0,30	-0,14	0,71	0,69	0,04	0,70		
Tryptophan metabolism - Mus musculus (mouse) - KEGG	47(39)	2 (5.1%)	0,60	0,35	-0,01	0,67	0,57	0,29	0,64		
Tryptophan metabolism - Wikipathways	44(41)	2 (4.9%)	0,66	0,06	0,07	0,65	0,66	0,08	0,65		
Type II diabetes mellitus - Mus musculus (mouse) - KEGG	50(48)	1 (2.1%)	0,05	0,24	-0,16	0,14	-0,02	0,08	0,07		
Tyrosine metabolism - Mus musculus (mouse) - KEGG	44(39)	4 (10.3%)	0,63	0,44	0,33	0,75	0,67	0,48	0,76		
Valine, leucine and isoleucine degradation - Mus musculus (mouse) - KEGG	53(47)	1 (2.1%)	0,39	0,34	0,16	0,45	0,40	0,35	0,45		
Vascular smooth muscle contraction - Mus musculus (mouse) - KEGG	128 (110)	1 (0.9%)	0,65	0,20	-0,04	0,65	0,62	0,14	0,60		
Vitamin B5 (pantothenate) metabolism - Reactome	9(9)	1 (11.1%)	-0,29	0,57	-0,86	-0,10	-0,49	-0,39	-0,33		
Vitamin B6 metabolism - Mus musculus (mouse) - KEGG	10(8)	1 (12.5%)	0,57	0,88	0,40	0,76	0,61	0,90	0,78		
Xenobiotics - Reactome	41(25)	3 (12.0%)	0,61	0,01	-0,40	0,55	0,51	-0,30	0,48		
XPoNet - protein-protein interactions in the podocyte expanded by STRING - Wikipathways	831(77 9)	1 (0.1%)	0,16	0,17	0,15	0,22	0,18	0,21	0,24		
ZBP1(DAI) mediated induction of type I IFNs - Reactome	31(28)	2 (7.1%)	-0,65	-0,72	-0,83	-0,75	-0,78	-0,81	-0,80		

U: Transcription factor binding sites (TFBs) with a z-score > 5 or a Fischer-score > 5 addressed by the upregulated CR genes and the overlap with the phytonutrients

(TF = transcription factor, R = resveratrol, P = phloridzin, Qd= quercetin (diet), Qi = quercetin (intraperitoneal), Gi = genistein (intraperitoneal))







V: Transcription factor binding sites (TFBs) with a z-score > 5 or a Fischer-score > 5 addressed by the downregulated CR genes and the overlap with the phytonutrients
 (TF = transcription factor, R = resveratrol, P = phloridzin, Qd= quercetin (diet), Qi = quercetin (intraperitoneal), Gi = genistein (intraperitoneal))

JASPAR ID	TF	R	P	Qd	Qi	Gi
MA0099.2	AP1					
MA0151.1	ARID3A					
PB0002.1	Arid5a_1					
MA0006.1	Arnt::Ahr	■		■	■	
PB0004.1	Atf1_1					
PB0007.1	Bhlhb2_1					
MA0102.2	CEBPA					
PB0113.1	E2F3_2	■	■			
PB0012.1	Elf3_1		■			
MA0136.1	ELF5			■		
PB0014.1	Esrra_1	■			■	
MA0148.1	FOXA1					
MA0047.2	Foxa2					
PB0015.1	Foxa2_1					
MA0031.1	FOXD1					
MA0041.1	Foxd3					
MA0042.1	FOXI1					
PB0016.1	Foxj1_1					
PB0017.1	Foxj3_1					
PB0121.1	Foxj3_2			■		
PB0018.1	Foxk1_1				■	
PB0122.1	Foxk1_2	■			■	
PB0019.1	Foxl1_1					
PB0123.1	Foxl1_2					
MA0157.1	FOXO3					■
MA0040.1	Foxq1					
MA0062.2	GABPA	■			■	
PB0020.1	Gabpa_1		■	■		
PB0021.1	Gata3_1			■	■	
PB0022.1	Gata5_1					
PB0028.1	Hbp1_1			■		
MA0043.1	HLF					
MA0046.1	HNF1A					
MA0153.1	HNF1B					
MA0114.1	HNF4A	■				■
PB0134.1	Hnf4a_2					
PB0135.1	Hoxa3_2					
MA0158.1	HOXA5					
PB0032.1	IRC900814_1					
MA0050.1	IRF1					
MA0051.1	IRF2					
PB0035.1	Irf5_1					
PB0037.1	Isgf3g_1					
PB0143.1	Klf7_2	■	■	■	■	
MA0135.1	Lhx3			■	■	
PB0145.1	Mafb_2					
PB0043.1	Max_1	■	■			
PB0147.1	Max_2	■	■			
MA0052.1	MEF2A			■		
MA0056.1	MZF1_1-4	■				
MA0152.1	NFATC2			■		



W: All mentioned functions / biological processes revealed by UniProt database for the transcription factors (TF), which address the top TFBs of caloric restriction

TF	Function	biological process
Ahr	Ligand-activated transcriptional activator. Binds to the XRE promoter region of genes it activates. Activates the expression of multiple phase I and II xenobiotic chemical metabolizing enzyme genes (such as the CYP1A1 gene). Mediates biochemical and toxic effects of halogenated aromatic hydrocarbons. Involved in	B cell differentiation; B (B-1) cell homeostasis; blood circulation; blood vessel development / morphogenesis / remodeling, camera-type eye development, cardiac left ventricle morphogenesis, cell cycle, cell morphogenesis, cellular response to cAMP, circadian regulation of gene expression, circumferential growth involved in left ventricle morphogenesis, common bile duct development, embryonic hemo-

	cell-cycle regulation. Likely to play an important role in the development and maturation of many tissues. Regulates the circadian clock by inhibiting the basal and circadian expression of the core circadian component PER1. Inhibits PER1 by repressing the CLOCK-ARNTL/BMAL1 heterodimer mediated transcriptional activation of PER1.	poiesis, gland development, glomerulus morphogenesis, immune system process, intracellular receptor signaling pathway, kidney morphogenesis, liver development, lymphocyte homeostasis, negative regulation of necrotic cell death, negative regulation of systemic arterial blood pressure, negative regulation of transcription from RNA polymerase II promoter, negative regulation of vasoconstriction, patterning of blood vessels, positive regulation of cell size, positive regulation of growth rate, positive regulation of RNA polymerase II transcriptional preinitiation complex assembly, post-embryonic hemopoiesis, prostate gland development, regulation of B cell proliferation, regulation of blood pressure, regulation of blood vessel size, regulation of gene expression, regulation of heart growth, regulation of transcription, DNA-templated, reproductive structure development, response to stress, response to toxic substance, response to xenobiotic stimulus, smooth muscle tissue development, spleen development, T cell homeostasis, venous blood vessel development, xenobiotic metabolic process
Arid5a	DNA-binding protein that may regulate transcription and act as a repressor by binding to AT-rich stretches in the promoter region of target genes	chondrocyte differentiation, positive regulation of histone acetylation, positive regulation of transcription from RNA polymerase II promoter, transcription, DNA-templated
Arnt	Required for activity of the Ah (dioxin) receptor. This protein is required for the ligand-binding subunit to translocate from the cytosol to the nucleus after ligand binding. The complex then initiates transcription of genes involved in the activation of PAH procarcinogens. The heterodimer with HIF1A or EPAS1/HIF2A functions as a transcriptional regulator of the adaptive response to hypoxia	cell differentiation, embryonic placenta development, intracellular receptor signaling pathway, positive regulation of hormone biosynthetic process, positive regulation of protein sumoylation, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, positive regulation of vascular endothelial growth factor production, regulation of transcription from RNA polymerase II promoter in response to oxidative stress, response to hypoxia, response to toxic substance, transcription, DNA-templated
Cebpa	Transcription factor that coordinates proliferation arrest and the differentiation of myeloid progenitors, adipocytes, hepatocytes, and cells of the lung and the placenta; Binds directly to the consensus DNA sequence 5'-T[TG]NNGNAA[TG]-3' acting as an activator on distinct target genes. During early embryogenesis, plays essential and redundant functions with CEBPB. Essential for the transition from common myeloid progenitors (CMP) to granulocyte/monocyte progenitors (GMP). Critical for the proper development of the liver and the lung. Necessary for terminal adipocyte differentiation, is required for postnatal maintenance of systemic energy homeostasis and lipid storage. To regulate these different processes at the proper moment and tissue, interplays with other transcription factors and modulators. Downregulates the expression of genes that maintain cells in an undifferentiated and proliferative state through E2F1 repression, which is critical for its ability to induce adipocyte and granulocyte terminal differentiation. Reciprocally E2F1 blocks adipocyte differentiation by binding to specific promoters and repressing CEBPA binding to its target gene promoters. Proliferation arrest also depends on a func-	brown fat cell differentiation; cell maturation, cellular response to lithium ion, cellular response to organic cyclic compound, cholesterol metabolic process, cytokine-mediated signaling pathway, embryonic placenta development, fat cell differentiation, glucose homeostasis, granulocyte differentiation, inner ear development, lipid homeostasis, liver development, lung development, macrophage differentiation, mitochondrion organization, myeloid cell differentiation, negative regulation of cell cycle, negative regulation of cell proliferation, negative regulation of transcription, DNA-templated, negative regulation of transcription from RNA polymerase II promoter, Notch signaling pathway, positive regulation of fat cell differentiation, positive regulation of osteoblast differentiation, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase III promoter, positive regulation of transcription from RNA polymerase II promoter, regulation of cell proliferation, regulation of transcription, DNA-templated, regulation of transcription from RNA polymerase II promoter, transcription, DNA-templated, transcription from RNA polymerase II promoter, transcription from RNA polymerase I promoter, urea cycle, white fat cell differentiation

	tional binding to SWI/SNF complex. In liver, regulates gluconeogenesis and lipogenesis through different mechanisms. To regulate gluconeogenesis, functionally cooperates with FOXO1 binding to IRE-controlled promoters and regulating the expression of target genes such as PCK1 or G6PC. To modulate lipogenesis, interacts and transcriptionally synergizes with SREBF1 in promoter activation of specific lipogenic target genes such as ACAS2. In adipose tissue, seems to act as FOXO1 coactivator accessing to ADIPOQ promoter through FOXO1 binding sites	
Esrrb	Nuclear receptor, may regulate ESR1 transcriptional activity. Induces the expression of PERM1 in the skeletal muscle	embryonic placenta development, in utero embryonic development, positive regulation of transcription from RNA polymerase II promoter, regulation of transcription, DNA-templated, stem cell population maintenance, trophectodermal cell proliferation / morphogenesis
Fos	Nuclear phosphoprotein which forms a tight but non-covalently linked complex with the JUN/AP-1 transcription factor. On TGF-beta activation, forms a multimeric SMAD3/SMAD4/JUN/FOS complex, at the AP1/SMAD-binding site to regulate TGF-beta-mediated signaling. Has a critical function in regulating the development of cells destined to form and maintain the skeleton. It is thought to have an important role in signal transduction, cell proliferation and differentiation. In growing cells, activates phospholipid synthesis, possibly by activating CDS1 and PI4K2A. This activity requires Tyr-dephosphorylation and association with the endoplasmic reticulum.	ageing, cellular response to calcium ion, cellular response to extracellular stimulus, cellular response to hormone stimulus, cellular response to reactive oxygen species, conditioned taste aversion, female pregnancy, nervous system development, positive regulation of osteoclast differentiation, positive regulation of pri-miRNA transcription from RNA polymerase II promoter, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, response to cAMP, response to cold, response to corticosterone, response to cytokine, response to drug, response to gravity, response to immobilization stress, response to light stimulus, response to lipopolysaccharide, response to muscle stretch, response to progesterone, response to toxic substance, skeletal muscle cell differentiation, sleep, SMAD protein signal transduction, transforming growth factor beta receptor signaling pathway
Foxd3	Binds to the consensus sequence 5'-A[AT]T[AG]TTTGTAA-3' and acts as a transcriptional repressor. Also acts as a transcriptional activator. Promotes development of neural crest cells from neural tube progenitors. Restricts neural progenitor cells to the neural crest lineage while suppressing interneuron differentiation. Required for maintenance of pluripotent cells in the pre-implantation and peri-implantation stages of embryogenesis	embryonic placenta development, in utero embryonic development, negative regulation of transcription from RNA polymerase II promoter, positive regulation of transcription from RNA polymerase II promoter, transcription, DNA-templated, trophectodermal cell differentiation
Foxj1	Transcription factor required for motile ciliogenesis	actin cytoskeleton organization, activation of GTPase activity, brain development, cell differentiation, central tolerance induction, cilium assembly, cilium morphogenesis, determination of left/right symmetry, epithelium development, establishment of apical/basal cell polarity, glomerular parietal epithelial cell development, heart development, humoral immune response, left/right pattern formation, leukocyte migration, metanephric part of ureteric bud development, negative regulation of B cell activation, negative regulation of germinal center formation, negative regulation of humoral immune response mediated by circulating immunoglobulin, negative regulation of interleukin-6 biosynthetic process, negative regulation of NF-kappaB transcription factor activity, negative regulation of T cell differentia-

		tion in thymus, negative regulation of T cell proliferation, negative regulation of transcription from RNA polymerase II promoter, positive regulation of central B cell tolerance induction, positive regulation of lung ciliated cell differentiation, positive regulation of transcription from RNA polymerase II promoter, regulation of epithelial cell differentiation, transcription from RNA polymerase II promoter
Foxj3	Transcriptional activator of MEF2C involved in the regulation of adult muscle fiber type identity and skeletal muscle regeneration	anatomical structure morphogenesis; cell differentiation; regulation of transcription from RNA polymerase II promoter; transcription, DNA-templated
Foxl1	Transcription factor required for proper proliferation and differentiation in the gastrointestinal epithelium. Target gene of the hedgehog (Hh) signaling pathway via GLI2 AND GLI3 transcription factors.	cell differentiation, heart development, Peyer's patch morphogenesis, proteoglycan biosynthetic process, regulation of transcription, DNA-templated, regulation of transcription from RNA polymerase II promoter, regulation of Wnt signaling pathway, transcription, DNA-templated, visceral mesoderm-endoderm interaction involved in midgut development
Hnf1a	Transcriptional activator that regulates the tissue specific expression of multiple genes, especially in pancreatic islet cells and in liver. Required for the expression of several liver specific genes. Binds to the inverted palindrome 5'- GTTAATNATTAAAC-3'.	bile acid and bile salt transport; bile acid biosynthetic process, blastocyst development, bone resorption, cholesterol metabolic process, chromatin remodeling, embryonic limb morphogenesis, endocrine pancreas development, fatty acid biosynthetic process, fatty acid transport, glucose homeostasis, glucose import, heme biosynthetic process, histone acetylation, insulin secretion, liver development, negative regulation of transcription from RNA polymerase II promoter, paraxial mesoderm formation, placenta development, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, positive regulation of transcription initiation from RNA polymerase II promoter, protein localization, regulation of hormone secretion, regulation of insulin secretion, regulation of transcription, DNA-templated, regulation of transcription from RNA polymerase II promoter, regulation of Wnt signaling pathway, renal glucose absorption, reproductive structure development, response to glucose, response to oxidative stress, reverse cholesterol transport, SMAD protein signal transduction
Hoxa5	Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'- CYYNATT[TG]Y-3'	anterior/posterior pattern specification; bronchiole development, cartilage morphogenesis, cell-cell signaling involved in mammary gland development, cell migration, embryonic skeletal system development, embryonic skeletal system morphogenesis, epithelial tube branching involved in lung morphogenesis, intestinal epithelial cell maturation, lobar bronchus epithelium development, lung alveolus development, lung-associated mesenchyme development, lung development, lung goblet cell differentiation, mammary gland alveolus development, mammary gland epithelial cell differentiation, mesenchymal-epithelial cell signaling, morphogenesis of an epithelium, multicellular organism growth, negative regulation of angiogenesis, negative regulation of erythrocyte differentiation, pattern specification process, positive regulation of apoptotic process, positive regulation of myeloid cell differentiation, positive regulation of receptor biosynthetic process, positive regulation of transcription from RNA polymerase II promoter, regulation of mammary gland epithelial cell proliferation, respiratory gaseous exchange, respiratory system process, skeletal system development, thyroid gland development, trachea

		cartilage morphogenesis, trachea morphogenesis, transcription from RNA polymerase II promoter
Jdp2	Component of the AP-1 transcription factor that represses transactivation mediated by the Jun family of proteins. Involved in a variety of transcriptional responses associated with AP-1, such as UV-induced apoptosis, cell differentiation, tumorigenesis and antitumorigenesis. Can also function as a repressor by recruiting histone deacetylase 3/HDAC3 to the promoter region of JUN. May control transcription via direct regulation of the modification of histones and the assembly of chromatin	negative regulation of fat cell differentiation, negative regulation of transcription from RNA polymerase II promoter, positive regulation of histone deacetylation, transcription, DNA-templated
Jun	Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'. Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation. Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells. Binds to the USP28 promoter in colorectal cancer (CRC) cells	ageing, angiogenesis, axon regeneration, cellular process, cellular response to calcium ion, cellular response to hormone stimulus, cellular response to potassium ion starvation, circadian rhythm, eyelid development in camera-type eye, leading edge cell differentiation, learning, liver development, membrane depolarization, microglial cell activation, monocyte differentiation, negative regulation by host of viral transcription, negative regulation of apoptotic process, negative regulation of cell proliferation, negative regulation of DNA binding, negative regulation of neuron apoptotic process, negative regulation of protein autophosphorylation, negative regulation of transcription, DNA-templated, negative regulation of transcription from RNA polymerase II promoter, negative regulation of transcription from RNA polymerase II promoter in response to endoplasmic reticulum stress, outflow tract morphogenesis, positive regulation by host of viral transcription, positive regulation of cell differentiation, positive regulation of DNA replication, positive regulation of DNA-templated transcription, initiation, positive regulation of endothelial cell proliferation, positive regulation of epithelial cell migration, positive regulation of ERK1 and ERK2 cascade, positive regulation of fibroblast proliferation, positive regulation of GTPase activity, positive regulation of monocyte differentiation, positive regulation of neuron apoptotic process, positive regulation of pri-miRNA transcription from RNA polymerase II promoter, positive regulation of smooth muscle cell proliferation, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, regulation of cell cycle, regulation of transcription, DNA-templated, release of cytochrome c from mitochondria, response to cAMP, response to cytokine, response to drug, response to hydrogen peroxide, response to lipopolysaccharide, response to muscle stretch, response to radiation, SMAD protein import into nucleus, SMAD protein signal transduction, transforming growth factor beta receptor signaling pathway
Klf4	Transcription factor; can act both as activator and as repressor. Binds the 5'-CACCC-3' core sequence. Binds to the promoter region of its own gene and can activate its own transcription. Regulates the expression of key transcription factors during embryonic development. Plays an important role in maintaining embryonic stem cells, and in preventing their differentiation. Required for establishing the barrier function of the skin and for postnatal	canonical Wnt signaling pathway, cell differentiation, cellular response to cycloheximide, cellular response to growth factor stimulus, cellular response to hydrogen peroxide, cellular response to laminar fluid shear stress, cellular response to peptide, cellular response to retinoic acid, epidermal cell differentiation, epidermis morphogenesis, fat cell differentiation, negative regulation of cell migration, negative regulation of cell migration involved in sprouting angiogenesis, negative regulation of cell proliferation, negative regulation of chemokine (C-X-C motif)

	maturation and maintenance of the ocular surface. Involved in the differentiation of epithelial cells and may also function in skeletal and kidney development. Contributes to the down-regulation of p53/TP53 transcription	ligand 2 production, negative regulation of cysteine-type endopeptidase activity involved in apoptotic process, negative regulation of ERK1 and ERK2 cascade, negative regulation of heterotypic cell-cell adhesion, negative regulation of interleukin-8 biosynthetic process, negative regulation of muscle hyperplasia, negative regulation of NF-kappaB transcription factor activity, negative regulation of phosphatidylinositol 3-kinase signaling, negative regulation of protein kinase B signaling, negative regulation of response to cytokine stimulus, negative regulation of sequence-specific DNA binding transcription factor activity, negative regulation of smooth muscle cell proliferation, negative regulation of transcription, DNA-templated, negative regulation of transcription from RNA polymerase II promoter, positive regulation of cellular protein metabolic process, positive regulation of hemoglobin biosynthetic process, positive regulation of nitric oxide biosynthetic process, positive regulation of protein metabolic process, positive regulation of telomerase activity, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, positive regulation of transcription regulatory region DNA binding, post-embryonic camera-type eye development, post-embryonic hemopoiesis, regulation of axon regeneration, regulation of cell differentiation, regulation of cell proliferation, regulation of transcription from RNA polymerase II promoter, response to organic substance, response to retinoic acid, stem cell population maintenance, transcription, DNA-templated, transcription from RNA polymerase II promoter
Klf7	Transcriptional activator. Binds specifically to an Ikaros core binding element that is crucial for in vivo NTRK1/TrkA enhancer function	axon guidance, axonogenesis, dendrite morphogenesis, positive regulation of transcription, DNA-templated, transcription, DNA-templated
Max	Transcription regulator. Forms a sequence-specific DNA-binding protein complex with MYC or MAD which recognizes the core sequence 5'-CAC[GA]TG-3'. The MYC:MAX complex is a transcriptional activator, whereas the MAD:MAX complex is a repressor. CpG methylation of the recognition site greatly inhibits DNA binding, suggesting that DNA methylation may regulate the MYC:MAX complex in vivo. May repress transcription via the recruitment of a chromatin remodeling complex containing H3 'Lys-9' histone methyltransferase activity	regulation of transcription, DNA-templated, transcription, DNA-templated
Myc	Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Activates the transcription of growth-related genes	activation of cysteine-type endopeptidase activity involved in apoptotic process, B cell apoptotic process, branching involved in ureteric bud morphogenesis, canonical Wnt signaling pathway, canonical Wnt signaling pathway involved in negative regulation of apoptotic process, canonical Wnt signaling pathway involved in positive regulation of apoptotic process, cell cycle arrest, cellular iron ion homeostasis, cellular response to DNA damage stimulus, cellular response to drug, cellular response to interferon-alpha, chromatin remodeling, chromosome organization, detection of mechanical stimulus involved in sensory perception of sound, intrinsic apoptotic signaling pathway in response to DNA damage, MAPK cascade, middle ear morphogenesis, negative regulation of cell division, negative

		regulation of fibroblast proliferation, negative regulation of monocyte differentiation, negative regulation of protein binding, negative regulation of transcription from RNA polymerase II promoter, pigmentation, positive regulation of apoptotic signaling pathway, positive regulation of B cell apoptotic process, positive regulation of catalytic activity, positive regulation of cell proliferation, positive regulation of cysteine-type endopeptidase activity involved in apoptotic process, positive regulation of DNA biosynthetic process, positive regulation of epithelial cell proliferation, positive regulation of fibroblast proliferation, positive regulation of mesenchymal cell proliferation, positive regulation of metanephric cap mesenchymal cell proliferation, positive regulation of response to DNA damage stimulus, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, protein processing, regulation of apoptotic process, regulation of gene expression, regulation of telomere maintenance, regulation of transcription, DNA-templated, response to alkaloid, response to gamma radiation, response to radiation, skeletal muscle cell differentiation, skeletal system morphogenesis, transcription from RNA polymerase II promoter, Wnt signaling pathway
Mycn		branching morphogenesis of an epithelial tube, cartilage condensation, embryonic digit morphogenesis, embryonic skeletal system morphogenesis, lung development, negative regulation of astrocyte differentiation, negative regulation of reactive oxygen species metabolic process, positive regulation of cell death, positive regulation of cell proliferation, positive regulation of mesenchymal cell proliferation, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, regulation of auditory receptor cell differentiation
Mzf1		negative regulation of transcription from RNA polymerase II promoter; positive regulation of transcription from RNA polymerase II promoter, regulation of transcription, DNA-templated, transcription from RNA polymerase II promoter
Nfyα	Component of the sequence-specific heterotrimeric transcription factor (NF-Y) which specifically recognizes a 5'-CCAAT-3' box motif found in the promoters of its target genes. NF-Y can function as both an activator and a repressor, depending on its interacting cofactors. NF-YA positively regulates the transcription of the core clock component ARNTL/BMAL1.	positive regulation of stem cell proliferation, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, regulation of stem cell population maintenance, regulation of transcription, DNA-templated, rhythmic process, transcription from RNA polymerase II promoter
Nkx2-5	Implicated in commitment to and/or differentiation of the myocardial lineage. Acts as a transcriptional activator of ANF in cooperation with GATA4. It is transcriptionally controlled by PBX1 and acts as a transcriptional repressor of CDKN2B. Together with PBX1, it is required for spleen development through a mechanism that involves CDKN2B repression.	adult heart development, apoptotic process involved in heart morphogenesis, atrial cardiac muscle cell development, atrial septum morphogenesis, atrioventricular node cell development, atrioventricular node cell fate commitment, atrioventricular node development, BMP signaling pathway, bundle of His development, canonical Wnt signaling pathway, cardiac conduction system development, cardiac muscle cell differentiation, cardiac muscle cell proliferation, cardiac muscle contraction, cardiac muscle tissue development, cardiac muscle tissue morphogenesis, cardiac ventricle formation, cardiac ventricle morphogenesis, cardiovascular system development, cell differentiation, embryonic heart tube

		development, embryonic heart tube left/right pattern formation, endoderm development, heart contraction, heart development, heart looping, heart morphogenesis, heart trabecula formation, hemopoiesis, negative regulation of apoptotic process, negative regulation of canonical Wnt signaling pathway, negative regulation of cardiac muscle cell apoptotic process, negative regulation of myotube differentiation, negative regulation of transcription from RNA polymerase II promoter, outflow tract morphogenesis, outflow tract septum morphogenesis, pharyngeal system development, positive regulation of cardioblast differentiation, positive regulation of cell proliferation, positive regulation of gene expression, positive regulation of heart contraction, positive regulation of neuron differentiation, positive regulation of sodium ion transport, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, positive regulation of transcription initiation from RNA polymerase II promoter, positive regulation of transcription via serum response element binding, proepicardium development, pulmonary myocardium development, Purkinje myocyte differentiation, regulation of cardiac conduction, regulation of cardiac muscle cell proliferation, regulation of cardiac muscle contraction, regulation of transcription from RNA polymerase II promoter, right ventricular cardiac muscle tissue morphogenesis, sarcomere organization, septum secundum development, spleen development, thyroid gland development, tongue development, transcription from RNA polymerase II promoter, vasculogenesis, ventricular cardiac muscle cell development, ventricular cardiac myofibril assembly, ventricular septum morphogenesis, ventricular trabecula myocardium morphogenesis
Nkx3-2	Transcriptional repressor that acts as a negative regulator of chondrocyte maturation. PLays a role in distal stomach development; required for proper antral-pyloric morphogenesis and development of antral-type epithelium. In concert with GSC, defines the structural components of the middle ear; required for tympanic ring and gonium development and in the regulation of the width of the malleus	animal organ development, determination of left/right symmetry, digestive system development, embryonic skeletal system development, intestinal epithelial cell development, middle ear morphogenesis, negative regulation of apoptotic process, negative regulation of chondrocyte differentiation, negative regulation of transcription from RNA polymerase II promoter, organ formation, pancreas development, skeletal system development, skeletal system morphogenesis, spleen development, transcription, DNA-templated
Pparg	Nuclear receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the nuclear receptor binds to DNA specific PPAR response elements (PPRE) and modulates the transcription of its target genes, such as acyl-CoA oxidase. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Key regulator of adipocyte differentiation and glucose homeostasis. ARF6 acts as a key regulator of the tissue-specific adipocyte P2 (aP2) enhancer. Acts as a critical regulator of gut homeostasis by suppressing NF-kappa-B-mediated proinflammatory responses. Plays a role in the regulation of cardiovascular circadian rhythms by regulating the transcription of ARNTL/BMAL1 in the blood vessels	activation of cysteine-type endopeptidase activity involved in apoptotic process, brown fat cell differentiation, cell fate commitment, cell maturation, cellular response to insulin stimulus, cellular response to lithium ion, cellular response to organic cyclic compound, diet induced thermogenesis, epithelial cell differentiation, fat cell differentiation, glucose homeostasis, inflammatory response, lipoprotein transport, long-chain fatty acid transport, low-density lipoprotein particle receptor biosynthetic process, monocyte differentiation, negative regulation of cell proliferation, negative regulation of cellular response to insulin stimulus, negative regulation of cholesterol storage, negative regulation of cytokine production, negative regulation of interferon-gamma-mediated signaling pathway, negative regulation of macrophage derived foam cell differentiation, negative regulation of peptide hormone secretion, negative regulation of receptor biosynthetic process, negative regulation of sequestering of triglyceride, negative

		regulation of smooth muscle cell proliferation, negative regulation of transcription, DNA-templated, negative regulation of transcription from RNA polymerase II promoter, peroxisome proliferator activated receptor signaling pathway, placenta development, positive regulation of fat cell differentiation, positive regulation of sequence-specific DNA binding transcription factor activity, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, regulation of blood pressure, regulation of circadian rhythm, regulation of fat cell differentiation, regulation of gene expression, regulation of transcription, DNA-templated, regulation of transcription involved in cell fate commitment, response to dietary excess, response to food, response to light stimulus, response to lipid, response to low-density lipoprotein particle, response to retinoic acid, rhythmic process, signal transduction, transcription from RNA polymerase II promoter, white fat cell differentiation
Rxra	Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes. The RAR/RXR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. The high affinity ligand for RXRs is 9-cis retinoic acid. RXRA serves as a common heterodimeric partner for a number of nuclear receptors. The RXR/RAR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. In the absence of ligand, the RXR-RAR heterodimers associate with a multiprotein complex containing transcription corepressors that induce histone acetylation, chromatin condensation and transcriptional suppression. On ligand binding, the corepressors dissociate from the receptors and associate with the coactivators leading to transcriptional activation. The RXRA/PPARA heterodimer is required for PPARA transcriptional activity on fatty acid oxidation genes such as ACOX1 and the P450 system genes.	ageing, angiogenesis involved in coronary vascular morphogenesis, axon regeneration, camera-type eye development, cardiac muscle cell differentiation, cardiac muscle cell proliferation, cellular response to insulin stimulus, embryo implantation, heart development, heart morphogenesis, inflammatory response, in utero embryonic development, liver development, maternal placenta development, mesenchyme development, midgut development, modulation by virus of host morphology or physiology, negative regulation of cell proliferation, negative regulation of gene expression, negative regulation of transcription from RNA polymerase II promoter, peroxisome proliferator activated receptor signaling pathway, placenta development, positive regulation of apoptotic process, positive regulation of transcription, DNA-templated, positive regulation of transcription from RNA polymerase II promoter, positive regulation of transcription from RNA polymerase II promoter involved in cellular response to chemical stimulus, positive regulation of translational initiation by iron, protein homotetramerization, regulation of branching involved in prostate gland morphogenesis, regulation of myelination, regulation of transcription, DNA-templated, regulation of transcription from RNA polymerase II promoter, response to ethanol, response to glucocorticoid, response to retinoic acid, response to selenium ion, response to vitamin A, response to vitamin D, retinoic acid receptor signaling pathway, secretory columnar luminal epithelial cell differentiation involved in prostate glandular acinus development, transcription, DNA-templated, ventricular cardiac muscle cell differentiation, ventricular cardiac muscle tissue morphogenesis, visceral serous pericardium development
Sp1	Transcription factor that can activate or repress transcription in response to physiological and pathological stimuli. Binds with high affinity to GC-rich motifs and regulates the expression of a large number of genes involved in a variety of processes such as cell growth, apoptosis, differentiation and immune responses. Highly regulated by post-translational modifications (phosphorylations, sumoylation, proteolytic cleavage, glycosylation and acetylation). Binds also the PDGFR-	definitive hemopoiesis, embryonic camera-type eye morphogenesis, embryonic placenta development, embryonic process involved in female pregnancy, embryonic skeletal system development, enucleate erythrocyte differentiation, in utero embryonic development, liver development, lung development, megakaryocyte differentiation, mRNA transcription from RNA polymerase II promoter, ossification, positive regulation by host of viral transcription, positive regulation of hydrogen sulfide biosynthetic process, positive regulation of transcription, DNA-templated, positive regulation of transcription from

	alpha G-box promoter. May have a role in modulating the cellular response to DNA damage. Implicated in chromatin remodeling. Plays a role in the recruitment of SMARCA4/BRG1 on the c-FOS promoter Plays an essential role in the regulation of FE65 gene expression (By similarity). Positively regulates the transcription of the core clock component ARNTL/BMAL1.	RNA polymerase II promoter, regulation of transcription, DNA-templated, rhythmic process, trophectodermal cell differentiation
Spib	Sequence specific transcriptional activator which binds to the PU-box, a purine-rich DNA sequence (5'-GAGGAA-3') that can act as a lymphoid-specific enhancer. Promotes development of plasmacytoid dendritic cells (pDCs), also known as type 2 DC precursors (pre-DC2) or natural interferon (IFN)-producing cells. These cells have the capacity to produce large amounts of interferon and block viral replication. Required for B-cell receptor (BCR) signaling, which is necessary for normal B-cell development and antigenic stimulation.	macrophage differentiation, regulation of transcription, DNA-templated, regulation of transcription from RNA polymerase II promoter, transcription, DNA-templated
Sry	Transcriptional regulator that controls a genetic switch in male development. It is necessary and sufficient for initiating male sex determination by directing the development of supporting cell precursors (pre-Sertoli cells) as Sertoli rather than granulosa cells. In male adult brain involved in the maintenance of motor functions of dopaminergic neurons (By similarity). Involved in different aspects of gene regulation including promoter activation or repression. SRY HMG box recognizes DNA by partial intercalation in the minor groove. Promotes DNA bending. Also involved in pre-mRNA splicing (By similarity). Binds to the DNA consensus sequence 5'-[AT]AACAA[AT]-3'.	cell differentiation, male gonad development, male sex determination, negative regulation of gene expression, negative regulation of transcription from RNA polymerase II promoter, positive regulation of male gonad development, positive regulation of transcription, DNA-templated, sex determination, transcription, DNA-templated
Tbp	General transcription factor that functions at the core of the DNA-binding multiprotein factor TFIID. Binding of TFIID to the TATA box is the initial transcriptional step of the pre-initiation complex (PIC), playing a role in the activation of eukaryotic genes transcribed by RNA polymerase II. Component of the transcription factor SL1/TIF-IB complex, which is involved in the assembly of the PIC (preinitiation complex) during RNA polymerase I-dependent transcription. The rate of PIC formation probably is primarily dependent on the rate of association of SL1 with the rDNA promoter. SL1 is involved in stabilization of nucleolar transcription factor 1/UBTF on rDNA	DNA-templated transcription, initiation, positive regulation of transcription, DNA-templated, spermatogenesis, transcription from RNA polymerase III promoter, transcription from RNA polymerase II promoter
Yy1	Multifunctional transcription factor that exhibits positive and negative control on a large number of cellular and viral genes by binding to sites overlapping the transcription start site. Binds to the consensus sequence 5'-CCGCCATNTT-3'; some genes have been shown to contain a longer binding motif allowing enhanced binding; the initial CG dinucleotide can be	anterior/posterior pattern specification, camera-type eye morphogenesis, cell differentiation, cellular response to DNA damage stimulus, cellular response to interleukin-1, cellular response to UV, chromosome organization, double-strand break repair via homologous recombination, negative regulation of gene expression, negative regulation of transcription from RNA polymerase II promoter, positive regulation of gene expression, positive

	<p>methylated greatly reducing the binding affinity. The effect on transcription regulation is depending upon the context in which it binds and diverse mechanisms of action include direct activation or repression, indirect activation or repression via cofactor recruitment, or activation or repression by disruption of binding sites or conformational DNA changes. Its activity is regulated by transcription factors and cytoplasmic proteins that have been shown to abrogate or completely inhibit YY1-mediated activation or repression. Binds to the upstream conserved region (UCR) (5'-CGCCATTTC-3') of Moloney murine leukemia virus (MuLV). Acts synergistically with the SMAD1 and SMAD4 in bone morphogenetic protein (BMP)-mediated cardiac-specific gene expression. Binds to SMAD binding elements (SBEs) (5'-GTCT/AGAC-3') within BMP response element (BMPRE) of cardiac activating regions. Proposed to recruit the PRC2/EED-EZH2 complex to target genes that are transcriptionally repressed. Involved in DNA repair. In vitro, binds to DNA recombination intermediate structures (Holliday junctions). Involved in spermatogenesis and may play a role in meiotic DNA double-strand break repair. Proposed core component of the chromatin remodeling INO80 complex which is involved in transcriptional regulation, DNA replication and probably DNA repair; proposed to target the INO80 complex to YY1-responsive elements.</p>	<p>regulation of transcription from RNA polymerase II promoter, regulation of transcription, DNA-templated, response to prostaglandin F, response to UV-C, RNA localization, spermatogenesis, transcription, DNA-templated</p>
Zeb1	<p>Acts as a transcriptional repressor. Binds to E-box sequences in the immunoglobulin heavy chain enhancer as well as in the regulatory regions of many other tissue-specific genes. Represses E-cadherin promoter and induces an epithelial-mesenchymal transition (EMT) by recruiting SMARCA4/BRG1. Represses BCL6 transcription in the presence of the corepressor CTBP1 (By similarity). Positively regulates neuronal differentiation. Represses RCOR1 transcription activation during neurogenesis. Represses transcription by binding to the E box (5'-CANNTG-3'). Promotes tumorigenicity by repressing stemness-inhibiting microRNAs.</p>	<p>animal organ development, cartilage development, cellular response to amino acid stimulus, central nervous system development, cochlea morphogenesis, embryonic camera-type eye morphogenesis, embryonic morphogenesis, embryonic skeletal system morphogenesis, negative regulation of cell proliferation, negative regulation of epithelial cell differentiation, negative regulation of transcription, DNA-templated, negative regulation of transcription from RNA polymerase II promoter, pattern specification process, positive regulation of neuron differentiation, positive regulation of transcription from RNA polymerase II promoter, regulation of mesenchymal cell proliferation, regulation of smooth muscle cell differentiation, regulation of T cell differentiation in thymus, regulation of transforming growth factor beta receptor signaling pathway, semicircular canal morphogenesis, transcription, DNA-templated</p>
Zfp105		regulation of transcription, DNA-templated
Zfx		fertilization, germ cell development, homeostasis of number of cells, multicellular organism growth, oocyte development, ovarian follicle development, parental behavior, post-embryonic development, regulation of transcription, DNA-templated, spermatogenesis, transcription, DNA-templated
Znf354c	<p>May function as a transcription repressor. Suppresses osteogenic effects of RUNX2. Binds to 5'-CCACA-3' core sequence. May be involved in osteoblastic differentiation</p>	regulation of transcription, DNA-templated, transcription, DNA-templated

Danksagung

Zum Gelingen meiner Dissertation hat eine Vielzahl von Menschen beigetragen, bei denen ich mich bedanken möchte.

Zuerst möchte ich meinem Doktorvater Herr Prof. Dr. Gerald Rimbach danken, dass er mir ermöglicht hat, meine Dissertation zu diesem spannenden Thema selbstständig und eigenverantwortlich bei ihm am Institut zu schreiben. In den regelmäßigen Gesprächen hat er mir immer wieder neue Anregungen gegeben und mich dazu gebracht, meine Fragestellung immer wieder aus anderen Perspektiven zu betrachten. Vielen Dank für die angenehme Zusammenarbeit.

Ganz besonderen Dank gilt auch Frau Dr. Patricia Hübbe, die mir während der gesamten Zeit zur Seite gestanden hat, bei Fragen immer ein offenes Ohr und zudem immer wieder neue Anregungen für mich gehabt hat, bspw. an welcher Stelle ich nochmal weiter ins Detail gehen könnte bzw. welche Analysemethoden ebenfalls sinnvoll wären. Gleichzeitig ist sie mir immer eine große Motivation gewesen, vor allem auch in Momenten, in denen ich einmal nicht weiter gekommen bin.

Weiterhin möchte ich Herrn Dr. Mario Hasler und Herrn Dr. Dieter-Christian Gottschling danken, die mir bei statistischen Fragen bzgl. Excel und R weitergeholfen haben.

Ein ganz besonderer Dank gilt weiterhin meinem Mann Mike, der mir während der gesamten Zeit immer wieder den Rücken freigehalten hat, damit ich neben meiner Arbeit die Zeit finden konnte, um an meiner Dissertation zu arbeiten. Ebenfalls möchte ich ihm, meinen Eltern und meinen Geschwistern dafür danken, dass sie mich immer wieder motiviert haben, wenn es mal schwierig wurde und gleichzeitig Verständnis dafür gehabt haben, wenn ich über Tage nicht richtig ansprechbar gewesen bin, weil ich in meine Analysen vertieft war. Gleichermaßen gilt für meine Freunde, die immer wieder die Geduld aufgebracht haben, mit mir über meine Dissertation zu sprechen und die mich immer wieder motiviert haben und Verständnis dafür gehabt haben, wenn meine zeitlichen Kapazitäten sehr knapp gewesen sind.

Danken möchte ich ebenfalls meinem Arbeitgeber Herrn Jochen Timmermann, der es mir strukturell ermöglicht hat, nebenberuflich an meiner Dissertation zu schreiben. Gleichzeitig möchte ich auch all meinen Kollegen danken, die immer wieder ein offenes Ohr für mich gehabt haben. Meine parallele praktische Tätigkeit ist eine wertvolle Bereicherung während meiner Dissertationszeit gewesen, zumal der regelmäßige Abstand zu meinen Analysen nicht selten einen wertvollen Beitrag geleistet hat, um anschließend mit neuen Ideen weiterarbeiten zu können.

Curriculum Vitae

Stefanie Rieper née Busch
born june 17th 1988
in Otterndorf, Germany

Education

07/2016	PhD oral examination
11/2012 - 07/2016	PhD, Division of Food Science Christian-Albrechts-University Kiel, Germany
10/2010 - 06/2012	Master of Science, Nutritional Science and Household Economics Christian-Albrechts-University Kiel, Germany Master thesis: "Einfluss von Bor auf die differentielle endotheliale Genexpression"
10/2007 - 08/2010	Bachelor of Science, Nutritional, Food and Consumer Sciences University of Applied Sciences Fulda, Germany Bachelor thesis: "Atopieprävention während der Schwangerschaft bei positiver Familienanamnese – aktuelle Studienergebnisse zur Relevanz einer mütterlichen Diät"

Professional Experience

Since 05/2012	Dietician Medical care unit for physiological and psychological health Timmermann and partner, Cuxhaven
---------------	--