



IL-22 Paucity in APECED Is Associated With Mucosal and Microbial Alterations in Oral Cavity

Epp Kaleviste^{1*}, Malte Rühlemann², Jaanika Kärner¹, Liis Haljasmägi¹, Liina Tserel¹, Elin Org³, Katarina Trebušak Podkrajšek⁴, Tadej Battelino⁵, Corinna Bang², Andre Franke², Pärt Peterson¹ and Kai Kisand^{1*}

¹ Department of Biomedicine, Institute of Biomedicine and Translational Medicine, University of Tartu, Tartu, Estonia, ² Institute of Clinical Molecular Biology, University of Kiel, Kiel, Germany, ³ Estonian Genome Centre, University of Tartu, Tartu, Estonia, ⁴ University Medical Centre Ljubljana, University Children's Hospital, Ljubljana, Slovenia, ⁵ Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia

OPEN ACCESS

Edited by:

Andrew R. Gennery,
Newcastle University, United Kingdom

Reviewed by:

Michail Lionakis,
National Institutes of Health (NIH),
United States
Desa Ljilic,
Newcastle University, United Kingdom

*Correspondence:

Epp Kaleviste
epp.kaleviste@gmail.com
Kai Kisand
kai.kisand@ut.ee

Specialty section:

This article was submitted to
Primary Immunodeficiencies,
a section of the journal
Frontiers in Immunology

Received: 17 January 2020

Accepted: 14 April 2020

Published: 08 May 2020

Citation:

Kaleviste E, Rühlemann M, Kärner J, Haljasmägi L, Tserel L, Org E, Trebušak Podkrajšek K, Battelino T, Bang C, Franke A, Peterson P and Kisand K (2020) IL-22 Paucity in APECED Is Associated With Mucosal and Microbial Alterations in Oral Cavity. *Front. Immunol.* 11:838. doi: 10.3389/fimmu.2020.00838

Autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED) is caused by recessive mutations in the *AIRE* gene. The hallmark of the disease is the production of highly neutralizing autoantibodies against type I interferons and IL-22. Considering the importance of IL-22 in maintaining mucosal barrier integrity and shaping its microbial community, we sought to study potential changes in the oral cavity in this model of human IL-22 paucity. We found that besides known Th22 cell deficiency, APECED patients have significantly fewer circulating MAIT cells with potential IL-22 secreting capacity. Saliva samples from APECED patients revealed local inflammation, the presence of autoantibodies against IFN- α and IL-22, and alterations in the oral microbiota. Moreover, gene expression data of buccal biopsy samples suggested impaired antimicrobial response and cell proliferation, both of which are processes regulated by IL-22. Our data complement the knowledge gained from mouse models and support the concept of IL-22 being a critical homeostatic cytokine in human mucosal sites.

Keywords: APECED, AIRE, IL-22, oral mucosa, microbiota

INTRODUCTION

Autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy syndrome is a rare recessive monogenic disorder (1). The responsible gene, *AIRE*, is expressed in the thymic medullary epithelial cells to support the negative selection of thymocytes that are potentially reactive to peripheral tissue-specific antigens (2–5). Nevertheless, among the autoimmune manifestations, the impaired clearance of *Candida* infection from the mucosal surfaces stands out as the most common and, usually, the first manifestation of the disease (6, 7). It is well established that CMC correlates with circulating autoantibodies against Th17 related cytokines IL-22 and IL-17F (8–10), and that the secretion of the respective cytokines is severely impaired in the circulating and skin-residing

Abbreviations: AHR, aryl hydrocarbon receptor; AIRE, autoimmune regulator; AMP, antimicrobial peptide; APECED, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy; ASV, amplicon sequence variant; CMC, chronic mucocutaneous candidiasis; IPA, ingenuity pathway analysis; LIPS, luciferase based immunoprecipitation system; MAIT, mucosal associated invariant T; NKT, natural killer T; TFIIH, transcription factor IIH.

CD4+ T cells (9, 11). The potential pathogenicity of the IL-22-neutralizing autoantibodies, isolated from APECED patients, has been confirmed in a mouse model of oropharyngeal candidiasis, where the antibody treatment caused delayed clearance of the yeast from the oral cavity (12).

IL-22 is essential for mucosal barrier function. It can protect from intestinal injury by supporting epithelial cell proliferation and wound healing, enhancing tight junctions, upregulating antimicrobial peptide, and mucus production (13–16). Moreover, IL-22 was recently shown to protect intestinal stem cells against genotoxic stress and thus against colon cancer (17). IL-22 is also capable of shaping gut microbiota (18). In contrast, the excessive production of IL-22 is associated with tissue inflammation in several immune-mediated inflammatory diseases, such as psoriasis, celiac disease, and rheumatoid arthritis (19–23).

While the target cells of IL-22 action are mostly epithelial cells, the best producers of IL-22 are various lymphoid cells: Th17 (24), Th22 (25), type 3 innate lymphoid cells (26), and several unconventional T cells, such as $\gamma\delta$ T (27), MAIT (28, 29), NKT (30), and invariant NKT cells (31). Unconventional T cells are crucial for the protection and homeostasis of the epithelial surfaces due to their immediate response to harmful agents. However, they are less studied in APECED patients than are conventional T cells.

Most of the knowledge about the functions of IL-22 have been derived from mouse and *in vitro* experiments. We reasoned that APECED can be viewed as a model disease that enables to study the consequences of IL-22 insufficiency in human oral mucosa. The shortage of IL-22 in APECED is thought to be associated with CMC, but considering the importance of this cytokine for epithelial cell homeostasis in the digestive tract, we hypothesized that it should lead to various other important consequences for epithelial homeostasis.

MATERIALS AND METHODS

Subjects

We studied 13 patients with APECED (9 males, 4 females) from Slovenia and Estonia and 16 control subjects, who were age and gender adjusted for the study, and recruited at the same time with the patients. Use of human material was approved by local ethics committees (Slovenia: National Medical Ethics Committee number 22/09/09 and 28/02/13; Estonia: Research Ethics Committee of the University of Tartu, 235/M-23). Informed consent was obtained from all participants or parents of participating children. Patient details are given in **Supplementary Table 1**.

Material

Peripheral blood was drawn into heparinized vacutainers, separated into plasma and peripheral blood mononuclear cells and stored at -20°C or liquid nitrogen, respectively, until usage. The saliva samples were provided using the passive-drool method, in which study participants allowed saliva to pool in the mouth and then drool it into a tube. Donors

did not eat or drink for 30 min before sample collection. Samples were stored at -80°C . Buccal biopsies were taken with surgical scalpel under aseptic conditions after local anesthesia. A core from the buccal mucosa of either the left or right cheek was obtained, rinsed in sterile PBS and snap frozen. None of the studied patients or controls received immunosuppressive treatment, reported sicca symptoms nor had difficulties in saliva collection.

Flow Cytometry

Surface marker expression on PBMCs was assessed by flow cytometry in 8 patients and 8 age matched control subjects. Cells were stained in flow cytometry buffer (PBS (pH 7.2), 2 mM EDTA, 0.5% BSA) for 20 min at room temperature in dark with antibodies listed in **Supplementary Table 2**. After staining, cells were analyzed using LSRFortessa flow cytometer (BD Biosciences) and FCS Express 5 Flow (*De Novo* Software). The optical detector configuration can be found in **Supplementary Table 3**. The gating strategy is depicted in **Supplementary Figure 1**.

Autoantibodies From Plasma and Saliva With LIPS

Luciferase based immunoprecipitation system assay was conducted on saliva samples (10 APECED patients and 10 controls) and on plasma samples (13 APECED patients and 7 controls). LIPS assay was performed as previously described (32). Briefly, IFN- α , IL-17A, IL-17F, and IL-22 coding sequences were cloned into modified pPK-CMV-F4 fusion vector (PromoCell GmbH, Germany) where Firefly luciferase was substituted with NanoLuc luciferase (Promega, United States). Cloned constructs were transfected into HEK293 cells and secreted luciferase fusion proteins were collected with culture medium after 48 h. IgG from plasma and saliva samples was captured onto Protein G Agarose beads (Exalpha Biologicals, United States) at room temperature for 1 h in 96-well microfilter plate (Merck Millipore, Germany). Next, 1×10^6 luminescence units (LU) of antigen was added per well. After 1 h of incubation the unbound antigen was washed away with vacuum system and Nano-Glo[®] Luciferase Assay Reagent (Promega, United States) was added. Luminescence intensity was measured by VICTOR X Multilabel Plate Reader (PerkinElmer Life Sciences, United States). The results were expressed as luminescence units (LU) representing the fold over the mean of the healthy control samples.

Cytokines From Saliva and Plasma

Cytokine quantification was conducted on saliva samples (10 Slovenian patients and 8 controls, with exception for TNF- α , IL-2, IL-1 β , IL-5, IL-6, IL-10, GM-CSF, and IFN- γ which were measured with 6 patient samples and 5–6 control samples) and on plasma samples (8 patients and 9 controls, with the exception of TNF- α , IL-2, IL-1 β , IL-5, IL-6, IL-10, GM-CSF, and IFN- γ , which were measured from 7 patients and 8 controls). Cytokine levels in saliva samples and EDTA-treated plasma samples were measured by the xMAP Technology on Luminex 200 (Luminex Corp.,

Austin, TX, United States) with Human Magnetic Luminescence Performance Assay Kits (R&D Systems, Minneapolis, MN, United States) (**Supplementary Table 4**). The cytokine levels were analyzed in accordance with the manufacturer's protocol. The cytokines studied are listed in **Supplementary Table 5** including their lower limits of detection. The values of IL-1 α in plasma samples and IL-2, IL-5, IL-17A, IL-17F, and IL-22 in both sample types fell below the detection limits in the majority of tested samples and therefore were excluded from further analysis.

Gene Expression From Buccal Biopsy

Buccal biopsy samples were derived from 4 patients and 4 age-matched healthy controls. However, after RNA isolation, 2 control samples did not pass the RNA quality control, and therefore, the gene expression analysis was conducted on 4 APECED patient and 2 control buccal biopsy samples. The biopsy specimens were homogenized in TRIzol (Invitrogen) and RNA in the aqueous phase was extracted with the miRNeasy Mini Kit combined with the RNase-free DNase I treatment (both from Qiagen). The concentration and quality of the RNA were assessed with an Agilent RNA 6000 Nano Kit on Agilent 2100 Bioanalyzer (Agilent Technologies, CA, United States). Next, cRNA was prepared from 300 ng of total RNA using an Illumina TotalPrep RNA Amplification Kit (Ambion Inc., TX, United States) according to the manufacturer's protocol and a genome-wide gene expression analysis was carried out using HumanHT-12 v4 Expression BeadChip (Illumina Inc., CA, United States) and the signals were scanned using a Beadscan (Illumina Inc.). Array data were analyzed using GenomeStudio software (Illumina, San Diego, CA, United States). Rank invariant normalization was applied. Genes with the FDR adjusted p -value < 0.05 and a fold change > 1.5 were considered to be differentially expressed. A positive diffScore represents upregulation, while a negative diffScore represents downregulation. The p -value for an observed expression difference between two analyzed groups of samples is < 0.05 for genes with diffScores of lower than -13 and higher than $+13$ (p -value < 0.01 is $-20 > \text{diffScore} > 20$; p -value < 0.001 is $-33 > \text{diffScore} > 33$). The data with negative average signal value was excluded from the list. Annotation analysis was implemented with gprofiler software (33) and IPA software (QIAGEN Inc)¹. Analysis data set included the upregulated and downregulated gene list with expression fold change. The core analysis was carried out on reference set Ingenuity[®] Knowledge Base (Genes only). The settings were by default including direct and indirect relationships between genes, 35 genes per network, 25 networks per analysis. Diseases and functions analysis was conducted, which predicted cellular processes and biological functions, based on gene expression. Z-score provides predictions about upstream or downstream processes. Network analysis generated the scores [p -score = $-\log_{10}$ (p -value)] according to the fit of the set of supplied genes and a list of biological functions. Focus Genes were those from the uploaded list that pass filters and are eligible

for generating networks. Only genes with a fold change of 1.5 and p -value ≤ 0.05 were considered. Upstream regulator analysis in IPA was used to predict the upstream transcriptional regulators from the dataset based on the literature and compiled in the Ingenuity Knowledge Base. Upstream regulator analysis calculated the causal networks. The causal networks scores were calculated using causal paths only. Two p -values were calculated: Fishers Exact Test of whether there was a greater than expected proportion of downstream data set genes than expected by chance. Network bias corrected p -value was a measure of how often a more significant result was seen in 10K iterations of selecting random data sets of genes with similar relationship number.

Candida Detection From Saliva

DNA from saliva samples and cultured laboratory strain SC5314 (ATCC, VA, United States) of *Candida albicans* was extracted using the DNeasy PowerSoil Kit (QIAGEN) according to manufacturer's protocol. Samples were homogenized with Precellys 24 Homogenizer (Bertin Instruments) with program 6500 rpm, 2 min' homogenization and 1-min break. Real time quantitative PCR was performed using Applied Biosystems[®] ViiA[™] 7 Real-Time PCR System with 384-Well Block (Life Technologies) and Maxima SYBR Green/ROX qPCR Master Mix (Fermentas). A primer pair of highly conserved *Candida* rDNA-coding region 28S was used (forward CGGCGAGTGAAGCGGCTAA, reverse ATTCCCAAACAACCTCGACTC) (34). The concentration of *Candida* cells in patient samples was calculated based on the calibration curve constructed according to the Ct values of serially diluted *C. albicans* probes (starting from 2E7 CFU/ml). The normal range of *Candida* concentration was calculated based on healthy control values: mean + 3 standard deviations.

Sequencing, Processing and Statistical Analysis of Bacterial 16S rDNA Sequences

DNA from saliva samples was manually extracted using the MoLYsis[™] Complete 5 Kit according to manufacturer's protocol with 20 μ l elution of DNA in water. The bacterial 16S rDNA variable region V1-V2 amplicon libraries were prepared using the 27F-338R primer combination in a dual indexing approach. Sequencing was performed on an Illumina MiSeq device using 2 \times 300 cycles and MiSeq Reagent Kit v3 (35). The Dataset included 6 Slovenian patient and 6 control samples. Raw-Data were processed using the DADA2 workflow² (v1.10) and subsequently analyzed in R using the vegan and phyloseq packages. Taxonomic annotation was performed using a Bayesian classifier and RDP training set 16. For each sample, 20k sequences were randomly sampled for a normalized read count. Analysis was performed on ASV, species and genus level. For differentially abundant taxonomic group all tests were performed using Wilcoxon

¹<https://www.qiagenbioinformatics.com/products/ingenuity-pathway-analysis/>

²<https://benjjneb.github.io/dada2/>




FIGURE 1 | Comparison of circulating lymphocyte subpopulations. Surface marker expression on PBMCs was assessed by flow cytometry in 8 APECED patients and 8 age-matched controls. The percentage of Vδ1+ and Vδ2+ γδ T cells, iNKT and MAIT cells from total T cell pool was compared between patient and control sample groups. In the violin plot the width of distribution of points is proportionate to the number of points at the value of the sample. Black discontinuous line represents median and colored discontinuous line represents quartiles. Statistical significance was assessed with *t*-test using R statistical software ($p < 0.05$). The outliers for different cell types did not overlap, with the exception of the APECED patient who had the highest proportion of both γδ T cell subtypes.

rank sum test using only taxa with median abundance > 100 sequence counts (0.5% rel. abundance) and prevalence > 0.3 (present in at least 4 samples). Alpha diversity (within sample diversity) was analyzed with Wilcoxon rank sum tests. Diversity indices used were as follows: Observed – Observed taxa with abundance > 0, Shannon – Shannon Diversity Index, Chao1 – Chao1 Estimator to estimate observed and unobserved species count. Differences in beta diversity (Whole Community Composition/inter-sample diversity), were assessed using Bray-Curtis similarity and permutational multivariate analysis of variance as implemented in the *adonis()* function of the *vegan* package for R. The sequences have been deposited in the NCBI Sequence Read Archive Database³ under accession number PRJNA601650.

Statistical Analysis

Statistical analysis was performed using the R statistical software⁴ and GraphPad Software (San Diego, CA, United States).

RESULTS

APECED Patients Have Decreased MAIT Cell Proportions in Their Circulation

With regard to the severe deficiency of Th22 cells in the blood circulation and skin of the APECED patients (9, 11), we enquired if the proportions of unconventional T cells, that are known for their IL-22 production capability, are also impaired in the patients. We used flow cytometry to enumerate these cells (Figure 1) (gating strategy displayed in

Supplementary Figure 1). Vδ1+ and Vδ2+ γδ T cell proportions were not significantly altered, in accordance with previous studies (36), neither did the iNKT cell numbers differ between patient and control samples. Nevertheless, the percentages of MAIT cells (Vα7.2 TCR+ and CD161+) among T cells were lower in patients than in healthy controls ($p < 0.05$).


Saliva From APECED Patients Contain Cytokine Autoantibodies and Reflect Inflammation

As IL-22 and MAIT cells are both important players in mucosal homeostasis and protection, we analyzed saliva samples from patients and controls to find signs of perturbations on oral mucosa. First, we tested the saliva and corresponding plasma samples for the presence of cytokine autoantibodies (Figure 2). Importantly, anti-IFN-α and anti-IL-22 autoantibodies were present in saliva from all the 10 APECED patients. Salivary anti-IL17A and anti-IL17F autoantibodies were detectable at very low levels in a fraction of the patients who tested positive for corresponding circulating autoantibodies. In addition, we detected LCN1 autoantibodies from the plasma of 3 of 13 APECED patients. We also tested for different anti-S100A subtypes (S100A7A, S100A7, S100A8, S100A9) in the plasma, but there was no difference between patient and control groups (Supplementary Figure 2). Collectively, these findings suggest that the local presence of IFN-α and IL-22 autoantibodies can aggravate the insufficiency of these cytokines on mucosal surfaces.

Next, we compared the salivary concentration of various cytokines in APECED patients with those in healthy controls (Supplementary Table 5). IL-22, IL-17A, and IL-17F levels remained undetectable in majority of the samples. However, proinflammatory cytokines TNF-α, GM-CSF, and

³<https://www.ncbi.nlm.nih.gov/sra>

⁴<https://www.r-project.org/>




IFN- γ were all significantly increased compared to control group (Figure 3A). This increase was reflective of the local inflammation as the difference was not recapitulated in the plasma samples (Figure 3B). From tested plasma cytokines, only CXCL10 was significantly increased in patients compared with that in controls, confirming an earlier report (37).

Furthermore, correlation analysis was conducted to compare the *Candida* concentrations (CFU/ml) with significantly increased cytokine levels and autoantibody levels from patient saliva samples (Supplementary Figure 3). There was no correlation between CFU values and proinflammatory cytokine concentrations. The only significant correlation was between salivary anti-IL-17A levels and *Candida* CFU, but this was caused by the one outlier – the patient with the highest

Candida CFU value had the highest IL-17A antibody level (Supplementary Figure 4).

Buccal Biopsy Transcriptomes Reveal Alterations in APECED Oral Mucosa

The gene expression analysis was conducted on 4 APECED patient and 2 control buccal biopsy samples. Differentially regulated genes were identified based on gene expression diffScore (Supplementary Table 6). Seventeen significantly up-regulated and 91 significantly down-regulated genes were identified. Among the upregulated genes, several transcripts were associated with tumorigenesis (*SNHG7*, *SHIP2*, *FANCG*, and *DDR2*). In contrast, several AMP genes (*DEFB103B*, *DEFB103A*, and *S100A12*) and genes important for epithelial barrier



function (*SPRR2C*, *SPRR2B*, and *TGM5*) were downregulated. The significantly downregulated genes were further analyzed with gProfiler database. Mitotic cell cycle was identified as an associated biological process. In addition, the gene lists together with fold changes were analyzed by IPA. We carried out a diseases and functions analysis, which predicted the biological processes (cellular processes, biological functions, etc.) affected by the expression of a specific gene. The main terms that emerged were connected with cell cycle and mitosis (Supplementary Table 7), the cell cycle process was predicted to be inhibited in patients (*p*-value 3.5E-08, activation *z*-score -2.256). Next, we carried out a network analysis, which predicts gene interaction map (Supplementary Table 8). We focused on the network with the highest score, which consisted of 23 focus molecules and 12 interconnecting molecules (“non-focused” proteins that were not present in our list). The main network included the following biological terms: antimicrobial response, cellular function and maintenance, and inflammatory response (Figure 4 and Supplementary Figure 5).

Finally, we conducted an upstream regulator analysis, which predicts the activity state of regulators by correlating literature

reported effects with observed gene expression. Participating regulators are molecules through which the upstream regulator molecule controls the expression of target molecules in dataset. The top two master regulators, identified in the dataset, were *TFIIH* and *AHR*, based on the highest *p*-value of overlap (Supplementary Table 9 and Supplementary Figure 6). Transcription factor IIH (TFIIH) is an important protein complex with many biological roles, ranging from DNA repair to transcription to cell cycle regulation (38). AHR is highly expressed by Th17 cells, and activation of AHR results in expansion of Th17 cells and enhanced production of Th17 cytokines, particularly IL-22 (15, 39, 40).

Taken together, the gene expression array results hint that IL-22-controlled pathways may be downregulated in the buccal biopsy samples of APECED patients.

Salivary Microbiota Analysis Is Consistent With Dysbiosis in APECED Oral Cavity

Salivary microbiota was analyzed in 6 APECED patients and 6 age-matched healthy controls. The microbial communities were significantly different in APECED patients in comparison to healthy controls on the whole community (beta-diversity) level (Figure 5). All differences were significant in permutational ANOVA (ASV: $R^2 = 14.6\%$, $p = 0.017$, species: $R^2 = 20.2\%$, $p = 0.036$, genus: $R^2 = 23.5\%$, $p = 0.021$). Interestingly, the 2 patients with elevated *Candida* levels in their saliva samples (Supplementary Table 1 and Supplementary Figure 3), clustered together at the genus and species level (Figure 5).


Next, we analyzed the taxonomic groups. The most abundant genus in both, patients and controls, was *Streptococcus* (Figure 6). Significantly higher abundance of *Fusobacterium* ($p < 0.05$) was revealed in APECED patients, and a relatively rare genus, *Lachnoanaerobaculum*, was more abundant in healthy individuals. At the species and ASV level, we noted significantly lower abundance of *Streptococcus salivarius* ($p < 0.05$) in APECED patient samples (Figure 7 and Supplementary Figure 7).

For alpha-diversity analysis, which estimates bacterial diversity within a sample (Figure 8), three different indices were used to estimate counts of observed and unobserved species: observed taxa with abundance over zero, Shannon Diversity Index, and Chao 1 Estimator. Interestingly, significantly increased Shannon diversity was found at the species ($p < 0.05$) and ASV ($p < 0.01$) levels in patient samples. Other alpha-diversity indices showed similar tendencies, but the differences were not significant in this small dataset.

Taken together, the results of the microbial 16S sequencing analysis indicated alterations in oral microbiota of APECED patients and increased alpha-diversity.

DISCUSSION


Since its discovery almost 20 years ago, IL-22 has gained considerable attention as an essential homeostatic cytokine on epithelial surfaces. IL-22, belonging to Th17 family of cytokines,



has similar and even synergistic effects with IL-17A, but it also exerts a plethora of unique effects to secure the barrier function of epithelium (13–16). In this study, we focused on a distinctive feature of APECED patients, namely, highly neutralizing autoantibodies against IL-22 and IFN- α . Previous studies have substantiated that due to neutralization of the low physiologic levels of IFN- α , blood cells of APECED patients have reduced levels of IFN-induced gene expression compared with that of healthy controls (37). Moreover, CMC in APECED patients is associated with the presence of circulating anti-IL-22 and IL-17F (11, 32, 41) and with the incapability of circulating and skin CD4+ T cells to secrete IL-22 after their stimulation *in vitro*. In addition, impaired mucosal barrier function has been suspected in APECED patients due to increased levels of

antibodies to commensals (42). In this study, we focused on the events occurring in the human oral mucosa.

The cause for the almost complete absence of IL-22 production by circulating and skin CD4+ T cells in APECED patients is still not known. Furthermore, the role of other lymphoid cells in Th17-related cytokine secreting capacity is not yet clear in this disease. However, there is a previous report suggesting that IL-17A secreting $\gamma\delta$ T cells are impaired in APECED (43). According to current study, the percentages of V δ 1+ and V δ 2+ $\gamma\delta$ T cells did not differ between patients and controls. The association of iNKT cells with AIRE-deficiency has remained controversial (10, 44, 45). Our study revealed no significant difference between patient and control samples for this cell type. However, we found significantly decreased




proportions of circulating MAIT cells in APECED patients. MAIT cells are reportedly present in oral mucosa and are capable of secreting Th17 cytokines (46). Considering this, it is possible that the paucity of MAIT cells contributes to the shortage of IL-22 in mucosal surfaces. The deficiency of MAIT cells in patients can be related to the presence of highly neutralizing autoantibodies against type I IFNs in APECED because MAIT cells are activated by type I IFNs, or by aberrations in microbiota that produce metabolites necessary for MAIT cell development in the thymus (47).

Although the main producers of IL-22 are distinct cell types from the lymphoid lineage (48), which are deficient in APECED patients, dendritic cells, and neutrophils can also contribute to the circulating or local concentration of IL-22 (49, 50). The remaining IL-22 from various sources can be neutralized by IL-22 autoantibodies. Interestingly, we demonstrated autoantibodies against IL-22 and IFN- α also in patients' saliva samples, levels of which varied a lot between patients. In theory, these local autoantibodies can further aggravate IL-22 shortage although we do not have formal proof for this. One could also hypothesize that local neutralization of IFN- α can further impair the recruitment of cells that could secrete IL-22, as recently reported (51).

As one of the key functions of IL-22 is to induce AMP secretion from epithelial cells in concert with IL-17A, it is possible that the secretion of these natural antibiotics is impaired in APECED patients. However, previous work has not found decreased beta and alpha defensin concentrations in APECED saliva samples (52). Our study confirmed abundant, and not impaired, LCN2 levels in the saliva of APECED patients. Gene expression analysis of buccal biopsy samples revealed equally high S100A8 and S100A9 expression in patients and controls, while some other AMPs (DEFB103B, DEFB103A, and S100A12) were significantly downregulated in the mucosa of


APECED patients. These results suggest specific but not universal impairment of AMP production in APECED oral cavity. Interestingly, autoantibodies against antimicrobial peptides, DEFA5 (53), LPO (54), BPIFA1 (54), BPIFA2 (54), and LCN1 (55), have been described previously. In addition, we detected LCN1 autoantibodies in the plasma of three of 13 APECED patients included in this study, but did not detect autoantibodies against the S100A family of antimicrobial peptides.

Considering the importance of the tight regulation of IL-22 levels to enable the symbiosis of the host with commensals (56), we hypothesized that IL-22 paucity could impact microbial community in the oral cavity. Indeed, the salivary microbiome of patients was significantly different from healthy controls (Figure 7) and showed higher diversity. The dysbiosis in APECED patients' oral cavity has been described in an earlier investigation with some overlaps in differentially abundant taxonomic units between the two studies, but also with discrepancies that can arise from various technical, analytical and biological differences between the two studies (57). Alterations in the microbiota could also be caused by medications, as well as Sjögren's-like syndrome, which is especially frequent among APECED patients in the United States (10). However, we think that neither factor had impact on our analysis as none of the study subjects received immunosuppressive drugs, and only one patient was on antifungals, all other medications were only hormone replacements. Moreover, objective sicca symptoms are rare in European APECED patients and were reportedly present only in patients over 33 years of age (52). Our study included relatively young patients (17.7 ± 10.9 years) and none of them reported dry mouth. *Candida* infection itself is also a potential modifier of microbiota. According to the beta-diversity analysis (Figure 5) the two patients with elevated *Candida* concentration




clustered together at the genus and species level but did not differ from the other patients in any other studied parameter. CMC is also a possible inducer of proinflammatory cytokines that were elevated in the patients' saliva that in turn can influence the changes in the microbial communities. Although we did not see any significant correlation between *Candida* CFU values and cytokine concentrations, we cannot

rule it out by our small study group. The cause of the salivary proinflammatory cytokine increase remains unknown. The oral cavity was regularly examined by the attended pediatric endocrinologist and no clinical signs of gingivitis and periodontitis were appreciated. The autoimmune attack toward mucosal antigens or preclinical inflammation in the salivary glands remains an option.



In addition to shaping the microbiome composition, IL-22 is important for epithelial cell proliferation, enhancing tight junctions and protecting cells from genotoxic stress (17). Indeed, our gene expression data from buccal biopsy samples suggested differences in the upstream and downstream molecular events of the IL-22 pathway. Notably, the upstream regulator analysis identified AHR, the transcription factor

necessary for IL-22 production, while network analysis revealed impaired antimicrobial response, cellular function and maintenance, and inflammatory response. The harmed epithelial barrier function can be suspected also from the reduced expression of several genes related with mitotic cell cycle in patients that can lead to defective wound healing on epithelial surfaces that can become especially



important during inflammatory processes. Importantly, chronic inflammation due to recalcitrant *Candida* infection has been regarded as a susceptibility factor for developing oral and esophageal squamous cell carcinoma in APECED patients (58). In line with this, the patient saliva samples revealed significantly increased levels of proinflammatory cytokines. As IL-22 was recently shown to protect intestinal stem cells against genotoxic stress and thus against colon cancer (17), we suggest that the susceptibility for oral and esophageal squamous cell carcinoma may result from the combination of persistent inflammation with the lack of protective function of IL-22.

Our study has several limitations including a small sample size and imprecise taxonomic annotation of the 16S amplicons due to the clade-dependent differences in resolution exhibited by the amplicon. Moreover, the usual weakness of the human studies is their descriptive nature and purely indirect reasoning to suggest some causalities.

CONCLUSION

According to our results, APECED patients are characterized by alterations in their oral mucosa and increased diversity of their salivary microbiota, both of which

can be in theory influenced by IL-22 shortage, possibly aggravated by local action of autoantibodies and paucity of circulating MAIT cells.

DATA AVAILABILITY STATEMENT

The datasets generated for this study can be found in the NCBI Sequence Read Archive Database, PRJNA601650.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by National Medical Ethics Committee number 22/09/09 and 28/02/13, Slovenia; Research Ethics Committee of the University of Tartu, 235/M-23, Estonia. Written informed consent to participate in this study was provided by the participant or/and the participants' legal guardian/next of kin.

AUTHOR CONTRIBUTIONS

EK and JK carried out flow cytometry experiments and analyzed the respective data. LH performed LIPS. MR, CB, AF,

and EO were responsible for the microbiota analysis. KK, LT, and EK analyzed the gene expression array data. KT and TB recruited the study subjects, collected the samples, and clinical data. KK and PP supervised the research. EK and KK wrote the manuscript with the support of all the other coauthors.

FUNDING

We thank financial support from the European Union through the European Regional Development Fund (Project No 2014-2020.4.01.15-0012) and the Centre of Excellence in Genomics (EXCEGEN framework), the Slovenian Research Agency (research core funding P3-0343), the University of Tartu Center

of Translational Genomics (SPIGVARENG), and the Estonian Research Council grants PUT1367 and PRG377.

ACKNOWLEDGMENTS

We express extreme gratitude to APECED patients who provided samples, attending physicians, and care-takers of the patients.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2020.00838/full#supplementary-material>

REFERENCES

- Kisand K, Peterson P. Autoimmune polyendocrinopathy candidiasis ectodermal dystrophy. *J Clin Immunol.* (2015) 35:463–78. doi: 10.1007/s10875-015-0176-y
- Heino M, Peterson P, Kudoh J, Nagamine K, Lagerstedt A, Ovod V, et al. Autoimmune regulator is expressed in the cells regulating immune tolerance in thymus medulla. *Biochem Biophys Res Commun.* (1999) 257:821–5. doi: 10.1006/bbrc.1999.0308
- Poliani PL, Kisand K, Marrella V, Ravanini M, Notarangelo LD, Villa A, et al. Human peripheral lymphoid tissues contain autoimmune regulator-expressing dendritic cells. *Am J Pathol.* (2010) 176:1104–12. doi: 10.2353/ajpath.2010.090956
- Anderson MS, Venanzi ES, Klein L, Chen Z, Berzins SP, Turley SJ, et al. Projection of an immunological self shadow within the thymus by the aire protein. *Science.* (2002) 298:1395–401. doi: 10.1126/science.1075958
- Yang S, Fujikado N, Kolodin D, Benoist C, Mathis D. Regulatory T cells generated early in life play a distinct role in maintaining self-tolerance. *Science.* (2015) 348:589–94. doi: 10.1126/science.aaa7017
- Ahonen P, Myllärniemi S, Sipilä I, Perheentupa J. Clinical variation of autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy (APECED) in a series of 68 patients. *N Engl J Med.* (1990) 322:1829–36. doi: 10.1056/NEJM199006283222601
- Perheentupa J. Autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy. *J Clin Endocrinol Metab.* (2006) 91:2843–50. doi: 10.1210/jc.2005-2611
- Puel A, Dffinger R, Navidad A, Chrabieh M, Barcenas-Morales G, Picard C, et al. Autoantibodies against IL-17A, IL-17F, and IL-22 in patients with chronic mucocutaneous candidiasis and autoimmune polyendocrine syndrome type I. *J Exp Med.* (2010) 207:291–7. doi: 10.1084/jem.20091983
- Kisand K, Boe Wolff AS, Podkrajsek KT, Tserel L, Link M, Kisand KV, et al. Chronic mucocutaneous candidiasis in APECED or thymoma patients correlates with autoimmunity to Th17-associated cytokines. *J Exp Med.* (2010) 207:299–308. doi: 10.1084/jem.20091669
- Ferre EMN, Rose SR, Rosenzweig SD, Burbelo PD, Romito KR, Niemela JE, et al. Redefined clinical features and diagnostic criteria in autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy. *JCI Insight.* (2016) 1:e88782. doi: 10.1172/JCI.INSIGHT.88782
- Laakso SM, Kekäläinen E, Heikkilä N, Mannerström H, Kisand K, Peterson P, et al. *In vivo* analysis of helper T cell responses in patients with autoimmune polyendocrinopathy – candidiasis – ectodermal dystrophy provides evidence in support of an IL-22 defect. *Autoimmunity.* (2014) 47:556–62. doi: 10.3109/08916934.2014.929666
- Bichele R, Kärner J, Truusalu K, Smidt I, Mändar R, Conti HR, et al. IL-22 neutralizing autoantibodies impair fungal clearance in murine oropharyngeal candidiasis model. *Eur J Immunol.* (2018) 48:464–70. doi: 10.1002/eji.201747209
- Costa MM, Saraceni PR, Forn-Cuní G, Dios S, Romero A, Figueras A, et al. IL-22 is a key player in the regulation of inflammation in fish and involves innate immune cells and PI3K signaling. *Dev Comp Immunol.* (2013) 41:746–55. doi: 10.1016/j.dci.2013.08.021
- Eidenschek C, Rutz S, Liesenfeld O, Ouyang W. Role of IL-22 in microbial host defense. In Fillatreau S, O'Garra A editors. *Interleukin-10 in Health and Disease. Current Topics in Microbiology and Immunology*, (Vol. 380), Berlin: Springer. (2014). p. 213–36. doi: 10.1007/978-3-662-43492-5_10
- Monteleone I, Rizzo A, Sarra M, Sica G, Sileri P, Biancone L, et al. Aryl hydrocarbon receptor-induced signals up-regulate IL-22 production and inhibit inflammation in the gastrointestinal tract. *Gastroenterology.* (2011) 141:237–48.e1. doi: 10.1053/j.gastro.2011.04.007
- Rubino SJ, Geddes K, Girardin SE. Innate IL-17 and IL-22 responses to enteric bacterial pathogens. *Trends Immunol.* (2012) 33:112–8. doi: 10.1016/j.IT.2012.01.003
- Gronke K, Hernández PP, Zimmermann J, Klose CSN, Kofoed-Branzk M, Guendel F, et al. Interleukin-22 protects intestinal stem cells against genotoxic stress. *Nature.* (2019) 566:249–53. doi: 10.1038/s41586-019-0899-7
- Fatkhullina AR, Peshkova IO, Dzutsev A, Aghayev T, McCulloch JA, Thovarai V, et al. An interleukin-23-interleukin-22 axis regulates intestinal microbial homeostasis to protect from diet-induced atherosclerosis. *Immunity.* (2018) 49:943–57.e9. doi: 10.1016/j.immuni.2018.09.011
- DePaolo RW, Abadie V, Tang F, Fehlner-Peach H, Hall JA, Wang W, et al. Co-adjuvant effects of retinoic acid and IL-15 induce inflammatory immunity to dietary antigens. *Nature.* (2011) 471:220–4. doi: 10.1038/nature09849
- Geboes L, Dumoutier L, Kelchtermans H, Schurgers E, Mitera T, Renaud J-C, et al. Proinflammatory role of the Th17 cytokine interleukin-22 in collagen-induced arthritis in C57BL/6 mice. *Arthritis Rheum.* (2009) 60:390–5. doi: 10.1002/art.24220
- Sonnenberg GF, Nair MG, Kirn TJ, Zaph C, Fouser LA, Artis D. Pathological versus protective functions of IL-22 in airway inflammation are regulated by IL-17A. *J Exp Med.* (2010) 207:1293–305. doi: 10.1084/jem.20092054
- Zheng Y, Danilenko DM, Valdez P, Kasman I, Eastham-Anderson J, Wu J, et al. Interleukin-22, a TH17 cytokine, mediates IL-23-induced dermal inflammation and acanthosis. *Nature.* (2007) 445:648–51. doi: 10.1038/nature05505
- Šahmatova L, Sügis E, Šunina M, Hermann H, Prans E, Pihlap M, et al. Signs of innate immune activation and premature immunosenescence in psoriasis patients. *Sci Rep.* (2017) 7:7553. doi: 10.1038/s41598-017-07975-2
- Liang SC, Tan X-Y, Luxenberg DP, Karim R, Dunussi-Joannopoulos K, Collins M, et al. Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J Exp Med.* (2006) 203:2271–9. doi: 10.1084/jem.20061308
- Basu R, O'Quinn DB, Silberger DJ, Schoeb TR, Fouser L, Ouyang W, et al. Th22 cells are an important source of IL-22 for host protection against enteropathogenic bacteria. *Immunity.* (2012) 37:1061–75. doi: 10.1016/j.immuni.2012.08.024

26. Cella M, Fuchs A, Vermi W, Facchetti F, Otero K, Lennerz JKM, et al. A human natural killer cell subset provides an innate source of IL-22 for mucosal immunity. *Nature*. (2009) 457:722–5. doi: 10.1038/nature07537
27. Ness-Schwickerath KJ, Morita CT. Regulation and function of IL-17A- and IL-22-producing $\gamma\delta$ T cells. *Cell Mol Life Sci*. (2011) 68:2371–90. doi: 10.1007/s00018-011-0700-z
28. Gibbs A, Leeansyah E, Introini A, Paquin-Proulx D, Hasselrot K, Andersson E, et al. MAIT cells reside in the female genital mucosa and are biased towards IL-17 and IL-22 production in response to bacterial stimulation. *Mucosal Immunol*. (2017) 10:35–45. doi: 10.1038/mi.2016.30
29. Leeansyah E, Loh L, Nixon DF, Sandberg JK. Acquisition of innate-like microbial reactivity in mucosal tissues during human fetal MAIT-cell development. *Nat Commun*. (2014) 5:3143. doi: 10.1038/ncomms4143
30. Witte F, Witte K, Warszawska K, Sabat R, Wolk K. Interleukin-22: a cytokine produced by T, NK and NKT cell subsets, with importance in the innate immune defense and tissue protection. *Cytokine Growth Factor Rev*. (2010) 21:365–79. doi: 10.1016/j.cytogfr.2010.08.002
31. Paget C, Ivanov S, Fontaine J, Renneson J, Blanc F, Pichavant M, et al. Interleukin-22 is produced by invariant natural killer T lymphocytes during influenza A virus infection. *J Biol Chem*. (2012) 287:8816–29. doi: 10.1074/jbc.M111.304758
32. Kärner J, Meager A, Laan M, Maslovskaja J, Pihlap M, Remm A, et al. Anti-cytokine autoantibodies suggest pathogenetic links with autoimmune regulator deficiency in humans and mice. *Clin Exp Immunol*. (2013) 171:263–72. doi: 10.1111/cei.12024
33. Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, et al. g:Profiler: a web server for functional enrichment analysis and conversions of gene lists. (2019 update). *Nucleic Acids Res*. (2019) 47:W191–8. doi: 10.1093/nar/gkz369
34. Zhang J, Hung G-C, Nagamine K, Li B, Tsai S, Lo S-C. Development of candida-specific real-time PCR assays for the detection and identification of eight medically important candida species. *Microbiol Insights*. (2016) 9:21–8. doi: 10.4137/MBI.S38517
35. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-throughput microbial community analysis on the illumina HiSeq and MiSeq platforms. *ISME J*. (2012) 6:1621–4. doi: 10.1038/ismej.2012.8
36. Tuovinen H, Prntynen N, Gylling M, Kekäläinen E, Perheentupa J, Miettinen A, et al. $\gamma\delta$ T cells develop independently of Aire. *Cell Immunol*. (2009) 257:5–12. doi: 10.1016/j.cellimm.2009.01.012
37. Kisand K, Link M, Wolff ASB, Meager A, Tserel L, Org T, et al. Interferon autoantibodies associated with AIRE deficiency decrease the expression of IFN-stimulated genes. *Blood*. (2008) 112:2657–66. doi: 10.1182/blood-2008-03-144634
38. Rimel JK, Taatjes DJ. The essential and multifunctional TFIIH complex. *Protein Sci*. (2018) 27:1018–37. doi: 10.1002/pro.3424
39. Veldhoen M, Hirota K, Westendorf AM, Buer J, Dumoutier L, Renaud J-C, et al. The aryl hydrocarbon receptor links TH17-cell-mediated autoimmunity to environmental toxins. *Nature*. (2008) 453:106–9. doi: 10.1038/nature06881
40. Kimura A, Naka T, Nohara K, Fujii-Kuriyama Y, Kishimoto T. Aryl hydrocarbon receptor regulates Stat1 activation and participates in the development of Th17 cells. *Proc Natl Acad Sci USA*. (2008) 105:9721–6. doi: 10.1073/pnas.0804231105
41. Kisand K, Lilic D, Casanova J-L, Peterson P, Meager A, Willcox N. Mucocutaneous candidiasis and autoimmunity against cytokines in APECED and thymoma patients: clinical and pathogenetic implications. *Eur J Immunol*. (2011) 41:1517–27. doi: 10.1002/eji.201041253
42. Hetemäki I, Jarva H, Kluger N, Baldauf H-M, Laakso S, Bratland E, et al. Anticommensal responses are associated with regulatory T cell defect in autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy patients. *J Immunol*. (2016) 196:2955–64. doi: 10.4049/jimmunol.1500301
43. Fujikado N, Mann AO, Bansal K, Romito KR, Ferre EMN, Rosenzweig SD, et al. Aire inhibits the generation of a perinatal population of interleukin-17A-producing $\gamma\delta$ T cells to promote immunologic tolerance. *Immunity*. (2016) 45:999–1012. doi: 10.1016/j.immuni.2016.10.023
44. Pitt LA, Hubert F-X, Scott HS, Godfrey DI, Berzins SP. NKT cell development in the absence of the autoimmune regulator gene (Aire). *Eur J Immunol*. (2008) 38:2689–96. doi: 10.1002/eji.200838553
45. Lindh E, Rosmaraki E, Berg L, Brauner H, Karlsson MCI, Peltonen L, et al. AIRE deficiency leads to impaired iNKT cell development. *J Autoimmun*. (2010) 34:66–72. doi: 10.1016/j.jaut.2009.07.002
46. Sobkowiak MJ, Davanian H, Heymann R, Gibbs A, Emgård J, Dias J, et al. Tissue-resident MAIT cell populations in human oral mucosa exhibit an activated profile and produce IL-17. *Eur J Immunol*. (2019) 49:133–43. doi: 10.1002/eji.201847759
47. Legoux F, Bellet D, Daviaud C, El Morr Y, Darbois A, Niort K, et al. Microbial metabolites control the thymic development of mucosal-associated invariant T cells. *Science*. (2019) 366:494–9. doi: 10.1126/science.aaw2719
48. Parks OB, Pociask DA, Hodzic Z, Kolls JK, Good M. Interleukin-22 signaling in the regulation of intestinal health and disease. *Front Cell Dev Biol*. (2015) 3:85. doi: 10.3389/fcell.2015.00085
49. Mann ER, Bernardo D, Ng SC, Rigby RJ, Al-Hassi HO, Landy J, et al. Human Gut dendritic cells drive aberrant gut-specific T-cell responses in ulcerative colitis, characterized by increased IL-4 production and loss of IL-22 and IFN γ . *Inflamm Bowel Dis*. (2014) 20:2299–307. doi: 10.1097/MIB.0000000000000223
50. Zindl CL, Lai J-F, Lee YK, Maynard CL, Harbour SN, Ouyang W, et al. IL-22-producing neutrophils contribute to antimicrobial defense and restitution of colonic epithelial integrity during colitis. *Proc Natl Acad Sci USA*. (2013) 110:12768–73. doi: 10.1073/pnas.1300318110
51. Neil JA, Matsuzawa-Ishimoto Y, Kernbauer-Hfzl E, Schuster SL, Sota S, Venzon M, et al. IFN-I and IL-22 mediate protective effects of intestinal viral infection. *Nat Microbiol*. (2019) 4:1737–49. doi: 10.1038/s41564-019-0470-1
52. Oftedal BE, Marthinussen MC, Erichsen MM, Tveitarås MK, Kjellesvik-Kristiansen A, Hammenfors D, et al. Impaired salivary gland activity in patients with autoimmune polyendocrine syndrome type I. *Autoimmunity*. (2017) 50:211–22. doi: 10.1080/08916934.2017.1344972
53. Dobeš J, Neuwirth A, Dobešová M, Vobošil M, Balounová J, Ballek O, et al. Gastrointestinal autoimmunity associated with loss of central tolerance to enteric α -defensins. *Gastroenterology*. (2015) 149:139–50. doi: 10.1053/j.gastro.2015.05.009
54. Burbelo PD, Ferré EMN, Chaturvedi A, Chiorini JA, Alevizos I, Lionakis MS, et al. Profiling autoantibodies against salivary proteins in sicca conditions. *J Dent Res*. (2019) 98:772–8. doi: 10.1177/0022034519850564
55. Fishman D, Kisand K, Hertel C, Rothe M, Remm A, Pihlap M, et al. Autoantibody repertoire in APECED patients targets two distinct subgroups of proteins. *Front Immunol*. (2017) 8:976. doi: 10.3389/fimmu.2017.00976
56. Bessman NJ, Sonnenberg GF. Emerging roles for antigen presentation in establishing host-microbiome symbiosis. *Immunol Rev*. (2016) 272:139–50. doi: 10.1111/imr.12425
57. Bruslerud Ø, Siddiqui H, Marthinussen MC, Chen T, Jonsson R, Oftedal BE, et al. Oral microbiota in autoimmune polyendocrine syndrome type I. *J Oral Microbiol*. (2018) 10:1442986. doi: 10.1080/20002297.2018.1442986
58. Rautemaa R, Hietanen J, Niissalo S, Pirinen S, Perheentupa J. Oral and oesophageal squamous cell carcinoma—a complication or component of autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED, APS-I). *Oral Oncol*. (2007) 43:607–13. doi: 10.1016/j.oraloncology.2006.07.005

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Kaleviste, Rühlemann, Kärner, Haljasmägi, Tserel, Org, Trebušak Podkrajšek, Battelino, Bang, Franke, Peterson and Kisand. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.