

Mendelian randomisation shows diverticular disease and irritable bowel syndrome increase the risk of haemorrhoidal disease

Recently in *Gut*, Zhu and colleagues¹ investigated causality of genetic risk effects across three genetically correlated conditions: haemorrhoidal disease (HEM), diverticular disease (DIV) and IBS. Using genome-wide association study (GWAS) summary statistics from our previous HEM GWAS² and other studies,^{3–5} they conducted unidirectional two-sample Mendelian randomisation (MR) and concluded ‘HEM might not be a trigger for developing diverticular disease or IBS, providing novel insights into a modest genetic correlation in conditions. Simultaneously, the observed causal (protective, ed.) effects of HEM on diverticular disease and IBS requires further exploration’. Although we did not originally make any claims of causality from the observed genetic correlations, the findings of Zhu *et al* are surprising and not in line with previous results,² hence we set out to investigate further the relationship between HEM, DIV and IBS. Here, we briefly report an analysis that differs from Zhu *et al* in that: (1) we used best practice two-sample MR analysis,⁶ (2) causality was studied bi-directionally (testing HEM both as exposure (cause) and outcome (effect) versus DIV and IBS risk) and (3) summary statistics from the largest, recent GWAS meta-analysis including >53 000 IBS cases were used to test genetic risk effects of HEM relative to IBS (online supplemental material).⁷ By these means, we obtained results different from Zhu *et al* and new evidence of causative effects of DIV and (to a lesser extent) IBS towards the risk of HEM. Figure 1, online supplemental figure S1 and S2 illustrate the outcome of our analyses, outlined below.

We first optimised the instrument variable (IV) selection for MR by using consensus thresholds and methods⁶ to select genetic variants that are: associated with the exposure ($p < 5 \times 10^{-8}$ or $p < 5 \times 10^{-5}$); not in linkage disequilibrium (LD) ($r^2 < 0.001$ across a 1 Mb window); and not horizontally pleiotropic (having no direct association with the outcome) ($p_{\text{MR-PRESSO}} < 0.05$). Then, using the selected variants as IVs, we performed bi-directional two-sample MR analyses^{8,9} using inverse variance weighted (IVW), MR Egger, weighted median, simple mode,




Figure 1 Bi-directional Mendelian randomisation (MR) analyses indicate a causal relationship between the effects of SNPs associated with DIV and IBS, respectively, on the development of HEM. The graph illustrates the associations between (A) IBS and HEM as well as (B) DIV and HEM, with ORs and their corresponding 95% CIs shown on the x-axis. The y-axis represents HEM as either an exposure or an outcome variable. MR analyses were conducted by using inverse variance weighted (IVW), MR Egger, weighted median, simple mode, weighted mode and CAUSE methods. Significance levels: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; else not significant. DIV, diverticular disease; HEM, haemorrhoidal disease.

weighted mode and CAUSE methods. Additionally, to ensure the robustness of the causal effects, we performed leave-one-out IVW regression analysis, which shows that causal effects are not driven by a few outlying genetic variants (online supplemental material).

Hence, this pipeline was first applied to testing HEM as exposure towards the risk of developing DIV or IBS (outcome) as in Zhu *et al*. In contrast to Zhu *et al*, no significant findings were obtained in our analysis in this direction (see HEM exposure in figure 1). However, when either DIV or IBS were tested as exposure on HEM as outcome, significant observations were made using multiple MR models (see HEM outcome in figure 1). In particular, MR based on the IVW method resulted in strongest and most significant findings, both for IBS and for DIV showing predisposing effects on HEM. Remarkably, DIV was associated with nearly five times increased HEM risk ($\text{OR} = 4.8$; $p = 2 \times 10^{-4}$). While their results were limited to testing HEM as exposure, discrepancies with Zhu *et al* are most likely due to the fact that they used IVs variants that are in LD and therefore not independent (as required for MR analyses; a detailed analysis of methodological issues is reported in the online supplemental material).

In summary, we show that DIV and IBS significantly predispose to HEM. This is in line with real-world data, as a most recent colonoscopy study also demonstrated: not only were patients with DIV more likely to have internal HEM ($\text{OR} = 2.21$) but also those with more than 10 diverticula had the highest risk of HEM ($\text{OR} = 3.33$).¹⁰ Indeed, DIV was deemed by the authors

even more important than diet or constipation for internal HEM development.

Simonas Juzenas^{1,2}, **David Ellinghaus**^{3,2}, **Serena Sanna**^{3,4}, **Andre Franke**^{2,5}, **Mauro D'Amato**^{6,7,8}

¹Institute of Biotechnology, Life Science Centre, Vilnius University, Vilnius, Lithuania

²Institute of Clinical Molecular Biology, Kiel University, Kiel, Germany

³Institute for Genetic and Biomedical Research (IGBR), Consiglio Nazionale delle Ricerche (CNR), Cagliari, Italy

⁴University of Groningen, Department of Genetics, University Medical Center Groningen, Groningen, The Netherlands

⁵University Hospital Schleswig-Holstein (UKSH), Kiel, Germany

⁶Department of Medicine and Surgery, LUM University, Casamassima, Italy

⁷Ikerbasque, Basque Foundation for Science, Bilbao, Spain

⁸Gastrointestinal Genetics Lab, CIC bioGUNE – BRTA, Derio, Spain

Correspondence to Professor Mauro D'Amato, Department of Medicine and Surgery, Università LUM Giuseppe Degennaro, Casamassima, Puglia, Italy; damato@lum.it

Twitter Simonas Juzenas @SimonasJuzenas and Mauro D'Amato @damato_mauro

Contributors SS, AF and MD designed and supervised the study; SJ and DE contributed to data acquisition, statistical and computational analyses; SJ, DE, SS, AF and MD analysed and interpreted data; SJ and MD drafted the manuscript, with input and critical revision from all other authors. All authors approved the final draft of the manuscript.

Funding The study received funds through AF's grantFR 2821/19-1 from the German Research Foundation (DFG). SS is supported by Next Generation EU grant Project Age-It (DM MUR 1557 11.10.2022), ERC Starting Grant 2022 (101075624) and NutrAGE grant (DM MUR 844 16.07.2021). SJ is supported by MSCA-IF-2020 grant (101030265). The study received infrastructure support from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Cluster of Excellence 2167 'Precision Medicine in Chronic Inflammation (PMI)' (EXC 2167-390884018).

Competing interests MD has received unrestricted research grants and consulting fees from QOL Medical LLC.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.



OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly

cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2023-330364>).



Check for updates

To cite Juzenas S, Ellinghaus D, Sanna S, *et al.* *Gut* Epub ahead of print: [please include Day Month Year]. doi:10.1136/gutjnl-2023-330364

Received 26 May 2023

Accepted 20 September 2023

Gut 2023;0:1–2. doi:10.1136/gutjnl-2023-330364

ORCID iDs

Simonas Juzenas <http://orcid.org/0000-0001-9293-0691>

David Ellinghaus <http://orcid.org/0000-0002-4332-6110>

Mauro D'Amato <http://orcid.org/0000-0003-2743-5197>

REFERENCES

- 1 Zhu Z, Chen X, Wang C, *et al.* Haemorrhoidal disease reduces the risk of Diverticular disease and irritable bowel syndrome: a Mendelian Randomisation study. *Gut* 2023;gutjnl-2022-329307.
- 2 Zheng T, Ellinghaus D, Juzenas S, *et al.* Genome-wide analysis of 944 133 individuals provides insights into the etiology of Haemorrhoidal disease. *Gut* 2021;70:1538–49.
- 3 Schafmayer C, Harrison JW, Buch S, *et al.* Genome-wide Association analysis of Diverticular disease points towards neuromuscular, connective tissue and epithelial Pathomechanisms. *Gut* 2019;68:854–65.
- 4 Dönertaş HM, Fabian DK, Valenzuela MF, *et al.* Common genetic associations between age-related diseases. *Nat Aging* 2021;1:400–12.
- 5 Jiang L, Zheng Z, Fang H, *et al.* A generalized linear mixed model Association tool for Biobank-scale data. *Nat Genet* 2021;53:1616–21.
- 6 Burgess S, Davey Smith G, Davies NM, *et al.* Guidelines for performing Mendelian randomization investigations. *Wellcome Open Res* 2020;4:186.
- 7 Eijbsbouts C, Zheng T, Kennedy NA, *et al.* Genome-wide analysis of 53,400 people with irritable bowel syndrome highlights shared genetic pathways with mood and anxiety disorders. *Nat Genet* 2021;53:1543–52.
- 8 Hemani G, Zheng J, Elsworth B, *et al.* The MR-base platform supports systematic causal inference across the human Phenome. *Elife* 2018;7:e34408.
- 9 Morrison J, Knoblauch N, Marcus JH, *et al.* Mendelian randomization accounting for correlated and Uncorrelated pleiotropic effects using genome-wide summary Statistics. *Nat Genet* 2020;52:740–7.
- 10 Sninsky JA, Galanko J, Sandler RS, *et al.* Diverticulosis is associated with internal hemorrhoids on colonoscopy: possible clues to etiology. *Clin Gastroenterol Hepatol* 2023;21:1097–9.